

1899

**T**ABLET TRITURATES, HYPO-  
DERMIC AND COMPRESS-  
ED TABLETS, CHOCOLATE  
COATED TABLETS, PHYSI-  
CIANS' PRIVATE FORMU-  
LAS, AND VIAL CASES.



NEW YORK  
THE FRASER TABLET  
TRITURATE MFG. COMPANY  
262, Fifth Avenue

*Manufacturing Department: 454 to 474 Eighteenth St.  
Brooklyn, New York*

*Where all orders should be addressed*

## TERMS OF SALE.

---

1. The prices on our list are for thousands, unless the price of 100 is given. Tablet Triturates are put up one thousand in a 4-ounce, French square, glass stoppered bottle. Compressed Tablets in thousands are put up in bottles ranging from 4 to 32 ounces, according to the size of tablet.

2. For five hundred Tablets, one-half the rate per thousand is charged, and five cents per 500 additional.

3. For one hundred Tablets, where the price for 100 is not given, the price is one-tenth of the price per 1000 and ten cents per 100 additional. The more expensive have the price per 100 attached.

4. Tablets sold in bulk, in quantities of five thousand or more in one container, will be charged for at five cents per thousand less than list prices, for those listed at less than one dollar. Those listed at more than one dollar will be charged for at five per cent less than list prices.

5. For those Chocolate Coated tablets which we manufacture, the price will be ten cents per thousand more than the list prices, for all those listed for less than two dollars per thousand. Tablets listed at two dollars or over, will be sold for the same price coated, excepting where the prices for coated tablets are given in the price list.

6. The tablets are always shipped in bottles, unless otherwise specified, and in any other container are at the risk of the customer and are not returnable.

7. The goods are invariably sold for cash, and no deviation from the list price will be made, except for marked fluctuations in the market price of the goods. The prices for tablets containing quinine are subject to change without notice.

8. We do not pay freight, mail or express charges, excepting on orders amounting to fifteen dollars or more, when sold at net list price.

9. We do not consign any goods.

10. As the tablets on our list have all been properly tested for quality, it must not be expected that they will be accepted back, either for credit or exchange, after they are once sold, unless for a just complaint of that quality.

11. When ordering, please give full address and shipping directions if preference should be given to any particular line or route.

NATIONAL LIBRARY OF MEDICINE  
Bethesda, Maryland

*Gift of*  
William H. Helfand



18-

# FRASER'S

## SOLUBLE

# Tablet Triturates

(As Suggested by DR. ROBT. M. FULLER.)

—AND—

## Compressed Tablets.

THESE TABLETS, INCLUDING HYPODERMICS, ETC.,  
GIVE TO THE PHYSICIAN AND DRUGGIST AN EASY,  
ECONOMICAL AND ACCURATE METHOD OF  
DISPENSING MEDICINES IN A COMPACT  
AND PALATABLE FORM.

—♦♦♦—

THE directions appended to the formulas in the following list represent the individual views of the authors of the prescriptions, or have been condensed from the authorities on therapeutics. We adopt this method of replying to the many inquiries received every day from physicians as to the composition, special uses and indications of certain tablets, and in so doing assume no responsibility other than that which attaches to a desire to afford them the assistance of such information as we gather from correspondence or conversation with their fellow-practitioners in all parts of the country.

When not otherwise stated, the dose is intended for an adult, and the correction for age may be made according to the usual tables. Where the same directions are used for tablets of two or more strengths of the same medicine, it indicates the conclusions or opinions of physicians of different experience or different classes, as to the unit of dosage to be employed in specific cases, or the variation which would naturally be governed by the circumstances of each case.

When a patient cannot swallow a tablet whole, a single one may be dissolved in a proper amount of water, or a number may be dissolved in any multiple of the amount, and the directions given as for a liquid medicine, except in the cases where the tablet contains insoluble elements. For example:—One tablet in a tablespoonful of water at a dose, or eight in eight tablespoonfuls, and of this a tablespoonful as often as necessary.

Suggestions for the use of any of our tablets or for the improvement of any directions that seem inadequate, will be gratefully received. Our endeavor is to make the list as complete as we can, and as useful as possible to physicians; and while we know that the first attempt of this kind may leave much to be desired, we feel sure that it will be helpful and suggestive, and we request the aid of the medical profession in our effort to make it of even more assistance.

## Per Thousand.

<b>ABSORBENT DYSPEPTIC</b> .....	Pepsin.....1 gr. Charcoal.....2 gr. Sodium Bicarb..2 1-2 gr.	\$1 85
One or two after meals, followed by another in one hour, for dyspepsia and acid stomach.		
<b>ACETANILID</b> .....	1 gr.....	75
Antipyretic analgesic.		
Give one every two hours, for fever, with pain and restlessness, in children; one every twenty minutes for adults until eight are taken, if necessary.		
<b>ACETANILID</b> .....	2 grs.....	1 00
Antipyretic.		
One every half hour or every hour until three or four are taken, for fever with pain.		
<b>ACETANILID</b> .....	3 grs.....	1 20
Give one every two hours as an antipyretic until the temperature falls; also, in the same dose, to control pain in neuralgia and muscular rheumatism.		
<b>ACETANILID</b> .....	4 grs.....	1 35
One, repeated in two hours, followed by another if necessary, for neuralgia, headache, and to control fever.		
<b>ACETANILID</b> .....	5 grs.....	1 50
One, repeated in two hours, followed by another if necessary, for neuralgia, headache, and to control fever.		
<b>ACETANILID ALKALINE</b> .....	Acetanilid.....3 gr. Spts. Ammon. Aro. 5 min.	2 00
One every hour for three doses, to control the fever and pain of polyarthritis; then every three hours.		
<b>ACETANILID AROMATIC</b> .....	Acetanilid.....5 gr. Oil Gaultheria.....q. s. Pink Coloring.....q. s.	1 65
Antipyretic.		
Give one every two hours as an antipyretic in typhoid, bronchitis and pneumonia; after the temperature falls, two to four a day.		
<b>ACETANILID, CANNABIS CO.</b> (Dr. H. C. Whistler.)	Acetanilid.....1 1-2 gr. Ext. Cannab. Ind..1-6 gr. Caffeine Alk.....1 gr.	3 00
One every hour until three are taken; afterwards one every two hours, in all headaches when face is flushed and eyes congested, or congestive headaches; in neuralgia of branches of fifth nerve.		
<b>ACETANILID COMP. No. 1</b> ....	Acetanilid.....2 gr. Fl. Ext. Gelsem....1 min.	1 30
(Dr. Pitcher.)		
Antipyretic, analgesic.		
One every two hours for three doses, then every two hours if necessary, for neuralgia and headache, and also fever with pain.		
<b>ACETANILID COMP. No. 3</b> ....	Acetanilid.....3 gr. Tully's Powder....3 gr.	2 15
Diaphoretic, antipyretic.		
One every two to four hours, for the fever of pneumonia, bronchitis and typhoid; and as an analgesic in rheumatism.		
<b>ACETANILID COMP. No. 5</b> ....	Acetanilid.....2 gr. Quinine Sulph.....1 gr.	2 50
(Dr. Morgan.)		
One every hour for four doses, then one every two hours until relieved. For influenza and fresh colds with severe pains in the head and body.		

<b>ACETANILID COMP No. 6</b> ..... (Dr. F. Preiss.)	{ Acetanilid.....5 gr. }	2 50
	{ Camphor Monobr...1 gr. }	
	{ Caffeine Cit.....2-3 gr. }	
	{ Capsicum Powd...1-20 gr. }	
	{ Tr. Gelsemium....2 min. }	
	{ Oil Wintergreen....q. s. }	

One tablet, repeated every hour for three doses, if necessary; for headache and neuralgia.

<b>ACETANILID &amp; CAFFEINE</b> <b>COMP</b> ..... (Dr. Hubbard.)	{ Acetanilid.....3 1-2 gr. }	4 50
	{ Caffeine .....1 gr. }	
	{ Sodium Brom....7 1-2 gr. }	

Give two tablets for the first dose, and repeat in four hours if necessary for headache and neuralgia.

<b>ACETANILID &amp; CALOMEL</b> <b>COMP</b> ..... (Dr. J. C. Gary.)	{ Acetanilid.....1 gr. }	1 00
	{ Sodium Bicarb....1-2 gr. }	
	{ Calomel.....1-10 gr. }	

Give a child from three to six years old a tablet every hour, for from three to six doses, or until bowels act. If they do not act in six hours, a saline cathartic or an enema is indicated, then give a tablet every two to four hours. For eruptive and irritative fevers, and typhoid in the first week; or in suspected cases.

In chronic cases, one every three to six hours for children or adults.

<b>ACETANILID &amp; GELSEMIN</b> <b>COMP</b> .....	{ Acetanilid.....2 1-2 gr. }	2 60
	{ Gelsemin.....1-10 gr. }	
	{ Caffeine Cit.....1 gr. }	

One every hour for not more than three doses, for headache with pain or fever.

<b>ACETANILID MODIFIED</b> .....	{ Acetanilid.....4 gr. }	1 65
	{ Tr. Digitalis.....2 min. }	

One tablet every two or three hours, for the fever and pain of pericarditis, pneumonia and bronchitis.

<b>ACETANILID SALICYLATED</b> .....	{ Acetanilid.....4 1-2 gr. }	1 60
	{ Sodium Salicyl...1-4 gr. }	
	{ Sodium Bicarb....1-4 gr. }	

One every hour for two or three doses, then every three hours, for acute articular rheumatism and its complications

<b>ACETANILID &amp; SALOL</b> ..... Antiseptic, antipyretic.	{ Acetanilid.....2 1-2 gr. }	4 00
	{ Salol.....2 1-2 gr. }	

One every hour until fever and pain abate, in acute articular rheumatism, and neuralgia of rheumatic origin.

<b>ACETANILID &amp; SODA</b> .....	{ Acetanilid.....4 gr. }	1 40
	{ Sodium Bicarb....1 gr. }	

One, repeated in two hours, followed by another if necessary, for headache and neuralgia.

<b>ACETANILID, SODA COMP</b> ... (Dr. R. H. Dingear.)	{ Acetanilid.....4 gr. }	2 35
	{ Sodium Bicarb....1 gr. }	
	{ Caffeine Citrate....1 gr. }	

One every two hours until symptoms are relieved, then increase intervals as indicated. Used in neuralgic headache and as an antipyretic.

<b>ACID ARSENOUS</b> .....1-1000 gr.....	75
--	----

Tonic alterative, antimalarial.

One tablet every two hours for an infant, as a tonic in chronic diarrhoea, and those forms of pain and headache which have their origin in malaria.

<b>ACID ARSENOUS</b> .....1-500 gr.....	75
---	----

One to three tablets every two hours, for eczema, psoriasis and other skin diseases; also for a general tonic in malaria, and those forms of headache and neuralgia of malarial origin.

<b>ACID ARSENOUS</b> .....	I-200 gr.....	75
One to two tablets every two hours, for eczema, psoriasis and other skin diseases; also for a general tonic in malaria, and those forms of headache and neuralgia of malarial origin.		
<b>ACID ARSENOUS</b> .....	I-100 gr.....	75
One tablet after each meal, increased by one tablet a day until effect is produced, or until fulness of the stomach with pain occurs; for a general and malarial tonic, skin diseases, and those forms of neuralgic pain of malarial origin.		
<b>ACID ARSENOUS</b> .....	I-60 gr.....	75
One to three tablets after each meal, stop when there are colicky pains or fulness in the stomach; for skin diseases, neuralgia, epilepsy and for a malarial and general tonic.		
<b>ACID ARSENOUS</b> .....	I-50 gr.....	75
One to two tablets after each meal, stop when there are colicky pains or fulness in the stomach; for skin diseases, neuralgia, epilepsy and for a malarial and general tonic.		
<b>ACID ARSENOUS</b> .....	I-40 gr.....	75
One after each meal, increased very gradually to two after each meal, for tonic in anæmia.		
<b>ACID ARSENOUS</b> .....	I-30 gr.....	75
One tablet after each meal, stop when there are colicky pains or fulness in the stomach; for skin diseases, neuralgia, epilepsy and for a malarial and general tonic.		
<b>ACID ARSENOUS</b> .....	I-20 gr.....	75
One three times a day after meals, for tonic in anæmia.		
<b>ACID ARSENOUS</b> .....	I-10 gr.....	75
One tablet after each meal, given with care; stop when there are colicky pains or fulness in the stomach; for skin diseases, neuralgia, epilepsy and for a malarial and general tonic.		
<b>ACID BENZOIC</b> .....	I-5 gr.....	75
One every half hour in summer diarrhœa, of infants; for aphthous and diphtheritic affections of the mouth, three or four an hour, dissolved in a teaspoonful of water.		
<b>ACID BENZOIC</b> .....	I gr.....	I 10
A tablet every hour for diseases of the respiratory passages, erysipelas, diphtheria, typhoid fever and acute rheumatism.		
<b>ACID BORIC</b> .....	I-10 gr.....	75
Antiseptic.		
Dissolve ten in half a glass of water, and give a teaspoonful every fifteen to thirty minutes, for an infant with diarrhœa or dysentery, and flatulence.		
<b>ACID BORIC</b> .....	I-2 gr.....	75
A tablet after each meal, increased to five or six at a dose, for catarrh of the ureters and bladder, and the uric acid diathesis.		
<b>ACID BORIC</b> .....	I gr.....	95
One every hour, dissolved on the tongue; for aphthæ and diphtheria, and for diarrhœa of children.		
<b>ACID BORIC</b> .....	3 grs.....	I 15
One every two hours for dysmenorrhœa, and vesical catarrh; three or four times a day as an intestinal disinfectant.		
<b>ACID BORIC</b> .....	5 grs.....	I 30
A tablet three or four times a day, for intestinal dyspepsia with diarrhœa, and in catarrhal inflammations of the urinary tract.		

<b>ACID CARBOLIC</b> .....	1-10 gr.....	75
One every hour for indigestion with nausea, and as an intestinal disinfectant in diarrhoea, dysentery and typhoid fever.		
<b>ACID SALICYLIC</b> .....	1 gr.....	1 10
Anti-rheumatic, antipyretic.		
One every hour for three doses, then every two or three hours, for acute rheumatism of children; discontinue the medication if nausea and vomiting occur.		
<b>ACID SALICYLIC</b> .....	3 grs.....	1 50
One tablet every hour for three doses, then every three hours; for acute rheumatism and its complications, neuralgia, tonsillitis and pleurisy.		
<b>ACID SALICYLIC</b> .....	5 grs.....	1 75
A tablet three or four times a day as an antipyretic; alternated with quinine to great advantage; used also in neuralgic affections, typhoid and hectic fever.		
<b>ACONITIA</b> .....	1-1000 gr.....	75
<b>ACONITIA</b> .....	1-500 gr.....	75
<b>ACONITIA</b> .....	1-400 gr.....	1 00
<b>ACONITIA</b> .....	1-200 gr.....	2 00
<b>ACONITIA</b> .....	1-150 gr.....	3 00
<b>ACONITIA</b> .....	1-100 gr.....	4 00
<b>ACONITIA</b> .....	1-50 gr.....	6 00
<b>ACONITE COMP</b> .....	{ Morph. Sulph....1-50 gr. Tartar Emetic..1-100 gr. Ext. Aconite....1-100 gr. }	1 25
One every three hours, for a child with bronchitis and high fever.		
<b>ACONITE COMP. No. 2</b> .....	{ Tr. Aconite.....1 min. Tr. Belladonna....2 min. Tr. Bryonia.....2 min. }	1 50
(Dr. W. E. Derry.)		
One every hour for three to six doses, then every two to three hours, in acute inflammation of the air passages, pharyngitis and tonsillitis.		
<b>ACONITE FL. EXT</b> .....	1 minim.....	75
Diaphoretic.		
Dissolve one in six tablespoonfuls of water; of this give one tablespoonful every hour, in neuralgia and fever.		
<b>ACONITE &amp; BELLADONNA</b> ..	{ Aconite Powd..1-100 gr. Belladon. Powd..1-100 gr. }	75
One crushed and taken on the tongue, or mixed with water on the end of a spoon, to a baby under three, at intervals of an hour, for bronchitis, pleurisy and fevers; for older children, one every half hour for four doses, then every hour.		
<b>ACONITE &amp; BELLADONNA</b> {	Tr. Aconite,....1-2 min. }	75
<b>No. 2</b> .....	Tr. Belladonna..1-2 min. }	
One every hour until effect is produced. For children, dissolve one tablet in ten teaspoonfuls of water, and give a teaspoonful every twenty minutes, for fever, tonsillitis and early stages of inflammatory diseases.		
<b>ACONITE &amp; BELLADONNA</b> {	Tr. Aconite,....1 min. }	1 00
<b>No. 3</b> .....	Tr. Belladonna...1 min. }	
One every hour for measles and other fevers of children, acute coughs and colds and acute rheumatism; after the sixth dose, give every three hours.		
<b>ACONITE &amp; BRYONIA</b> .....	{ Aconite Powd..1-100 gr. Bryonia Powd..1-100 gr. }	75
Sedative, antipyretic, membrane stimulant.		
One every half hour as a sedative and antipyretic in the early stages of fever, and cold with fever.		

<b>ACONITE &amp; BRYONIA, No. 2.</b>		{ Tr. Aconite.....1 min. Tr. Bryonia .....1 min. }	1 00
One, three or four times a day, increasing carefully according to the effect; for pneumonia and for pleurisy with effusion.			
<b>ACONITE &amp; BRYONIA COMP.</b> (Dr. Judd.)		{ Tr. Aconite.....1 min. Tr. Gelsem.....1 7-8 min. Tr. Bryonia....1 1-4 min. }	1 35
Dissolve ten tablets in four ounces of water and give a teaspoonful every hour in all febrile conditions, and especially in those due to inflammation of the air-passages.			
<b>ACONITE &amp; GELSEMIUM.....</b>		{ Tr. Aconite.....2 min. Tr. Gelsemium....1 min. Tr. Belladonna....2 min. }	1 50
Give one in water, for sick headache and neuralgia; repeat hourly for two doses if necessary.			
<b>ACONITE &amp; IPECAC.....</b>		{ Tr. Aconite.....1-2 min. Sedative, diaphoretic, expectorant. { Wine of Ipecac...1 min. }	1 00
One every half hour for five doses, then one every hour or two, as a sedative and diaphoretic; for colds with hoarseness and dryness of the throat.			
<b>ACONITE &amp; MERCURIUS.....</b>		{ Aconite Powd...1-100 gr. Sedative, alterative. { Merc. Sol. H....1-100 gr. }	75
One every two hours for a child, for fever, or cold with fever and congestion.			
<b>ACONITE RAD. PULV... ..</b>		1-200 gr.....	75
Antipyretic, sedative.			
One every hour in the fevers of children, when pulse is rapid and of high tension.			
<b>ACONITE RAD. PULV.....</b>		1-100 gr.....	75
Give one tablet every hour or two, for elevated temperature, strong pulse and restlessness, and for colds of children, with fever.			
<b>ACONITE RAD. PULV.....</b>		1-10 gr.....	75
One tablet every two or three hours, for colds with fever.			
<b>ACONITE &amp; STIBIUM.....</b>		{ Aconite Powd....1-50 gr. (Dr. Wood.) { Tartar Emetic...1-50 gr. }	75
Sedative expectorant.			
One or two every half hour for four doses; repeat after two hours, in cases of dry and obstinate cough, asthma and chronic bronchitis.			
<b>AGARICIN.....</b>		1-10 gr.....	1 50
One or two at night, for the profuse sweating of phthisical and gouty patients, and for exhausting cough.			
<b>AIKEN'S TONIC.....</b>		{ Quinine Sulph.....1 gr. Iron Powd.....2-3 gr. Acid Arsenous...1-50 gr. Strych. Sulph....1-50 gr. Ext. Gentian.....1-4 gr. }	
		Plain or Chocolate Coated.....	2 00
One to two tablets, three times a day, after meals.			
<b>ALKALINE CHARCOAL.....</b>		{ Charcoal.....1 gr. Absorbent alkaline. { Sodium Bicarb....1 gr. }	75
Give frequently with a little water, for heartburn, acidity of the stomach and flatulence.			

<b>ALKALINE IODIDE</b> .....	{ Potass. Iodide.....5 gr. }	6 00
Alterative.	{ Potass. Bicarb.....5 gr. }	
One after eating, for depressed and weakened conditions of the system, poverty of the blood, and chronic syphilitic conditions.		
<b>ALKALINE LAXATIVE</b> .....	{ Ext. Belladonna...1-8 gr. }	3 00
	{ Ext. Nux Vom....1-4 gr. }	
	{ Ext. Colocy. Co..1 1-2 gr. }	
	{ Sodium Bicarb.....1 gr. }	
One every night, for chronic constipation and the intestinal dyspepsia of rheumatic and gouty subjects.		
<b>ALKALINE LAXATIVE No. 2.</b>	{ Rhubarb Powd. ....2 gr. }	1 15
(Dr. J. W. Malone.)	{ Rochelle Salt.....4 gr. }	
	{ Sodium Bicarb.....4 gr. }	
One tablet after meals.		
<b>ALOES, COLOCYNTH COMP.</b>	{ Aloes .....1 gr. }	2 50
(Dr. Duncan.)	{ Ext. Colocynth Co..2 gr. }	
	{ Ext. Belladonna...1-8 gr. }	
	<b>Plain or Chocolate Coated</b>	
One tablet before dinner and two at bedtime, followed for several days by one or two at bedtime, to relieve torpidity or passive congestion of the liver.		
<b>ALOIN</b> .....	1-10 gr.....	75
	<b>Chocolate Coated</b> .....	85
One tablet at night occasionally, for constipation of children due to errors in diet.		
<b>ALOIN</b> .....	1-5 gr.....	75
	<b>Chocolate Coated</b> .....	85
One or two at night in habitual constipation.		
<b>ALOIN</b> .....	1-4 gr.....	75
	<b>Chocolate Coated</b> .....	85
One or two at night in chronic constipation of sedentary people.		
<b>ALOIN</b> .....	1-2 gr.....	1 00
	<b>Chocolate Coated</b> .....	1 10
One to four tablets at night, for constipation.		
<b>ALOIN COMP. No. 1</b> .....	{ Aloin.....1-8 gr. }	1 10
Cathartic, hepatic.	{ Podophyllin .....1-8 gr. }	
	<b>Chocolate Coated</b> .....	1 20
One to four at night, as an hepatic stimulant in chronic constipation; gradually reduce the dose.		
<b>ALOIN COMP. No. 2</b> .....	{ Aloin .....1-4 gr. }	1 25
	{ Podophyllin .....1-4 gr. }	
	<b>Chocolate Coated</b> .....	1 35
One or two at night as a laxative, where the action of the liver is sluggish.		
<b>ALOIN &amp; BELLADONNA CO.</b>	{ Aloin.....1-5 gr. }	1 10
Tonic laxative.	{ Ext. Belladonna...1-8 gr. }	
	{ Strych. Sulph....1-60 gr. }	1 20
	<b>Chocolate Coated</b> .....	
One or two tablets at night, repeated every three hours the next day if necessary; for torpid liver and as a laxative in chronic constipation.		
<b>ALOIN &amp; BELLADONNA CO.</b>	{ Aloin.....1-5 gr. }	1 10
No. 2.....	{ Ext. Belladonna...1-8 gr. }	
	{ Strych. Sulph....1-120 gr. }	1 20
	<b>Chocolate Coated</b> .....	
One or two at night, to be repeated the next morning if necessary; as a tonic laxative.		

ALOID, BELLADONNA AND NUX..... (Dr. Willard.)	Aloid.....	1-5 gr.	1 25
	Ext. Belladonna..	1-8 gr.	
	Ext. Nux Vomica	1-6 gr.	
	Chocolate Coated.....		1 35

One every night, decreasing to one every second night, in cases of habitual constipation.

ALOID, BELLADONNA AND PODOPHYLLIN..... (Dr. LeMoyne.)	Aloid.....	1-8 gr.	1 25
	Ext. Belladonna..	1-8 gr.	
	Podophyllin .....	1-8 gr.	
Hepatic, tonic laxative.	Chocolate Coated.....		1 35

ALOID, BELLADONNA AND PODOPHYLLIN COMP.....	Aloid.....	1-10 gr.	1 25
	Ext. Belladonna..	1-10 gr.	
	Podophyllin.....	1-10 gr.	
	Ext. Nux Vom..	1-10 gr.	
	Chocolate Coated.....		1 35

One or two at bedtime as an hepatic laxative, repeated every three hours the next day, as a tonic to liver; then one every second or third night for constipation.

ALOID, BELLADONNA AND STRYCH. with CASCARA..	Aloid.....	1-5 gr.	1 75
	Ext. Belladonna..	1-16 gr.	
	Strych. Sulph.....	1-60 gr.	
	Ext. Cascara.....	1-2 gr.	
	Chocolate Coated.....		1 85

One, night and morning, for constipation and as an hepatic stimulant.

ALOID, CASCARIN COMP.... (Dr. Duncan.)	Aloid.....	1-5 gr.	1 50
	Ext. Belladonna..	1-8 gr.	
	Cascarin.....	1-4 gr.	
	Strych. Sulph.....	1-60 gr.	
Tonic hepatic.	Chocolate Coated.....		1 60

ALOID, CASCARIN CO. No. 2 (Dr. Duncan.)	Aloid.....	1-4 gr.	1 60
	Podophyllin.....	1-4 gr.	
	Cascarin.....	1-4 gr.	
	Ext. Belladonna..	1-8 gr.	
	Chocolate Coated.....		1 70

ALOID, HYOSCYAMUS CO... (Dr. Wilkinson.)	Aloid.....	1-4 gr.	1 40
	Ext. Belladonna..	1-4 gr.	
	Ext. Hyoscyamus	1-4 gr.	
	Chocolate Coated.....		1 50

One or two every night, for cases of constipation with intestinal discomfort and hemorrhoids.

ALOID, IRON & STRYCHNINE (Dr. E. Jay Fisk.)	Aloid.....	1-10 gr.	1 10
	Iron Powd.....	1 gr.	
	Strych. Sulph.....	1-60 gr.	
	Chocolate Coated.....		1 20

One or two after each meal, dose to be governed by the laxative effect, for the atonic dyspepsia and non-assimilation of the anæmic; also in anæmia and general debility with constipation.

ALOID, PODOPHYLLIN AND HYOSCYAMUS..... (Dr. Clark.)	Aloid.....	1-4 gr.	1 25
	Podophyllin.....	1-4 gr.	
	Ext. Hyoscyam..	1-10 gr.	
	Chocolate Coated.....		1 35

One or two at night, followed by one next morning if necessary, for constipation.

When ordering or prescribing

ALOIN, SAGRADIN COMP. .... (Dr. Morris.)	{ Aloin.....1-8 gr. }	1 25
	{ Strych. Sulph....1-120 gr. }	
	{ Cascarin.....1-6 gr. }	
	{ Ext. Hyoscyam....1-4 gr. }	
	Chocolate Coated.....	1 35

One or two at night, followed by one every three hours the next day until catharsis is produced; afterwards one at night as necessary; in chronic constipation and any cholæmic condition where a mild but good cholagogue is needed.

ALOIN & STRYCHNINE CO.. Laxative, stomachic.	{ Aloin.....1-5 gr. }	1 10
	{ Ext. Belladonna...1-8 gr. }	
	{ Strych. Sulph....1-60 gr. }	
	{ Ipecac Powd....1-16 gr. }	
	Chocolate Coated.....	1 20

One before supper and at bedtime, followed by one every two hours next day if necessary, for constipation, sick headache and as a tonic to the liver.

ALOIN STRYCHNINE CO. No 2	{ Aloin.....1-5 gr. }	1 10
	{ Ext. Belladonna...1-16 gr. }	
	{ Strych. Sulph....1-60 gr. }	
	{ Ipecac Powd....1-16 gr. }	
	Chocolate Coated.....	1 20

One before supper and at bedtime, followed by one every two hours next day if necessary, for constipation, sick headache and as a tonic to the liver.

ALTERANS..... (Dr. Hamilton.)	{ Acid Arsenous...1-20 gr. }	75
	{ Strych. Sulph....1-30 gr. }	
	{ Merc. Bichlor....1-40 gr. }	
	{ Iron Powd.....1 gr. }	
	Chocolate Coated.....	85

A tonic in certain intractable nerve diseases, to be used under proper indications.

ALUM COMP.....	{ Alum.....2 1-2 gr. }	2 00
	{ Zinc Sulphate.....2 gr. }	
	{ Ex. Hydras.colorless 1 gr. }	
	{ Morph. Sulph....1-32 gr. }	

Dissolve one in two to four tablespoonfuls of water for use as an injection in gonorrhœa and leucorrhœa.

AMMONIUM BROMIDE.....5 grs.....	1 10
---------------------------------	------

Sedative.

One or two at night, repeated in an hour, to quiet the nerves and induce sleep.

AMMONIUM BROMIDE.....10 grs.....	2 00
----------------------------------	------

Give one tablet every two hours, repeated as many as three times, if necessary, for cerebral congestion and insomnia.

AMMONIUM MURIATE.... 1-10 gr.....	75
-----------------------------------	----

For an infant, one tablet crushed in a little water, every half hour, for bronchitis and difficult breathing; one every fifteen to thirty minutes to induce expectoration in the adult.

AMMONIUM MURIATE.....1-4 gr.....	75
----------------------------------	----

Give one every half hour for five doses, in the coughs and colds of children.

AMMONIUM MURIATE.....1-2 gr.....	75
----------------------------------	----

A tablet allowed to dissolve on the tongue every twenty to sixty minutes as a gentle expectorant in dryness of the throat and slight colds.



<b>AMMON. SALICYLATE COMP</b>	{ Ammon. Salicylate. 1 gr. }	7 50
	{ Phenacetine ..... 1 gr. }	
	{ Caffeine ..... 1-2 gr. }	
	{ Salicin ..... 1-2 gr. }	

One to two tablets every two or three hours as an analgesic and antipyretic.

<b>AMENORRHŒA, ANÆMIC</b> ...	{ Mangan. Binox. .... 1 gr. }	2 00
	{ Iron Powd. .... 1 gr. }	
	{ Ext. Gentian. .... 1 gr. }	
	{ Ext. Cascara. .... 1-2 gr. }	
	{ Ext. Nux Vom. .... 1-6 gr. }	
	{ Capsicum Powd. .... 1-8 gr. }	
	{ Acid Arsenous. .... 1-50 gr. }	
	<b>Chocolate Coated only</b>	

One or two, three times a day, after meals. For anæmia, chlorosis amenorrhœa and atonic dyspepsia.

<b>ANÆMIA</b> ..... (Dr. Porter.)	{ Hæmogallol. .... 3 gr. }	16 00
	{ Ox Gall. .... 1 gr. }	
	{ Ext. Pancreatis. .... 3 gr. }	
	{ Strych. Sulph. .... 1-40 gr. }	
	{ Caffeine Mur. .... 1-2 gr. }	
	{ Ext. Colocynth. .... 1-2 gr. }	
	<b>Per 100</b> .....	1 65

Two tablets, three times a day after meals, for one day; then one tablet after meals.

<b>ANÆMIA No. 2</b> .....	{ Blaud's Mass. .... 3 gr. }	1 50
	{ Acid Arsenous. .... 1-60 gr. }	
	{ Mangan. Binoxide. 1 gr. }	
	<b>Chocolate Coated</b> .....	1 60

One to two, three times a day, after meals, for anæmia, chlorosis and menstrual disorders.

<b>ANÆSTHETIC</b> .....	{ Camphor. .... 1-4 gr. }	1 50
	{ Morphine Mur. .... 1-24 gr. }	
	{ Oil Cajeput. .... 1-24 min. }	

One every hour for diarrhœa.

<b>ANALGESIC</b> .....	{ Phenacetine. .... 2 1-2 gr. }	15 00
	{ Morph. Sulph. .... 1-16 gr. }	
	{ Tr. Aconite. .... 1 min. }	
	{ Fl. Ext. Bryonia. 1 1-2 min. }	
	{ Fl. Ext. Hyoscy. 2 1-2 min. }	
	<b>Per 100</b> .....	1 50

Give two at a dose if in severe pain, otherwise one every hour for four doses, then one every two to four hours; for rheumatism, neuralgia and grip.

<b>ANALGESIC No. 2</b> .....	{ Acetanilid. .... 2 gr. }	4 00
	{ Ammon. Salicylate. 1 gr. }	
	{ Quin. Sulphate. .... 1 gr. }	
	{ Codeine. .... 1-16 gr. }	

Two tablets every two or three hours. Relieves pain, reduces fever, allays cough, supports the heart.

<b>ANAPHRODISIAC</b> .....	{ Camphor. .... 2 gr. }	1 75
	{ Ext. Hyoscyamus. .... 1 gr. }	
	<b>Chocolate Coated</b> .....	1 85

One tablet morning and night.

<b>ANODYNE</b> .....	{ Camphor. .... 1-8 gr. }	1 10
	{ Ext. Hyoscyam. .... 1-8 gr. }	
	{ Morph. Sulph. .... 1-60 gr. }	
	{ Oleores. Capsic. .... 1-60 gr. }	
	<b>Chocolate Coated</b> .....	1 20

One tablet every half hour until drowsiness occurs; for insomnia, restlessness and neuralgia.

ANODYNE No. 2 . . . . . (Dr. H. P. Loomis.)	Ext. Hyoscyam. . . . . 2 gr.	3 50
	Camphor. . . . . 1 gr.	
	Morph. Sulph. . . . . 1-6 gr.	
<b>Chocolate Coated only</b>		

One, repeated every hour for not more than three doses, in cases of intense pain, neuralgia and tic-doloureux, intestinal and uterine colic.

ANODYNE No. 3 . . . . . (Dr. Clemens.)	Camphor . . . . . 1 gr.	2 25
	Ext. Hyoscyamus . . . 1 gr.	
	Morph. Acetate. . . 1-20 gr.	
	Oleores. Capsic. . . 1-20 gr.	
<b>Chocolate Coated only</b>		

One every half hour while in pain; stop after the fourth dose.

ANODYNE No. 4 . . . . . (Dr. Loving.)	Morphine Sulph. . . 1-16 gr.	2 25
	Ext. Cannab. Ind. . . 1-8 gr.	
	Oil Peppermint. . . 1-6 min.	
	Oleores. Capsic. . . 1-16 gr.	

One every half hour for cholera morbus and nausea, from pregnancy or other cause; for colic one every twenty minutes until four have been taken or until relief.

ANODYNE, For Infants. . . . . (Dr. Waugh.)	Nickel Bromide. . . 1-134 gr.	1 10
	Codeine Sulph. . . 1-67 gr.	
	Ipecac Powd. . . 1-134 gr.	
	Lithium Carb. . . 1-25 gr.	
	Oil Anise. . . . . 1-134 gr.	

One or two dissolved in hot water every fifteen to thirty minutes until effect is produced; for flatulent colic, restlessness, teething, etc.

ANTACID . . . . .	Calcium Carb. . . . 3 1-2 gr.	1 00
	Magnesia Carb. . . 2 1-2 gr.	
	Sodium Chlor. . . . . 1 gr.	

One, as required with a little water, for pyrosis and stomach indigestion.

ANTIASTHMATIC . . . . .	Potass. Iodide. . . . . 2 gr.	3 75
	Potass. Bromide. . . . 2 gr.	
	Fl. Ext. Euphor. Pil. 3 min.	
	Tinct. Lobelia. . . . . 2 min.	
	Nitroglycerin. . . . 1-200 gr.	

One tablet every three or four hours, for paroxysmal cough, bronchial spasm and dyspnoea.

ANTIASTHMATIC No. 2 . . . . . (Dr. L. T. Stevens.)	Potass. Iodide. . . . . 5 gr.	6 00
	Fowler's Solution. . . 1 min.	
	Tr. Lobelia. . . . . 4 min.	
	Tr. Belladonna. . . 1-2 min.	
	Tr. Opii Deod. . . . . 1-2 min.	
	Oil Wintergreen . . . 1-4 min.	
	Oil Sassafras. . . . . 1-4 min.	

One every three hours, for asthmatic cough and asthma.

ANTIASTHMATIC No. 3 . . . . . (Dr. Bennett.)	Potass Iodide. . . . . 2 1-2 gr.	3 25
	Potass Arsen. . . . . 1-100 gr.	
	Atropine Sulph. . . 1-350 gr.	
	Morph. Sulph. . . . . 1-120 gr.	

One to two tablets with water every three hours, for asthmatic cough and asthma.

ANTIBRONCHITIS . . . . . (Dr. O. A. Gordon.)	Potass. Nitrate. . . 2 1-2 gr.	1 10
	Tartar Emetic . . . 1-64 gr.	
	Dover's Powd . . . . . 1-4 gr.	
	Tr. Bryonia. . . . . 1-10 min.	
	Tr. Aconite . . . . . 1-4 min.	
	Oil Gaultheria . . . . . q. s.	

One every hour for four hours, then one as needed for cough.

When ordering or prescribing

<b>ANTICOLD.</b> ..... (Dr. O. A. Gordon.)	Morph. Sulph....	1-100 gr.	1 10
	Tartar Emetic....	1-100 gr.	
	Tr. Aconite....	1-2 min.	
	Camphor Powd....	1-8 gr.	
	Fl. Ext. Bellad.	1-32 min.	
		Oil Anise .....	1-20 min.
One tablet every hour until relieved.			

<b>ANTICONSTIPATION</b> .....	Ext. Cascara Sag....	1 gr.	2 10
	Ext. Nux Vom....	1-8 gr.	
	Ext. Belladonna....	1-8 gr.	
	Ipecac Powd....	1-8 gr.	
	Podophyllin.....	1-8 gr.	
<b>Plain or Chocolate Coated</b>			

One to three at night, or in divided doses through the day; for sluggishness of the intestinal glands and loss of tone in the muscular coat.

<b>ANTICONSTIPATION No. 2</b> .... (Dr. Dalton.)	Rhubarb Powd....	1 gr.	1 50
	Aloes.....	1-2 gr.	
	Jalap.....	1-2 gr.	
	Ipecac Powd....	1-4 gr.	
	Castile Soap.....	1-4 gr.	
		Strych. Sulph....	1-100 gr.
<b>Chocolate Coated</b> .....			1 60

One or two at night.

<b>ANTICONSTIPATION No. 3</b> .... (Dr. Hickling.)	Fel Bovis, Exsic....	2 gr.	3 50
	Strych. Sulph....	1-60 gr.	
	Ext. Cascara Sag....	3 gr.	
	Ext. Bellad....	1-16 gr.	
	Ipecac Powd....	1-16 gr.	
<b>Plain or Chocolate Coated</b>			

One at night, repeated next morning if necessary, for constipation.

ANTIDYSPEPTIC..... (Dr. Fothergill.)	{	Strych. Sulph....	1-120 gr.	I 10
		Ipecac Powd.....	1-3 gr.	
		Black Pepper Po....	1-8 gr.	
		Oil Gaultheria....	1-10 min.	
		Chocolate Coated.....		

One tablet after eating, repeated at the end of the hour, as a stimulant in retarded and deficient activity of the stomach.

<b>ANTIDYSPEPTIC No. 2</b> .....	Strych. Sulph....	1-60 gr.	1 25
	Ipecac Powd....	1-10 gr.	
	Capsicum Powd....	1-4 gr.	
	Ext. Rhubarb....	1-4 gr.	
	Ext. Gentian....	1-2 gr.	
		Sodium Bicarb....	2 gr.
<b>Chocolate Coated</b> .....			1 35

One tablet in water half an hour before meals to stimulate the appetite, and repeat half an hour after eating, to hasten digestion.

<b>ANTIFEBRIN COMP</b> ..... (Dr. Hoag.)	Caffeine Citrate....	1-2 gr.	3 75
	Sodium Bromide....	5 gr.	
	Antifebrin.....	2 gr.	

One tablet with fifteen drops aromatic spts. ammonia in water, and repeat every half hour if necessary, until four doses have been taken.

<b>ANTIFERMENTATIVE</b> .....	Sodium Salicylate....	2 gr.	2 00
	Ginger Powd.....	2 gr.	
	Capsicum Powd....	1-10 gr.	
	Cardamom Powd....	1-4 gr.	
<b>Plain or Chocolate Coated</b>			

One tablet every two or three hours for diarrhoea, flatulence and dysentery, and for the chronic intestinal dyspepsia of neurotic patients.

<b>ANTIMALARIAL</b> ..... (Dr. S. C. James.)	{ Nux Vom. Powd...1-4 gr. }	4 25
	{ Capsicum Powd...1-2 gr. }	
	{ Ext. Hyoscyam...1-2 gr. }	
	{ Quinine Sulph.....3 gr. }	
<b>Plain or Chocolate Coated</b>		

Give a tablet every two or three hours with plenty of water, in dumb ague and general malarial disorders; give one at six, four and two hours before the chills in acute attacks.

<b>ANTIMALARIAL No. 2</b> ..... (Dr. Guernsey.)	{ Cinchonidia Sulph...3 gr. }	2 65
	{ Capsicum Powd...1-2 gr. }	
	{ Nux Vomica.....1-5 gr. }	
<b>Plain or Chocolate Coated</b>		

Give one after each meal with plenty of water, in malaria of delicate women.

<b>ANTIMALARIAL No. 3</b> ..... (Dr. Bennett.)	{ Quin. Sulph.....1 1-2 gr. }	2 65
	{ Acid Arsen.....1-32 gr. }	
	{ Powd. Capsicum...1-2 gr. }	
<b>Plain or Chocolate Coated</b>		

One for a child, ten to fourteen years old, and two for an adult every three hours, for chronic and acute malaria.

<b>ANTIMALARIAL No. 4</b> ..... (Dr. Bennett.)	{ Quin. Sulph.....1 1-2 gr. }	2 70
	{ Acid Arsen.....1-32 gr. }	
	{ Strych. Sulph....1-30 gr. }	
	{ Capsicum Powd...1-2 gr. }	
<b>Plain or Chocolate Coated</b>		

One for a child, ten to fourteen years old, and two for an adult every three hours, for chronic and acute malaria.

<b>ANTIMONY &amp; IPECAC</b> ..... (Dr. Fowler.)	{ Tartar Emetic...1-100 gr. }	75
	{ Ipecac Powd....1-100 gr. }	

One every two hours in the secondary stages of a cold, and as a remedy for asthma of the "dry" form.

<b>ANTIPYRINE</b> .....1 gr.....	2 00
Antipyretic.	

One tablet every hour for a child; stop administration as temperature falls. For adults, five every hour with a glass of water, for three doses if necessary, for headache or fever.

<b>ANTIPYRINE</b> .....2 grs.....	3 00
Give a tablet every hour or two for headaches and fever.	

<b>ANTIPYRINE</b> .....5 grs.....	6 10
-----------------------------------	------

One tablet with a glass of water, every two or three hours while fever is high; for headache, one tablet every hour, for three doses if necessary.

<b>ANTIRHACHITIC</b> ..... (Dr. L. Weber.)	{ Precip. Chalk.....3 gr. }	1 25
	{ Calc. Phosph....1 1-2 gr. }	
	{ Iron Lactate.....2-5 gr. }	

One tablet three times a day, after eating, as a constructive in rickets and imperfect or retarded bone formation.

<b>ANTIRHACHITIC No. 2</b> ..... (Dr. L. Weber.)	{ Precip. Chalk...1 1-2 gr. }	1 00
	{ Calc. Phosph.....3-4 gr. }	
	{ Iron Lactate.....1-5 gr. }	

One to two tablets according to age, three times a day, after meals, as a constructive in rickets and imperfect or retarded bone formation.

<b>ANTIRHEUMATIC</b> .....	{ Acid Salicylic.....3 gr. }	1 85
	{ Sodium Bicarb...2 1-2 gr. }	
	{ Wine Colch. Seed.1 min. }	

One tablet after each meal, and one at bedtime, for chronic rheumatism; give more frequently in acute cases.

ANTIRHEUMATIC No. 2. ....	{ Potass. Iodide.....2 gr. Ext. Phytolacca.....1 gr. Resin Guaiac.....3 gr. Colchicin.....1-100 gr. Digitalia .....1-100 gr.	4 00
---------------------------	--	------

Give one every two hours in a recent attack, until its severity abates, then one every four hours for not more than three doses; three times a day for chronic cases.

ANTIRHEUMATIC No. 3. ....	{ Acid Salicylic .....3 gr. Sodium Bicarb. 2 1-2 gr. Wine Colch. Seed.1 min. Guaiac. ....2 gr.	2 00
---------------------------	---	------

One tablet after each meal, and one at bedtime, for chronic rheumatism; give more frequently in acute cases.

ANTISEPTIC DYSPEPTIC.....	{ Pepsin.....1 gr. Bismuth Subnit.....5 gr. Salol .....1 gr.	5 50
---------------------------	--	------

A tablet a few minutes after eating, to check fermentation of food and the production of gas.

ANTISEPTIC DYSPEPTIC, AN- ODYNE.....	{ Pepsin.....1 gr. Bismuth Subnit.....5 gr. Salol.....1 gr. Cocaine Hydro.1-12 gr.	7 50
---	---	------

One or two tablets after meals; for dyspepsia with gastric distress or pain.

ANTISEPTIC No. 1 .....	{ Corrosive Sub..7 3-10 gr. White or Blue. / Ammon. Mur..7 7-10 gr.	3 75
Per doz. screw cap bottles of 25 each, net		2 00

Dissolve one tablet in a pint of water; the strength of corrosive sublimate in the resulting solution is 1 to 1000.

ANTISEPTIC No. 2 .....	{ Corrosive Sub..3 7-10 gr. (White or Blue.) / Ammon. Mur..3 9-10 gr.	2 80
Per doz. screw cap bottles of 35 each, net		2 00

Dissolve one tablet in one half pint of water; the strength of corrosive sublimate in the resulting solution is 1 to 1000.

ANTISEPTIC No. 3 .....	{ Corrosive Sub..7 3-10 gr. White or Blue. / Sodium Chlor..7 7-10 gr.	3 75
Per doz. bottles of 25 each, net		2 00

Dissolve one tablet in one pint of water; the strength of corrosive sublimate in the resulting solution is 1 to 1000.

ANTISEPTIC No. 4.....	{ Corrosive Sub..3 7-10 gr. White or Blue. / Sodium Chlor..3 6-10 gr.	2 80
Per doz. bottles of 35 each, net		2 00

Dissolve one tablet in one half pint of water; the strength of corrosive sublimate in the resulting solution is 1 to 1000.

ANTISEPTIC PASTILLES. ....		1 60
(Dr. Carl Seiler,)		

For spray, mouth wash and nasal douche.

Dissolve one tablet in four tablespoonfuls of water, and use freely as a spray, mouth wash, gargle or nasal douche.

ANTISEPTIC IMPROVED.....		2 00
--------------------------	--	------

For spray, mouth wash and nasal douche.

Formula similar to Antiseptic Pastilles with addition of Hydrastine and Sanguinarine Nitrate.

Dissolve one tablet in four tablespoonfuls of water and use freely as a spray, mouth wash, gargle or nasal douche.

<b>APERIENT</b> ..... (Dr. Drysdale.)	{ Rhubarb Powd...1-4 gr. }	1 75
	{ Aloes Powd.....1-4 gr. }	
	{ Ipecac Powd.....5-12 gr. }	
	{ Nux Vomica.....1-2 gr. }	
	<b>Chocolate Coated</b> .....	1 85

A tablet at night, repeated next morning if necessary, as a gentle laxative.

<b>APHRODISIAC</b> .....	{ Ext. Cannab. Ind..1-2 gr. }	4 25
	{ Ext. Nux Vom.....1-2 gr. }	
	{ Ext. Ergot Aq.....2 gr. }	
	<b>Plain or Chocolate Coated</b>	

One tablet morning and night.

<b>APHRODISIAC No. 2</b> ..... (Dr. Landers.)	{ Ext. Nux Vom....1-8 gr. }	3 00
	{ Ext. Damiana.....1-4 gr. }	
	{ Phosphorus.....1-100 gr. }	
	{ Gold & Sod. Chl..1-20 gr. }	
	<b>Chocolate Coated only</b>	

One tablet after eating, three times a day, indicated in neurasthenia, functional impotence, spermatorrhœa and reflex nervous disorders, as a powerful nerve stimulant.

<b>APHRODISIAC No. 3</b> .....	{ Ext. Saw Palmet. 2 1-2 gr. }	2 75
	{ Phosphorus.....1-150 gr. }	
	{ Strych. Sulph....1-60 gr. }	
	{ Iron Phosphate..1-2 gr. }	
	<b>Chocolate Coated only</b>	

One after each meal as an aphrodisiac and general tonic.

<b>APOMORPHINE MUR. Cryst.</b> ..1-100 gr.....	1 30
<b>APOMORPHINE MUR. Cryst.</b> ..1-50 gr.....	1 65
<b>APOMORPHINE MUR. Cryst.</b> ..1-20 gr.....	2 50
<b>APOMORPHINE MUR. Cryst.</b> ..1-10 gr.....	4 00
<b>ARSENIC BROMIDE</b> .....1-100 gr.....	75

One tablet three times a day, as an alterative in sluggish skin diseases of children; stop when colic or fulness in the stomach appears.

<b>ARSENIC BROMIDE</b> .....1-50 gr.....	75
--	----

Give one three times a day, to control sugar excretion in diabetes, and as a stimulant in chronic skin affections. Discontinue the treatment if stomach fulness or colic appears.

<b>ARSENIC BROMIDE</b> .....1-40 gr.....	75
--	----

A tablet three or four times a day, for chronic eczema, psoriasis and malarial poisoning. Withhold the medicine if pain in the chest or abdomen occurs.

<b>ARSENIC BROMIDE</b> .....1-30 gr.....	75
--	----

One every three or four hours; for chronic skin diseases, epilepsy and malarial poisoning. Discontinue the administration if pain in the chest or abdomen occurs.

<b>ARSENIC BROMIDE</b> .....1-20 gr.....	75
--	----

One tablet every four hours given with care; the medicine to be withheld if pain in the stomach or abdomen should intervene; for chronic malarial infection, epilepsy and scaly skin diseases, where a tonic and stimulant are needed.

<b>ARSENIC CHLOROPHOS</b> .....1-50 gr.....	75
---	----

One tablet every three hours as a general tonic, and for malaria in subacute and chronic forms; stop if unpleasant symptoms occur.

<b>ARSENIC CHLOROPHOS</b> .....1-20 gr.....	75
---	----

One three times a day, stopping if pains in the chest or abdomen appear; for neuralgia, malaria and chronic skin diseases.

<b>ARSENICUM COMP.</b> .....	{ Acid Arsenous... 1-100 gr. }	75
(Dr. F. X. Spranger, Sr.)	{ Merc. Sol. II... 1-100 gr. }	
	{ Sulphur ..... 1-100 gr. }	

One tablet every three hours, for skin and mucous diseases, dependent on and resultant of a strumous or scrofulous diathesis.

<b>ARSENIC IODIDE</b> .....	1-1000 gr.	75
Alterative.		

One every three hours, for an infant with malaria or chronic disease of the skin, or atonic diarrhoea.

<b>ARSENIC IODIDE</b> .....	1-100 gr.	75
-----------------------------	-----------	----

One every three hours, as a stimulant tonic to the cutaneous functions, and for malarial headache. Decrease the frequency of the dose if colic or pain in the chest appear.

<b>ARSENIC IODIDE</b> .....	1-50 gr.	75
-----------------------------	----------	----

Give one or two tablets after eating, as a general tonic in malaria, and as an alterative in diseases of the blood. Stop if pains in the chest or stomach develop.

<b>ARSENIC IODIDE</b> .....	1-25 gr.	75
-----------------------------	----------	----

One after each meal; stop when pains occur in chest or abdomen; for malaria, epilepsy, psoriasis and chronic skin diseases.

<b>ARSENIC IODIDE</b> .....	1-10 gr.	75
-----------------------------	----------	----

One tablet after meals, given with care, as a general tonic and for malaria, epilepsy, psoriasis and chronic eczema.

<b>ARSENIC &amp; IRON No. 1.</b> .....	{ Acid Arsenous... 1-30 gr. }	75
Tonic, hæmatic.	{ Iron Powd. .... 1 gr. }	

Give a tablet three times a day, for anæmia and where a general tonic is indicated.

<b>ARSENIC &amp; IRON No. 2.</b> .....	{ Acid Arsenous... 1-60 gr. }	75
	{ Iron Powd. .... 1-5 gr. }	

One or two after each meal, for malaria, anæmia and chlorosis. Stop on the occurrence of pain in the stomach or abdomen.

<b>ARSENIC &amp; IRON No. 3.</b> .....	{ Acid Arsenous... 1-100 gr. }	75
	{ Iron Powd. .... 1 gr. }	

A tablet every two or three hours, for simple and pernicious anæmia, where a general tonic is required, and during convalescence after wasting disease.

<b>ARSENIC &amp; IRON No. 4.</b> .....	{ Acid Arsenous... 1-60 gr. }	75
	{ Iron Powd. .... 1 gr. }	

One tablet after meals, for anæmia and chlorosis, malarial poisoning, and as a general tonic.

<b>ARSENIC RED SULPHURET.</b>	1-100 gr.	75
-------------------------------	-----------	----

Alterative.

**Chocolate Coated**.....

One three times a day, increasing one every second or third day; for affections of the skin accompanied with suppuration. Stop when the physiological effect is produced, or pain or fulness of the stomach occurs.

<b>ARSENIC RED SULPHURET.</b>	1-50 gr.	75
-------------------------------	----------	----

**Chocolate Coated**.....

One after each meal, gradually increasing to three after meals; for chronic eczema and psoriasis. Stop when the effect is produced, or pain or fulness of the stomach occurs.

<b>ARSENIC RED SULPHURET.</b>	1-40 gr.	75
-------------------------------	----------	----

**Chocolate Coated**.....

One three times daily, as a general alterative for diseases of the blood and skin. Stop when the effect is produced, or pain or fulness of the stomach occurs.

<b>ARSENIC RED SULPHURET.</b> 1-30 gr .....	75
<b>Chocolate Coated</b> .....	85
A tablet after each meal, for boils and suppurative skin diseases. Stop when the effect is produced, or pain or fulness of the stomach occurs.	
<b>ARSENIC RED SULPHURET.</b> 1-20 gr .....	75
<b>Chocolate Coated</b> .....	85
One three times a day, in eczema, psoriasis and other chronic diseases of the skin. Stop when the effect is produced, or pain or fulness of the stomach occurs.	
<b>ARSENIC &amp; STRYCHNINE</b> .... ( Acid Arsenous..1-100 gr. )	75
Tonic, alterative. } Strych. Sulph....1-60 gr. }	
<b>Chocolate Coated</b> .....	85
One tablet after eating, three times a day; as a general tonic and alterative, especially in obscure malarial affections.	
<b>ARTHRITIC</b> ..... { Sodium Salicylate...5 gr. }	3 00
..... { Tr. Cimicifuga....5 min. }	
..... { Tr. Gelsemium....5 min. }	
One every three hours for rheumatism.	
<b>ASAFCETIDA</b> .....2 grs.....	1 15
<b>Chocolate Coated only.</b>	
One to three after meals and at bedtime, as a carminative, and an antispasmodic in nervous dyspepsia. Valuable also in allaying the nervous symptoms of neurasthenic and hysterical patients.	
<b>ASAFCETIDA</b> .....3 grs.....	1 35
<b>Chocolate Coated only.</b>	
One to three after meals and at bedtime, as a carminative, and an antispasmodic in nervous dyspepsia. Valuable also in allaying the nervous symptoms of neurasthenic and hysterical patients.	
<b>ASAFCETIDA</b> .....4 grs.....	1 50
<b>Chocolate Coated only.</b>	
One to two after meals and at bedtime, as a carminative, and an antispasmodic in nervous dyspepsia. Valuable also in allaying the nervous symptoms of neurasthenic and hysterical patients.	
<b>ASAFCETIDA</b> .....5 grs.....	1 50
<b>Chocolate Coated only.</b>	
One to two after meals and at bedtime, as a carminative, and an antispasmodic in nervous dyspepsia. Valuable also in allaying the nervous symptoms of neurasthenic and hysterical patients.	
<b>ASAFCETIDA &amp; NUX VOMICA.</b> { Asafoetida.....3 gr. }	1 50
{ Ext. Nux Vomica..1-4 gr. }	
<b>Chocolate Coated only</b>	
One after meals and at bedtime, as a carminative, and antispasmodic in nervous dyspepsia, and a nerve tonic in neurasthenia and hysteria.	
<b>ASTRINGENT COMP.</b> ..... { Ext. Hæmatoxylon..1 gr. }	2 00
..... { Tr. Opii Camph....6 min. }	
..... { Acid Sulph. Arom...1 gr. }	
..... { Pulv. Aromatic.....1 gr. }	
One to three tablets according to the severity of the case, repeated in one or two hours; for diarrhœa and dysentery, and before the stage of collapse in cholera infantum.	
<b>ASTRINGENT WASH</b> ..... { Lead Acetate.....2 gr. }	2 00
..... { Zinc Acetate.....2 gr. }	
..... { Hydrastin Mur....1-20 gr. }	
..... { Morph. Acetate...1-32 gr. }	
One in from four to eight tablespoonfuls of water for use as an injection in gonorrhœa, and also as a local application for sores, ulcers and moist skin diseases.	

When ordering or prescribing

<b>ATONIC DYSPEPTIC</b> .....	{ Warburg's Tr...10 min. F. E. Viburn. Prun. 10 min. F. E. Hydrastis. 10 min. }	
(Dr. J. Fuhs.)	<b>Plain or Chocolate Coated</b>	6 00
Two tablets three times a day after meals, for atonic dyspepsia.		
<b>ATONIC DYSPEPTIC No. 2</b> ....	{ Pepsin Powd.....2 gr. Pancreatin.....2 gr. Oleoresin Capsic. 1-4 min. Oleores. Ginger. 1-4 min. Creosote B'wood. 1-2 min. }	
(Dr. Hare.)	<b>Plain or Chocolate Coated</b>	3 65
One tablet an hour before each meal in atonic dyspepsia		
<b>ATROPINE COMP</b> .....	{ Atropine Sulph. 1-400 gr. }	
(Dr. Gilbert.)	{ Tartar Emetic. 1-100 gr. }	75
Respiratory corrective.		
One every three hours in bronchitis with asthmatic complications.		
<b>ATROPINE, GELSEMIUM CO.</b>	{ Fl. Ext. Gelsem...1 min. Atropine Sulph. 1-400 gr. Camphor. ....1-2 gr. Morphine Sulph. 1-32 gr. }	
(Dr. J. E. Riggs.)	<b>Chocolate Coated</b> .....	1 60 1 70
One every two hours until four are taken, for coryza.		
<b>ATROPINE SULPHATE</b> .....	1-1000 gr ..	75
One tablet every half hour for three doses, then every three hours, for an infant with fresh cold.		
<b>ATROPINE SULPHATE</b> .....	1-500 gr.....	75
For children, one tablet every three hours, for incontinence of urine, recent colds, and to lessen the sweating of consumption and rheumatism.		
<b>ATROPINE SULPHATE</b> .....	1-400 gr.....	75
A tablet every half hour to relieve the pain of muscular rheumatism and neuralgia, and to control sweating.		
<b>ATROPINE SULPHATE</b> .....	1-200 gr.....	75
One every hour for four doses, then one every two hours, for a fresh cold.		
<b>ATROPINE SULPHATE</b> .....	1-150 gr.....	90
One every three hours for five doses, to allay vesical spasm and neuralgia, and in recent cold.		
<b>ATROPINE SULPHATE</b> .....	1-120 gr.....	90
One every three hours to relieve neuralgia and sweating of phthisical patients, and in fresh colds with abundant secretion.		
<b>ATROPINE SULPHATE</b> .....	1-100 gr.....	1 00
One every two hours for three doses, to allay vesical spasm and neuralgia, and in the early stages of colds.		
<b>ATROPINE SULPHATE</b> .....	1-50 gr ..	1 25
One tablet as an anodyne, repeated cautiously after an interval of three hours; also to relieve dyspnœa in inflammations of thoracic viscera.		
<b>BAPTISIA FL. EXT.</b> .....	1 minim.....	1 00
Diaphoretic, local astringent.		
One tablet every three hours, as a diaphoretic, and to stimulate hepatic and intestinal functions.		
<b>BELLADONNA COMP</b> .....	{ Corrosive Sub...1-100 gr. }	
(Dr. Rankin.)	{ Tr. Belladonna,...1 min. }	1 10
	{ Ipecac Powd.....1-10 gr. }	
One every hour through the day, for ulcerated sore throat, first stages of tonsillitis and pharyngitis.		

<b>BELLADONNA EXTRACT</b> ....	1-100 gr. ....	75
A tablet every two hours for six doses, for an infant with diarrhœa and fever.		
<b>BELLADONNA EXTRACT</b> ....	1-50 gr. ....	75
One every half hour for six doses, for facial neuralgia.		
<b>BELLADONNA EXTRACT</b> ....	1-10 gr. ....	75
One every hour or two to control pain in neuralgia, and for profuse sweating in phthisis, and in recent colds.		
<b>BELLADONNA EXTRACT</b> ....	1-8 gr. ....	75
A tablet every hour in recent inflammations of the respiratory tract, until three doses are taken, or dryness of the throat occurs.		
<b>BELLADONNA EXTRACT</b> ....	1-5 gr. ....	75
Give one every two hours in recent colds, rhinitis or cold in the head, and to relieve neuralgia.		
<b>BELLADONNA EXTRACT</b> ....	1-4 gr. ....	75
One tablet every three hours in pharyngitis, tonsillitis and neuralgia of the head and face.		
<b>BELLADONNA FL. EXT</b> .....	1 minim. ....	1 10
Dissolve one tablet in ten tablespoonfuls of water, and give one tablespoonful of the solution every half hour to a child with fever and headache, or recent cold.		
<b>BETA NAPHTOL</b> .....	1 gr. ....	1 00
One every half hour, with antipyretics, for tubercular and other ulceration of the bowels; when symptoms improve, give one tablet every two or three hours.		
<b>BETA NAPHTOL CO. No. 2</b> ....	<div> <div> { Beta Naphtol.....1-2 gr. Bismuth Subnit.2 1-2 gr. Sodium Bicarb.2 1-2 gr. Chocolate Coated..... } </div> </div>	<div> <div>1 75</div> <div>1 85</div> </div>
One or two after meals for flatulent dyspepsia, and as an intestinal antiseptic in diarrhœa.		
<b>BISMUTH AROMATIC</b> .....	<div> <div> { Bismuth Subnit.....2 gr. Paregoric.....3 min. Oil Cinnamon...1-20 min. } </div> </div>	<div> <div>2 25</div> </div>
(Dr. Parrish, St. Louis.)		
For a child one to three years of age, one tablet every hour until diarrhœa is controlled. For older children, two every hour. For all cases of inflammatory diarrhœa.		
<b>BISMUTH, CARBOLIC COMP.</b>	<div> <div> { Bismuth Subnit.....1 gr. Calomel....1-40 gr. Ipecac Powd....1-40 gr. Acid Carbolie....1-15 gr. } </div> </div>	<div> <div>1 50</div> </div>
(Dr. Ross Foster.)		
One every half to one hour in vomiting of adults or children. One every hour in cholera infantum.		
<b>BISMUTH CARBONATE CO</b> ....	<div> <div> { Bismuth Subcarb....1 gr. Sodium Bicarb.....1 gr. Oil Peppermint.1-50 min. } </div> </div>	<div> <div>1 50</div> </div>
One every half hour, with a little water on the end of a spoon, for sour stomach of infants, with "spitting up" of fermented milk. Meanwhile give no milk to child, only hot water or barley water.		
<b>BISMUTH, CATECHU CO</b> .....	<div> <div> { Salol .....2 1-2 gr. Bismuth Subnit.2 1-2 gr. Opium Powd.....1-2 gr. Catechu Powd....1-2 gr. Acid Hydrochl.Dil 1-2 gr. Oil Cinnamon ...1-8 min. } </div> </div>	<div> <div>5 75</div> </div>
(Dr. C. C. Hoyt.)		
A tablet after meals and one at night, in chronic and subacute intestinal dyspepsia, and irritative diarrhœa.		

When ordering or prescribing

<b>BISMUTH COMP.</b> . . . . .	{ Bismuth Subnit. . . 1-2 gr. } { Cerium Oxalate. . . 1-2 gr. }	1 20
One or two, repeated every hour for six doses, or until effect is produced; for nausea and as a sedative for sick headache.		
<b>BISMUTH &amp; CALOMEL CO.</b> . . . . .	{ Bismuth Subnit. . . . 1 gr. } { Calomel . . . . . 1-40 gr. } { Ipecac Powd. . . . 1-60 gr. }	1 50
Alterative stomachic.		
A tablet or two after meals, as a gentle alterative and tonic for the stomach, in dyspepsia from ice-water and overeating.		
<b>BISMUTH &amp; CHARCOAL.</b> . . . . .	{ Bismuth Subnit. . . . 2 gr. } { Charcoal . . . . . 5 gr. }	2 00
Absorbent stomachic.		
One to three tablets after meals, for dyspepsia with fermentation and formation of gas.		
<b>BISMUTH, CHARCOAL CO.</b> . . . . .	{ Bismuth Subcarb. . . 2 gr. } { Charcoal . . . . . 2 gr. } { Sodium Bicarb. . . . 3 gr. } { Oil Peppermint. 1-10 min. }	2 00
One after eating, repeated in half an hour, for flatulent dyspepsia and heartburn. Avoid starchy food.		
<b>BISMUTH &amp; IPECAC COMP.</b> . . . . .	{ Bismuth Subnit. . . . 1 gr. } { Calomel . . . . . 1-10 gr. } { Ipecac Powd. . . . 1-20 gr. } { Opium Powd. . . . 1-40 gr. }	1 50
Alterative stomachic.		
One every fifteen to thirty minutes, for diarrhœa, to check the discharges easily and gradually, and at the same time to act as an intestinal antiseptic.		
<b>BISMUTH &amp; IPECAC CO. No. 2</b> . . . . .	{ Bismuth Subnit. . . 1-2 gr. } { Cerium Oxalate. . . 1-2 gr. } { Ipecac Powd. . . . 1-20 gr. }	1 30
A tablet as indicated in dyspepsia with nausea and headache.		
<b>BISMUTH &amp; MORPHINE.</b> . . . . .	{ Bismuth Subnit. . . . 4 gr. } { Morphine Sulph. . . 1-16 gr. }	3 65
One to three tablets for vomiting and gastric pain, repeated if necessary in an hour.		
<b>BISMUTH &amp; NUX VOMICA.</b> . . . . .	{ Bismuth Subnit. . . . 1 gr. } { Ext. Nux Vom. . . . 1-8 gr. }	1 50
One to three tablets, three times a day for atonic dyspepsia with lack of appetite.		
<b>BISMUTH OPIUM COMP.</b> . . . . .	{ Bismuth Subnit. . . . 4 gr. } { Corros. Sub. . . . 1-96 gr. } { Opium Powd. . . . 1-4 gr. } { Oil Anise . . . . . q. s. }	3 25
One every two or three hours for colliquative diarrhœa.		
<b>BISMUTH SALICIN COMP.</b> . . . . .	{ Bism. Subgall. . . 2 1-2 gr. } { Salicin . . . . . 5 gr. } { Strych. Sulph. . . 1-100 gr. }	6 50
(Dr. Hubbard.)		
Antifermentative and tonic.		
Two tablets four times a day, after meals and at bedtime, for atonic dyspepsia.		
<b>BISMUTH SALICYLATE.</b> . . . . . 1 gr. . . . .		1 50
Stomachic.		
Give a tablet every hour or two, for intestinal fermentation; allow plenty of drinking water.		
<b>BISMUTH SALICYLATE.</b> . . . . . 2 grs. . . . .		2 50
Give a tablet every hour or two, for intestinal fermentation; allow plenty of drinking water.		

**BISMUTH SALICYLATE**.....5 grs..... 3 75

A tablet every two or three hours, for inflammation of the intestines, typhoid fever and dysentery; allow a free supply of drinking water.

**BISMUTH SALICYLATE AROMATIC**.....  
 { Bism. Salicyl. basic. 3 gr. }  
 { Creta Prep. .... 2 gr. } 3 25  
 { Pulv. Aromat. .... 1 gr. }  
 { Zinc Sulphocarb. 1 1-4 gr. }

Give one tablet three or four times a day, in atony of the intestines, with loss of function.

**BISMUTH SALICYLATE CO.** { Bismuth Salicyl. .... 3 gr. }  
**No. 2**..... { Salol ..... 3 gr. } 5 75  
 { Sodium Bicarb. .... 2 gr. }

One after meals for stomach or intestinal indigestion, accompanied by excessive flatulence or diarrhoea.

**BISMUTH SUBCARBONATE**..1 gr..... 1 50

A tablet every hour until six are taken, for a child with vomiting and the diarrhoea of teething.

**BISMUTH SUBCARBONATE**..2 grs..... 2 00

One or two according to symptoms, for nausea, pain in the stomach and vomiting; repeated every half hour for three doses.

**BISMUTH SUBCARBONATE**..5 grs..... 3 75

One tablet, and repeat as necessary, for diarrhoea, dyspepsia, nausea and vomiting.

**BISMUTH SUBGALLATE**...1 gr..... 1 50

One every hour or two, for vomiting and diarrhoea of infants.

**BISMUTH SUBGALLATE**.....2 grs..... 2 00

One to three, according to age, three times a day, after meals, for flatulent or fermentative dyspepsia.

**BISMUTH SUBGALLATE**.....3 grs..... 2 60

One or two, three times a day, for fermentative or flatulent dyspepsia.

**BISMUTH SUBGALLATE**.....4 grs..... 3 25

One, three times a day, for flatulent or fermentative dyspepsia.

**BISMUTH SUBGALLATE**.....5 grs..... 3 75

One, three times a day, for flatulent or fermentative dyspepsia.

**BISMUTH SUBGALLATE CO.** { Bismuth Subgall. .... 5 gr. }  
 (Dr. Aulde.) { Pepsin Pure ..... 1 gr. } 4 00  
 { Ext. Nux Vom. .... 1-4 gr. }

One tablet before and after eating, for cases of slow or imperfect digestion of food in the stomach.

**BISMUTH SUBGALLATE CO.** { Bism. Subgall. ... 3 1-2 gr. }  
**No. 2.** (Dr. Aulde.) { Ext. Pancreatis.. 1 1-2 gr. } 4 25  
 { Ext. Nux Vom. .... 1-4 gr. }

One tablet before meals, and two hours after, for intestinal indigestion.

**BISMUTH SUBNITRATE**.....1 gr..... 1 25

Stomachic.

One tablet every two hours, for the vomiting and diarrhoea of teething children; give only hot water and sterilized milk in small quantities until symptoms abate.

**BISMUTH SUBNITRATE**.....2 grs..... 1 75

A tablet half an hour after eating, repeated every half hour for four doses, as a stomachic tonic and for intestinal dyspepsia.

**BISMUTH SUBNITRATE**.....5 grs..... 3 00

One, three times a day, in painful stomach dyspepsia; one every two hours, for diarrhoea.

---

When ordering or prescribing

<b>BISMUTH SUBNITRATE</b> .....10 grs.....		5 00
One as needed, to allay nausea, vomiting, and gastralgia.		
<b>BITTER TONIC</b> .....	{ Wine Ipecac.....1 min. Tr. Capsicum.....2 min. Tr. Nux Vom.....5 min. Tr. Gentian Co.....5 min. }	
(Dr. Frank E. West.)		
	<b>Plain or Chocolate Coated</b>	2 00
One tablet just before eating, three times a day.		
<b>BLADDER</b> .....	{ Sodium Benzoate...5 gr. } { Sodium Biborate...5 gr. }	2 25
(Dr. Noble.)		
One every two hours, followed by a glass of water; for low grade cystitis with alkaline urine.		
<b>BLANK TABLETS</b> .....	White, Yellow and Pink....	50
<b>BLAUD'S PILL</b> .....1 gr.....		75
Tonic.	<b>Chocolate Coated</b> .....	85
One after each meal, increasing one a day until five are taken three times a day, in anæmia and chlorotic affections.		
<b>BLAUD'S PILL COMP.</b> .....	{ Bland's Mass.....3 gr. } { Acid Arsenous...1-60 gr. }	1 35
	<b>Chocolate Coated</b> .....	1 45
One, three times a day, for a tonic.		
<b>BLAUD'S PILL COMP. No. 2.</b> ..	{ Bland's Mass.....5 gr. } { Acid Arsenous...1-40 gr. }	1 50
	<b>Chocolate Coated</b> .....	1 60
One, three times a day, for a tonic.		
<b>BLAUD'S PILL CO. No. 4.</b> .....	{ Bland's Mass.....5 gr. } { Corros. Sublim...1-80 gr. } { Strychnine Sulph...1-60 gr. } { Acid Arsenous...1-50 gr. }	1 65
	<b>Chocolate Coated</b> .....	1 75
One, three times a day, as an iron tonic and alterative in anæmia and general debility.		
<b>BLAUD'S PILL COMP. No. 5.</b> ..	{ Bland's Mass.....5 gr. } { Strych. Sulph....1-40 gr. }	1 70
(Dr. Raymond Clark.)	<b>Chocolate Coated</b> .....	1 80
One after meals as a general tonic in anæmia.		
<b>BLAUD'S PILL COMP. IMP.</b> ...	{ Bland's Mass.....5 gr. } { Acid Arsen....1-50 gr. } { Ext. Nux Vomica...1-8 gr. } { Ext. Cascara.....1 gr. }	
	<b>Plain or Chocolate Coated</b>	2 10
One after meals as a general tonic in anæmia with constipation.		
<b>BLAUD'S &amp; NUX VOMICA</b> ....	{ Bland's Mass.....3 gr. } { Ext. Nux Vom...1-10 gr. }	1 60
	<b>Chocolate Coated</b> .....	1 70
One, three times a day, as an iron and general tonic.		
<b>BLAUD'S TABLETS</b> .....3 grs.....		1 30
	<b>Chocolate Coated</b> .....	1 40
One, three times a day, as an iron tonic.		
<b>BLAUD'S TABLETS</b> .....5 grs.....		1 50
	<b>Chocolate Coated</b> .....	1 60
One, three times a day, as an iron tonic.		
<b>BLAUD'S TONIC LAXATIVE.</b> ..	{ Bland's Mass.....3 gr. } { Quinine Sulph....1-2 gr. } { Nux Vom. Ext...1-10 gr. } { Aloin.....1-5 gr. }	
	<b>Chocolate Coated only</b>	2 00
One to two, three times a day after meals, as a tonic and hæmatic, in anæmia and chlorosis.		

BLAUD'S TONIC LAXATIVE. No. 2.....	{	Blaud's Mass.....	3 gr.	}
		Quinine Sulph....	1-2 gr.	
		Ext. Nux Vom.,	1-10 gr.	
		Aloin.....	1-10 gr.	
Chocolate Coated only				2 00

One to two, three times a day after meals, as a tonic and hæmatic, in anæmia and chlorosis.

<b>BLUE MASS</b> .....	1-2 gr.	75
Hepatic stimulant.		

One tablet every two or three hours until six are taken, and repeat after an interval of a day; for torpidity of the liver and beginning biliousness.

<b>BLUE MASS</b> .....	1 gr.	1 00
------------------------	-------	------

One tablet every three hours until effect is produced, then twice or three times daily; for torpor of the liver, biliousness and cirrhosis in the early stages.

<b>BLUE MASS</b> .....	5 grs.	2 15
------------------------	--------	------

One or two at night; followed by a seidlitz powder or a bottle of citrate of magnesia next morning, for a purge.

<b>BLUE MASS. COMP.</b> ..... (Dr. L. D. Rockwell)	{	Blue Mass.....	1 gr.	} 1 30
		Rhubarb.....	1 gr.	
		Sodium Bicarb.....	1 gr.	
		Oil Peppermint.....	q. s.	

One tablet every three or four hours until effect is produced, then one morning and night for torpid liver, biliousness and constipation.

<b>BORAX</b> .....	3 grs.	75
--------------------	--------	----

Dissolve a tablet in six tablespoonfuls of water, and of this give one teaspoonful every half hour, for summer dyspepsia and diarrhœa of infants.

<b>BORAX</b> .....	5 grs.	90
--------------------	--------	----

One tablet after meals, to correct acidity of the stomach and intestinal fermentation; one on the tongue for sore throat.

BORAX, ACONITE COMP..... (Dr. Sadler.)	{	Ammon Mur.....	1 gr.	} 1 15
		Borax Powd.....	2 gr.	
		Sugar.....	2 gr.	
		Tr. Aconite Rad..	1 min.	
		Oil Gaultheria..	1-2 min.	

One to two every two hours, slowly dissolved in the mouth, for sore throat, or dissolved in two teaspoonfuls of water, and used with douche, for inflammation of nasal passages.

BORAX COMP..... (Fraser & Co.)	{	Borax.....	1 gr.	I 10
		Saccharum Alb.....	5 gr.	
		Oil Gaultheria.	I-20 min.	

One, dissolved slowly on the tongue every hour or two, for sore throat, cough, and sore mouth.

BREAKFAST TABLET..... (Dr. J. G. Wiltshire.)	{ Aloes Socot.....1 gr. }	1 50
	{ Ext. Nux Vom....1-2 gr. }	
	{ Ext. Hyoscyamus. 1-2 gr. }	

One after breakfast, as often as necessary, for chronic constipation.

<b>BROMIDE COMP.</b> .....	{ Morph. Brom.....1-8 gr. }	7 50
(Dr. Hubbard.)	{ Hyosc. Hydrob., 1-100 gr. }	
Anodyne.	{ Camph. Monobr.... 1 gr. }	

One or two at night to quiet the nerves and induce sleep; one every half hour for three doses to produce a strong anodyne action.

<b>BROMIDE COMP. No. 2</b> ..... (Dr. Hubbard.)	{	Morph. Brom....	1-16 gr.	4 50
		Hyosc. Hydrob.,	1-200 gr.	
		Camph. Monobr.	1-2 gr.	

A tablet every hour for three doses, for insomnia caused by pain; also for irritable bladder.

When ordering or prescribing

BRONCHIAL .....	Ammon Mur. ....	1-3 gr.	1 00
	Ext. Licorice.....	1 gr.	
	Balsam Tolu.....	1-5 gr.	
	Oleores. Cubebs....	1-3 gr.	
	Henbane Powd....	1-4 gr.	
	Senega Powd.....	1-5 gr.	
	Ipecac Powd.....	1-50 gr.	

One tablet, slowly dissolved in the mouth, every hour or two, for bronchial irritation.

BRONCHIAL No. 2..... (Dr. Wood.)	Ext. Hyoscyamus..	1-5 gr.	1 50
	Morph. Sulph....	1-100 gr.	
	Tartar Emetic....	1-100 gr.	
	Ext. Aconite.....	1-100 gr.	
	Ext. Bellad.....	1-100 gr.	
	Merc. Iod. Red....	1-100 gr.	
	Potass. Bichrom..	1-300 gr.	
	Bryonia.....	1-200 gr.	
	Oil Anise.....	1-20 min.	

One or two as needed, every hour, for inflammation of throat and bronchial tubes.

BRONCHIAL No. 3..... (Dr. E. G. Rankin.)	Tartar Emetic....	1-50 gr.	1 00
	Tr. Bryonia.....	1-2 min.	

One tablet hourly, or every two hours, for bronchial affections with muscular or pleuritic pain in chest.

BRONCHITIS.....	Ext. Belladonna..	1-40 gr.	1 50
	Dover's Powder..	1-10 gr.	
	Ipecac Powd....	1-20 gr.	
	Quinine Sulph....	1-4 gr.	
	Chocolate Coated.....		1 60

One every hour for four or five doses, then one every two hours for bronchial cough.

BRONCHITIS, Without Quinine	Ext. Belladonna..	1-40 gr.	1 00
	Dover's Powder..	1-10 gr.	
	Ipecac Powd....	1-20 gr.	

One every hour for four or five doses, then one every two hours, for bronchial cough.

BRONCHITIS No. 2, Acute..... (Dr. T. G. Davis.)	Tr. Aconite.....	1-5 min.	1 00
	Tr. Bryonia.....	1-10 min.	
	Tr. Belladon....	1-10 min.	
	Antimony Sul....	1-100 gr.	
	Potass. Bichrom..	1-100 gr.	

One every half hour until the bronchial secretions act normally, then every two hours as long as benefit is experienced in bronchial cough and irritation; also useful in the chronic bronchitis of the aged; one tablet four times daily.

BRONCHITIS No. 3..... (Dr. N. L. North.)	Tinct. Digitalis....	1 min.	2 00
	Terpin Hydrate....	1 gr.	
	Dover's Powder....	1-2 gr.	
	Tinct. Belladon....	1 min.	
	Tinct. Sanguinar..	1 min.	
	Ammon. Mur.....	1 gr.	
	Acid Benzoic.....	1-2 gr.	

One tablet, best allowed to dissolve slowly on the back of the tongue or dissolved in water and slowly swallowed, three or four times a day, for winter cough or bronchitis of the aged, or for abortive effects when given in first stages of grip.

BRONCHITIS No. 4..... (Dr. J. C. Bierwirth)	Tr. Sanguinar..	2 1-2 min.	1 50
	Wine Ipecac....	2 1-2 min.	
	Tr. Aconite.....	1 min.	

One tablet every two hours for bronchitis in the aged.

<b>BROWN AMMONIA</b> .....	{ Ammon. Mur.....4 gr. Brown Mixture...1 teasp. Oil Anise.....1-10 min. }	1 50
----------------------------	---	------

One tablet every three hours, for a cough.

<b>BROWN MIXTURE</b> ..... (One teaspoonful.) Anodyne expectorant.	{ Ext. Licorice.....1-10 gr. Camphor.....1-25 gr. Acid Benzoic.....1-25 gr. Oil Anise.....1-25 min. Opium Powd....1-25 gr. Tartar Emetic...1-60 gr. }	1 00
--	--	------

One on the tongue every hour, for bronchial cough. Stop when oppression of chest is relieved and secretion well established.

<b>BROWN MIXTURE</b> ..... (One-half teaspoonful.)	{ Ext. Licorice.....1-20 gr. Camphor.....1-50 gr. Acid Benzoic.....1-50 gr. Oil Anise.....1-50 min. Opium Powd....1-50 gr. Tartar Emetic...1-120 gr. }	75
---	---	----

Dissolve one in the mouth every half hour, stopping as the symptoms lessen and the cough becomes free; for bronchial coughs and colds.

<b>BROWN MIXTURE COMP</b> ....	{ Brown Mixture..1 teasp. Ammon. Mur.....1 gr. }	1 25
--------------------------------	---	------

A tablet every two hours, allowed to dissolve in the mouth, for coughs and colds with thick and difficult expectoration; lessen the frequency of the dose as the symptoms improve.

<b>BROWN MIXTURE CO. No. 2.</b>	{ Brown Mixture..1 teasp. Ammon. Mur.....5 gr. }	1 35
---------------------------------	---	------

Take one tablet with water, or dissolve it in the mouth, every two or three hours; for bronchitis and severe cough with scanty sputa; continue until the bronchial secretions are restored.

<b>BRUCIA</b> .....	1-100 gr.....	75
---------------------	---------------	----

Tonic.

One to two tablets, three times a day, for a stimulant to the stomach and a general nerve tonic.

<b>BRUCIA</b> .....	1-50 gr.....	75
---------------------	--------------	----

One tablet three times a day, for a stimulant to the stomach and a general nerve tonic.

<b>BRUCIA COMP</b> ..... (Dr. Wood.) Tonic antiperiodic.	{ Quinine Sulph....1-2 gr. Brucia.....1-100 gr. Acid Arsenous...1-100 gr. }	1 75
--	---	------

Give one after each meal as a general tonic.

<b>BRYONIA ALB</b> .....	1-200 gr.....	75
--------------------------	---------------	----

Membranous stimulant.

One to three tablets, repeated every two hours as necessary, for infantile pleurisy and bronchitis.

<b>BRYONIA ALB</b> .....	1-100 gr.....	75
--------------------------	---------------	----

Give one or two tablets every hour in bronchitis and pleurisy.

<b>CACTUS COMPOUND</b> .....	{ Cactus Fl. Ext....1 min. Tr. Strophanthus..2 min. }	2 35
------------------------------	--	------

A tablet every three or four hours, according to the severity of the symptoms, in cardiac weakness and irregularity, and as a diuretic.

<b>CACTUS FL. EXTRACT</b> .....	1 minim.....	1 70
---------------------------------	--------------	------

Heart stimulant.

One or two tablets three times daily, as a heart stimulant in weak and irregular action from valvular disease, or myocardium.

When ordering or prescribing

<b>CAFFEINE</b> .....1-10 gr.....	1 00
One every hour as a stimulant to the heart, or as a diuretic.	
<b>CAFFEINE</b> .....1-2 gr.....	1 75
A tablet three times a day as a cardiac stimulant and as an antiperiodic; useful also in nephritis with partial suppression.	
<b>CAFFEINE</b> .....1 gr.....	2 50
One tablet three times a day, as a stimulant to the cardiac and the renal functions; useful in the after treatment of alcohol and opium poisoning.	
<b>CAFFEINE CITRATE</b> .....1-10 gr.....	1 00
One every hour as a stimulant to the heart, or as a diuretic.	
<b>CAFFEINE CITRATE</b> .....1-2 gr.....	1 50
One or two tablets every three hours as a stimulant to the heart and the kidneys, and for bronchial asthma.	
<b>CAFFEINE CITRATE</b> .....1 gr.....	2 00
A tablet three times a day to stimulate heart action and the excretion of urine.	
<b>CAFFEINE CITRATE</b> .....2 grs.....	3 00
One tablet three times a day, as a diuretic in cardiac dropsy; one every two hours for three doses, for headache.	
<b>CAFFEINE COMP.</b> .....	2 25
( Caffeine Citrate....1 gr. / Nitroglycerin...1-200 gr. )	
One tablet three or four times a day, for cardiac dilatation, arteriosclerosis, and as a diuretic.	
<b>CAFFEINE COMP. No. 2</b> .....	2 35
( Caffeine Citrate....1 gr. / Nitroglycerin...1-200 gr. / Fl.Ext.Gelsem...1-2 min. )	
A tablet every three or four hours as a heart tonic and stimulant, to avert the attacks of angina pectoris and lessen their severity, and to promote diuresis.	
<b>CALCIUM SULPHIDE</b> .....1-1000 gr.....	75
A tablet every two or three hours, for the suppurative eruptions of infancy.	
<b>CALCIUM SULPHIDE</b> .....1-100 gr.....	75
One every two hours for discharge from the ear, boils and suppurating skin affections in children; also for conjunctivitis.	
<b>CALCIUM SULPHIDE</b> .....1-50 gr.....	75
A tablet every hour or two for boils, carbuncles, discharges from the ear and diseases of the skin.	
<b>CALCIUM SULPHIDE</b> .....1-20 gr.....	75
<b>Chocolate Coated</b> .....	85
One to three after meals for boils and other suppurating diseases of the cutaneous glands.	
<b>CALCIUM SULPHIDE</b> .....1-10 gr.....	75
<b>Chocolate Coated</b> .....	85
One every two hours for boils, carbuncles and suppurative conditions.	
<b>CALCIUM SULPHIDE</b> .....1-5 gr.....	75
<b>Chocolate Coated</b> .....	85
One after meals for carbuncles, furuncles, and acne; increase gradually until the full systemic effect is reached.	
<b>CALCIUM SULPHIDE</b> .....1-5 gr.....	75
<b>Chocolate Coated</b> .....	85
One to two tablets three times a day for boils, carbuncles, discharges from the ear, and suppurative conditions.	

<b>CALCIUM SULPHIDE</b> .....	1-4 gr.....	75
Chocolate Coated.....		85
One to two tablets every three hours for boils, carbuncles, discharges from the ear, and suppurative conditions.		
<b>CALCIUM SULPHIDE</b> .....	1-2 gr.....	75
Chocolate Coated.....		85
One, three times a day, for boils, carbuncles and suppurative conditions.		
<b>CALCIUM SULPHIDE</b> .....	1 gr.....	1 00
Chocolate Coated.....		1 10
A tablet three or four times daily, for suppurating affections of the skin, carbuncles, boils and otorrhœa.		
<b>CALOMEL</b> .....	1-1000 gr.....	75
Alterative, purgative, antiseptic		
One every hour for an infant, for hepatic and intestinal tonic.		
<b>CALOMEL</b> .....	1-500 gr.....	75
One every hour for an infant, for hepatic and intestinal tonic.		
<b>CALOMEL</b> .....	1-200 gr.....	75
One every hour for an infant, for hepatic and intestinal tonic.		
<b>CALOMEL</b> .....	1-100 gr.....	75
One every two hours to an infant for hepatic and intestinal tonic.		
<b>CALOMEL</b> .....	1-50 gr.....	75
One every hour for a child until effect is produced; for hepatic and intestinal tonic, and corrective in some forms of diarrhœa.		
<b>CALOMEL</b> .....	1-40 gr.....	75
One every hour for a child until effect is produced; for hepatic and intestinal tonic and corrective of diarrhœa.		
<b>CALOMEL</b> .....	1-20 gr.....	75
One every hour for a child until bowels act, as corrective to liver, and laxative.		
<b>CALOMEL</b> .....	1-16 gr.....	75
One every hour for a child until bowels act, as corrective to liver, and laxative.		
<b>CALOMEL</b> .....	1-12 gr.....	75
One every half hour for adults until bowels move or twelve are taken; for hepatic alterative and laxative in biliousness or bilious headache.		
<b>CALOMEL</b> .....	1-10 gr.....	75
One every hour until the bowels move, or until ten have been taken; for biliousness, headache and congestion.		
<b>CALOMEL</b> .....	1-10 gr.....	75
Flavored with Wintergreen.		
One every hour until the bowels move, or until ten have been taken; for biliousness, headache and congestion.		
<b>CALOMEL</b> .....	1-8 gr.....	75
One every hour until the bowels move, or until eight have been taken.		
<b>CALOMEL</b> .....	1-8 gr.....	75
Flavored with Wintergreen.		
One every hour until the bowels move, or until eight have been taken.		
<b>CALOMEL</b> .....	1-6 gr.....	75
One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia or laxative mineral water. For biliousness and torpid liver.		

---

When ordering or prescribing

**CALOMEL** . . . . . 1-6 gr. . . . . 75

Flavored with Wintergreen.

One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia, or laxative mineral water. For biliousness and torpid liver.

**CALOMEL** . . . . . 1-5 gr. . . . . 75

One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia, or laxative mineral water. For biliousness and torpid liver.

**CALOMEL** . . . . . 1-5 gr. . . . . 75

Flavored with Wintergreen.

One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia, or laxative mineral water. For biliousness and torpid liver.

**CALOMEL** . . . . . 1-4 gr. . . . . 75

One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia, or laxative mineral water. For biliousness and torpid liver.

**CALOMEL** . . . . . 1-4 gr. . . . . 75

Flavored with Wintergreen.

One every two hours, until cathartic action is produced, followed by a seidlitz powder, citrate of magnesia, or laxative mineral water. For biliousness and torpid liver.

**CALOMEL** . . . . . 1-2 gr. . . . . 90

One every two hours until effect is produced, followed by a saline cathartic.

**CALOMEL** . . . . . 1-2 gr. . . . . 90

Flavored with Wintergreen.

One every two hours until effect is produced, followed by a saline cathartic.

**CALOMEL** . . . . . 1 gr. . . . . 1 00

One every three hours until bowels move freely, followed by a saline cathartic.

**CALOMEL** . . . . . 1 gr. . . . . 1 00

Flavored with Wintergreen.

One every three hours until bowels move freely, followed by a saline cathartic.

**CALOMEL** . . . . . 2 gr. . . . . 1 30

One or two tablets at night as a purge, taken with a ten grain tablet of bicarbonate of sodium, followed next morning by a saline cathartic.

**CALOMEL** . . . . . 2 gr. . . . . 1 30

Flavored with Wintergreen.

One or two tablets at night as a purge, taken with a ten grain tablet of bicarbonate of sodium, followed next morning by a saline cathartic.

**CALOMEL** . . . . . 5 grs. . . . . 2 00

Alterative.

One at night with a ten grain tablet of bicarbonate of sodium, followed by a dose of citrate of magnesia in the morning for a purge, and as an alterative to liver.

**CALOMEL** . . . . . 10 grs. . . . . 3 00

One at night with a ten grain tablet of bicarbonate of sodium, followed by a dose of citrate of magnesia in the morning for a purge, and as an alterative to liver.

<b>CALOMEL, ALOIN &amp; PODO-PHYLLIN</b> .....	{ Calomel..... 1-10 gr. }	1 10
(Dr. Underwood.)	{ Aloin..... 1-10 gr. }	
Hepatic.	{ Podophyllin.... 1-10 gr. }	
	<b>Chocolate Coated</b> .....	1 20

One to five at night, according to the requirements of the case, in chronic constipation, atony of the lower bowel, and as a hepatic stimulant. Decrease the dose gradually.

<b>CALOMEL COMP.</b> .....	{ Calomel..... 1-50 gr. }	1 10
(Dr. Spencer.)	{ Morph. Sulph.... 1-100 gr. }	
	{ Tartar Emetic... 1-100 gr. }	

One every hour or two until cough becomes easier.

<b>CALOMEL COMP. No. 2</b> .....	{ Calomel..... 1-4 gr. }	75
Alkaline hepatic.	{ Podophyllin.... 1-12 gr. }	
	{ Sodium Bicarb... 1-2 gr. }	
	<b>Chocolate Coated</b> .....	85

One every three hours until biliousness is relieved.

<b>CALOMEL COMP. No. 3</b> .....	{ Calomel..... 1-10 gr. }	75
	{ Sodium Bicarb.... 1 gr. }	
	<b>Chocolate Coated</b> .....	85

One every hour until laxative effect is produced, for biliousness, headache and dull feeling with coated tongue.

<b>CALOMEL COMP. No. 4</b> .....	{ Calomel..... 1 gr. }	1 10
	{ Sodium Bicarb.... 1 gr. }	

One tablet at night and one in the morning, for a week, then omit the morning dose for a week and decrease to one every second or third night; for biliousness, sick headache and light attacks of jaundice.

<b>CALOMEL COMP. No. 5</b> .....	{ Calomel..... 1-2 gr. }	1 00
	{ Podophyllin.... 1-12 gr. }	
	{ Sodium Bicarb... 1-2 gr. }	

One to three at night, followed by a teaspoonful of Epsom salts in cold water in the morning, as a stimulant for hepatic engorgement and to relieve hemorrhoids.

<b>CALOMEL, MODIFIED</b> .....	{ Calomel..... 1-5 gr. }	1 00
(Dr. G. F. Brooks.)	{ Ext. Belladon.... 1-25 gr. }	

One tablet at 7 p. m. repeated every half hour for five doses, for an hepatic stimulant and laxative.

<b>CALOMEL &amp; BISMUTH</b> .....	{ Calomel..... 1-6 gr. }	1 70
Alterative stomachic.	{ Dover's Powder.... 1 gr. }	
	{ Sodium Bicarb.... 2 gr. }	
	{ Bismuth Subnit.... 1 gr. }	

A tablet every three hours, as an alterative in disorders of the gastric functions.

<b>CALOMEL &amp; BISMUTH No. 2</b> .....	{ Calomel..... 1-10 gr. }	75
	{ Bism. Subnit.... 1-10 gr. }	
	{ Sugar Milk..... q. s. }	

One every two or three hours as an alterative in disorders of the gastric functions.

<b>CALOMEL &amp; CODEINE</b> .....	{ Calomel..... 1-10 gr. }	3 75
Sedative hepatic.	{ Codeine..... 1-8 gr. }	

A tablet after meals, for cases of biliousness, with coated tongue and pain referred to the liver.

<b>CALOMEL HYDRASTIS COMP</b> .....	{ Calomel..... 1-12 gr. }	1 00
	{ Nux Vomica.... 1-12 gr. }	
	{ Podophyllin.... 1-12 gr. }	
	{ Calcium Carb.... 1-12 gr. }	
	{ Golden Seal..... 1-12 gr. }	

From one to three tablets with meals and at bedtime, for indigestion with acidity, and torpidity of the liver with constipation.

When ordering or prescribing

<b>CALOMEL &amp; IPECAC</b> . . . . .	{ Calomel . . . . . 1-4 gr. }	75
Modified hepatic and stomachic	{ Ipecac Powd . . . . . 1-4 gr. }	
One or two tablets half an hour after eating, and one at bed-time if necessary, for biliousness and bilious headache, indigestion and vertigo.		
<b>CALOMEL &amp; IPECAC No. 2</b> . . . . .	{ Calomel . . . . . 1-8 gr. }	75
	{ Ipecac Powd . . . . . 1-8 gr. }	
One every two hours for diarrhoea with green movements; for infants. Discontinue after four or five hours.		
<b>CALOMEL &amp; IPECAC No. 4</b> . . . . .	{ Calomel . . . . . 1-100 gr. }	75
(Dr. Boardman Reed.)	{ Ipecac . . . . . 1-100 gr. }	
One, dry, on the tongue every half hour for bilious vomiting.		
<b>CALOMEL &amp; IPECAC COMP.</b> . . . . .	{ Calomel . . . . . 1-4 gr. }	1 35
<b>No 2</b> . . . . .	{ Ipecac Powd . . . . . 1-2 gr. }	
	{ Rhubarb Powd . . . . . 1 gr. }	
One or two after each meal, with an occasional seidlitz powder in the morning; as a tonic laxative for jaundice, hemorrhoids and hepatic torpidity.		
<b>CALOMEL, IPECAC &amp; SODA</b> . . . . .	{ Calomel . . . . . 1-5 gr. }	75
Stomachic.	{ Ipecac Powd . . . . . 1-10 gr. }	
	{ Sodium Bicarb . . . . . 1 gr. }	
One or two tablets after meals, for stomach indigestion with coated tongue and headache.		
<b>CALOMEL, IPECAC &amp; SODA</b> . . . . .	{ Calomel . . . . . 1-8 gr. }	75
<b>No. 2</b> . . . . .	{ Ipecac Powd . . . . . 1-8 gr. }	
	{ Sodium Bicarb . . . . . 1 gr. }	
One to three tablets after eating; for dyspepsia, atony of the stomach and constipation.		
<b>CALOMEL &amp; LEPTANDRIN</b> . . . . .	{ Calomel . . . . . 1-3 gr. }	1 60
(Dr. J. B. Hawes.)	{ Leptandrin . . . . . 1-2 gr. }	
Give one night and morning for two days, and one on the night and morning of the fourth day, for biliousness.		
<b>CALOMEL &amp; MORPHINE CO.</b> . . . . .	{ Calomel . . . . . 1-2 gr. }	3 50
Sedative hepatic.	{ Morphine Sulph . . . . . 1-4 gr. }	
	{ Tartar Emetic . . . . . 1-16 gr. }	
One tablet at night with an occasional saline laxative in the morning; for disorders of the liver accompanied by pain and neurotic symptoms.		
<b>CALOMEL &amp; OPIUM No. 1</b> . . . . .	{ Calomel . . . . . 1-4 gr. }	1 15
(Dr. McEwen.)	{ Opium Powd . . . . . 1-4 gr. }	
One every two hours until a laxative action is produced; for bilious colic, and as an hepatic stimulant.		
<b>CALOMEL &amp; OPIUM No. 2</b> . . . . .	{ Calomel . . . . . 1-8 gr. }	1 00
(Dr. McEwen.)	{ Opium Powd . . . . . 1-8 gr. }	
One every hour until cathartic action is produced; for biliousness, headache and constipation.		
<b>CALOMEL &amp; OPIUM CO. No. 1</b> . . . . .	{ Calomel . . . . . 1-2 gr. }	1 25
	{ Opium Powd . . . . . 1-8 gr. }	
	{ Ipecac Powd . . . . . 1-4 gr. }	
One every three hours, followed next morning by a saline laxative, for biliousness, jaundice and the early stages of cirrhosis.		
<b>CALOMEL &amp; OPIUM CO. No. 2</b> . . . . .	{ Calomel . . . . . 1-4 gr. }	75
	{ Opium Powd . . . . . 1-16 gr. }	
	{ Ipecac Powd . . . . . 1-8 gr. }	
One every two or three hours, for bilious disorders and sick headache, followed by a saline laxative.		
<b>CALOMEL &amp; PODOPHYLLIN</b> . . . . .	{ Calomel . . . . . 1-8 gr. }	75
(Dr. Steenberg.)	{ Podophyllin . . . . . 1-8 gr. }	
One or two at night, or one after meals until the bowels are freely opened; for the habitual constipation of the sedentary.		

<b>CALOMEL &amp; PODOPHYLLIN COMP.</b> .....	{ Calomel.....1 gr. }	1 60
	{ Sodium Bicarb.....2 gr. }	
	{ Podophyllin.....1-4 gr. }	
	{ Ipecac Powd.....1-4 gr. }	

One three times a day, and one or two at night, for habitual constipation, plethora and tendency to piles.

<b>CALOMEL &amp; RHUBARB CO.</b> ... Alterative, laxative.	{ Calomel.....2 1-2 gr. }	2 00
	{ Rhubarb Powd. 2 1-2 gr. }	
	{ Cinnamon Powd....1 gr. }	

One or two at night for a week, then one every second or third night, as an alterative; an occasional one at night as a laxative, for bilious tendency and intestinal dyspepsia.

<b>CALOMEL &amp; SODA</b> .....	{ Calomel.....1-2 gr. }	95
	{ Sodium Bicarb.....3 gr. }	
	<b>Chocolate Coated</b> .....	

One every two hours, for children's diarrhoea with green movements. Discontinue after two or three doses. Two at night for adult hepatic.

<b>CALOMEL &amp; SODA No. 2</b> .....	{ Calomel.....1-4 gr. }	75
	{ Sodium Bicarb.....1 gr. }	

One every hour until laxative effort is produced, for bilious headache with coated tongue.

<b>CAMPBOR COMP</b> .....	{ Camphor Monobr...1 gr. }	1 65
	{ Acetanilid.....2 gr. }	

Analgesic.  
One every three hours when in pain, for headache, muscular rheumatism and neuralgia.

<b>CAMPBOR, CAPSICUM CO.</b> ....	{ Camphor.....1-2 gr. }	2 25
	{ Opium Powd.....1-2 gr. }	
	{ Capsicum Powd....1 gr. }	
	{ Dover's Powder....3 gr. }	

A tablet half an hour after eating, for indigestion with gastralgia, headache and intestinal distress.

<b>CAMPBOR MONOBROMIDE</b> ..1-10 gr.....	75
---	----

Sedative.  
One every hour as a sedative in whooping cough, for a child under seven years.

<b>CAMPBOR MONOBROMIDE</b> ..1-2 gr.....	1 25
--	------

A tablet every hour or two, to lessen the violence of the attack in whooping cough, and quiet restlessness.

<b>CAMPBOR MONOBROMIDE</b> ..1 gr.....	1 50
--	------

One at night, repeated once or twice at intervals of an hour, to quiet nerves and induce sleep.

<b>CAMPBOR MONOBROMIDE</b> ..2 grs.....	1 85
---	------

A tablet as required to diminish the intensity of asthmatic paroxysms; also as an hypnotic, and to control delirium.

<b>CANNABIS COMP. No. 1</b> ..... (Dr. Patterson.) Anodyne.	{ Camphor.....1 gr. }	1 00
	{ Ext. Can. Ind.....1-4 gr. }	
	{ Ext. Hyoscyam...1-2 gr. }	
	{ Sugar of Milk.....q. s. }	

**Chocolate Coated**..... 2 00

Give one every two or three hours, or as often as every hour, as long as necessary to relieve pain of ovarian neuralgia and of the genito-urinary region.

<b>CANNABIS COMP. No. 2</b> .....	{ Ext. Can. Ind.....1-8 gr. }	1 70
	{ Morph. Sulph....1-20 gr. }	
	{ Strych. Sulph....1-120 gr. }	
	{ Acid Arsenous...1-80 gr. }	
	{ Ext. Aconite.....1-8 gr. }	

One tablet as required, for the pain of neuralgias or inflammations; repeat in an hour if necessary. Stop at the third dose.

When ordering or prescribing

<b>CANNABIS COMP. No. 3</b> .....	{ Fl. Ext. Can. Ind. . . 1 min. }	<b>I 60</b>
(Dr. Sadler.)	{ Fl. Ext. Gelsem. . . 2 min. }	
One to three tablets every three to six hours, as long as pain exists. Indicated in trigeminal neuralgias and nervous disturbances arising from uterine irregularities.		
<b>CANNABIS IND. EXT.</b> .....	1-20 gr. ....	<b>75</b>
Hypnotic and anodyne.		
One every two hours, to control the twitchings and restlessness of chorea, and to induce sleep.		
<b>CANNABIS IND. EXT.</b> .....	1-10 gr. ....	<b>I 00</b>
A tablet after meals and one at night, for hypochondriasis and anæmic headache, and to control intense pruritus.		
<b>CANNABIS IND. EXT.</b> .....	1-4 gr. ....	<b>I 50</b>
Give one every three or four hours, for depression of spirits, melancholia, and some forms of headache, and to lessen the pain of rheumatism.		
<b>CANNABIS &amp; CODEINE</b> .....	{ Tr. Cannabis Ind. . . 1 min. }	<b>I 60</b>
Sedative.	{ Codeine. .... 1-25 gr. }	
One to four tablets at intervals of half an hour, to quiet the nerves and produce sleep. In chorea, one tablet after meals.		
<b>CANTHARIDES</b> .....	1-100 gr. ....	<b>75</b>
Stimulant diuretic.		
One tablet every hour, for congestion of urinary tract; especially to avert threatened suppression of urine. Allow plenty of drinking water.		
<b>CANTHARIDES</b> .....	1-50 gr. ....	<b>75</b>
A tablet every hour for congestion of urinary tract. Administer drinking water freely.		
<b>CANTHARIDES</b> .....	1-20 gr. ....	<b>75</b>
A tablet every three hours in renal congestion, and to prevent suppression of urine. Give drinking water freely.		
<b>CANTHARIDES</b> .....	1-10 gr. ....	<b>75</b>
Give one every three or four hours, in renal congestion with threatened anuria, with free use of drinking water.		
<b>CAPSICUM</b> .....	1-100 gr. ....	<b>75</b>
Stomachic.		
One after meals for children, as an aid to digestion; may be repeated every hour.		
<b>CAPSICUM</b> .....	1-50 gr. ....	<b>75</b>
One after meals for children, as an aid to digestion; may be repeated every hour.		
<b>CAPSICUM</b> .....	1-20 gr. ....	<b>75</b>
One to three after meals, repeat one every hour afterwards, as an aid to digestion, and to relieve epigastric oppression after eating.		
<b>CAPSICUM</b> .....	1-10 gr. ....	<b>75</b>
One or two after meals, as a tonic and aid to the digestive organs and a relief to epigastric fulness and oppression after eating. May be repeated for two doses at half hour intervals.		
<b>CAPSICUM</b> .....	1-4 gr. ....	<b>75</b>
One or two after meals, repeated in an hour if necessary as a tonic and aid to the digestive organs, and a relief to epigastric fulness and oppression after eating.		
<b>CAPSICUM</b> .....	1-2 gr. ....	<b>75</b>
One after each meal, repeated in an hour if necessary as a tonic and aid to the digestive organs, and a relief to epigastric fulness and oppression after eating; also good in the morning nausea of dipsomaniacs.		

<b>CAPSICUM COMP.</b> .....	{ Capsicum.....1-10 gr. }	75
(Dr. Lindsley.)	{ Nux Vomica.....1-4 gr. }	85
Stimulant stomachic.	Chocolate Coated.....	
One or two tablets half an hour after meals; if necessary repeat at half hour intervals for three doses, for stomach indigestion.		
<b>CARBO VEG.</b> .....	{ .....Carbo Veg..... }	75
(Fraser.)	{ .....Saccharin..... }	
Antacid.		
Use freely as required, in cases of eructation, heartburn and flatulence.		
<b>CARBO VEG.</b> .....	1-100 gr.....	75
Give one every half hour to a infant with dyspepsia and wind colic.		
<b>CARBO VEG.</b> .....	1-10 gr.....	75
Give one to three after meals, for dyspepsia and flatulence of children after errors in diet. Give an occasional saline laxative.		
<b>CARBO VEG.</b> .....	1-2 gr.....	75
A tablet or two as required, for heartburn, eructation and flatulence. Take a seidlitz powder in the morning every few days.		
<b>CARDIAC</b> .....	{ Sparteine Sulph...1-10 gr. }	3 40
(Dr. Waldstein.)	{ Tr. Strophanthus...3 min. }	
	{ Caffeine Cit.....1-2 gr. }	
	{ Codeine...1-20 gr. }	
Give a tablet every three hours, to control rapid and tumultuous heart action, irregular pulse, difficult breathing and tendency to œdema. Avoid overloading the stomach.		
<b>CARDIAC No. 2</b> .....	{ Ext. Valerian.....2 gr. }	3 75
(Dr. Heineman.)	{ Ext. Digitalis.....1-2 gr. }	
	{ Ergotin.....2 gr. }	
One, three or four times a day, for irritable, feeble heart.		
<b>CARDIAC TONIC</b> .....	{ Strych. Arsen...1-100 gr. }	2 25
(Dr. Bartley.)	{ Sparteine Sulph...1-8 gr. }	
	{ Quinine Sulph...1-2 gr. }	
One or two tablets three or four times daily, for senile heart and fatty degeneration of the heart.		
<b>CARDIANE</b> .....	{ Digitalia.....1-120 gr. }	2 75
(Dr. Pallen.)	{ Tr. Strophanthus.2 min. }	
	{ Tr. Cactus.....1 min. }	
	{ Sparteine Sulph...1-20 gr. }	
A tablet three times a day, to regulate the action of the heart, diminish undue frequency and relieve dyspnœa; for valvular cardiac disease and begining dilatation.		
<b>CASCARA COMP</b> .....	{ Ext. Cascara.....1-2 gr. }	1 60
	{ Aloin.....1-8 gr. }	
	{ Podophyllin.....1-10 gr. }	
	{ Oil Peppermint.1-10 min. }	
	Chocolate Coated.....	1 70
One or two at night, followed by one after each meal if necessary, for a laxative.		
<b>CASCARA COMP. No. 2</b> .....	{ Ext. Cascara Sag...2 gr. }	2 50
	{ Podophyllin.....1-8 gr. }	
	{ Ext. Belladonna...1-16 gr. }	
	Plain or Chocolate Coated.....	
Two tablets at night, as a mild tonic cathartic.		
<b>CASCARA COMP. No. 3</b> .....	{ Cascarin.....1-4 gr. }	1 05
(Dr. Hinckle.)	{ Aloin.....1-4 gr. }	
	{ Podophyllin.....1-6 gr. }	
	{ Ext. Belladonna...1-8 gr. }	
	{ Strych. Sulph...1-60 gr. }	
	{ Gingerine...1-8 gr. }	
	Chocolate Coated.....	1 75
One or two at night, followed by one in the morning and at noon, if necessary, for laxative.		

When ordering or prescribing

<b>CASCARA SAGRADA Fl. Ext.</b> .....5 min.....		1 75
Laxative.	<b>Plain or Chocolate Coated.</b>	
One or two at night, or every three hours during the day, increasing intervals as constipation is relieved.		
<b>CASCARIN COMP.</b> .....	{ Aloin.....1-4 gr. } Podophyllin.....1-4 gr. }	1 40
Tonic, hepatic and laxative.	{ Cascarin.....1-4 gr. }	
	<b>Chocolate Coated.</b> .....	1 50
A tablet after meals, with free use of drinking water, for habitual and atonic constipation.		
<b>CASCARIN COMP No. 2.</b> .....	{ Ext. Cascara Sag.1-2 gr. } Aloin.....1-8 gr. } Podophyllin.....1-10 gr. }	1 85
(Dr. Gilbert.)	{ Oil Peppermint..1-10 min. } Oil Cajeput.....1-2 min. }	
	<b>Chocolate Coated.</b> .....	1 95
One after each meal, until they act freely; then one at night as necessary for habitual constipation accompanied with flatulence.		
	{ Cascarin.....1-2 gr. } Podophyllin.....1-12 gr. } Euonymin.....1-4 gr. }	
<b>CASCARIN COMP. No. 3.</b> .....	{ Strychnine.....1-80 gr. } Aloin.....1-10 gr. }	
(Dr. Martin, Clinton, Iowa.)	{ Ext. Belladonna..1-8 gr. } Oil Black Pepper..1-4 gr. }	
	Gingerine.....1-16 gr. }	
	<b>Plain or Chocolate Coated</b>	2 00
One, rarely two, on going to bed, for hepatic and intestinal tonic and aperient.		
<b>CASCARIN TABLETS</b> .....	{ Salicin.....3 gr. } { Cascarin.....2 gr. }	4 25
(Dr. Waldstein.)		
Tonic cathartic.		
One at night, for loss of tone in the intestine with consequent habitual constipation.		
<b>CATARRHAL</b> .....	{ Tr. Aconite.....1-2 min. } Ipecac Powd.....1-8 gr. }	1 10
(Dr. F. W. Koehler.)	{ Potass. Iodide...1-10 gr. }	
One every half hour for three doses, then every hour, to start the secretions dried up by an inflammation of the throat, in its first stages.		
	{ Leptandrin.....1-32 gr. } Aloin.....1-16 gr. } Podophyllin.....1-6 gr. }	
<b>CATHARTIC No. 1.</b> .....	{ Gamboge.....1-64 gr. } Ext. Hyoscyam..1-16 gr. }	1 00
	{ Oleores. Capsic.1-128 gr. } Oil Pepperm't.1-128 min. }	
	<b>Chocolate Coated.</b> .....	1 10
One to three tablets at night.		
	{ Aloin.....1-10 gr. } Ext. Colocy. Co..1-10 gr. } Ext. Nux Vom...1-10 gr. }	
<b>CATHARTIC No. 2.</b> .....	{ Podophyllin.....1-10 gr. } Croton Oil.....1-30 gr. }	1 25
	{ Oleores. Capsic.1-256 gr. } Oil Pepperm't.1-128 min. }	
	<b>Chocolate Coated.</b> .....	1 35
One tablet every two hours until bowels move, or one to three at night, followed in the morning by another if necessary.		

<b>CATHARTIC No. 3</b> .....	{ Scammony Res. .2 1-2 gr. }	4 00
(Dr. Russell.)	{ Gray Powder....2 1-2 gr. }	

Give one to four at night, according to age, as a cathartic when an hepatic is indicated.

<b>CATHARTIC No. 4</b> .....	{ Pil. Rhubarb Comp. .4 gr. }	2 25
	{ Blue Mass. ....3 gr. }	

Two at night and one every seventh night afterwards, until liver and bowels act regularly.

<b>CATHARTIC No. 5</b> .....	{ Aloes Powd.....1 1-8 gr. }	1 35
	{ Gamboge Powd..3-16 gr. }	
	{ Podophyllin.....1-6 gr. }	
	{ Capsicum Powd..1-10 gr. }	
	{ Croton Oil.....1-10 gr. }	
	{ Ext. Hyoscyamus..1-8 gr. }	
	<b>Chocolate Coated</b>	1 45

One to three at night for active purge.

<b>CATHARTIC COMP. Active</b> ....	{ Aloes Purified...1 1-8 gr. }	1 50
	{ Gamboge.....3-16 gr. }	
	{ Podophyllin.....1-8 gr. }	
	{ Capsicum Powd..1-12 gr. }	
	{ Croton Oil.....1-50 gr. }	
	<b>Chocolate Coated</b> .....	1 60

One to three tablets at night for a cathartic.

<b>CATHARTIC COMP. Improved</b>	{ Ext. Colocy. Co. ....1 gr. }	2 75
	{ Ext. Jalap.....1-2 gr. }	
	{ Podophyllin.....1-4 gr. }	
	{ Ext. Gentian.....1-2 gr. }	
	{ Leptandrin.....1-4 gr. }	
	{ Ext. Hyoscyam...1-4 gr. }	
	{ Oil Peppermint.....q. s. }	
	<b>Plain or Chocolate Coated</b>	

One or two at bedtime, as a laxative; two to three as a cathartic.

<b>CATHARTIC COMP. U. S. P.</b> ....	{ Ext. Colocy. Co. .1 1-4 gr. }	2 50
	{ Calomel.....1 gr. }	
	{ Ext. Jalap.....1-2 gr. }	
	{ Gamboge Powd..1-4 gr. }	
	<b>Plain or Chocolate Coated</b>	

One at bedtime as a laxative, two as a cathartic, three as a purge; for indigestion and biliousness.

<b>CATHARTIC CO. (Pill U. S. P.)</b> ..1-10 gr.....	75
---	----

Give one every half hour until the bowels move, or six at night, according to the requirements of the case.

<b>CATHARTIC CO. (Pill U. S. P.)</b> ..1-5 gr.....	75
<b>Chocolate Coated</b> .....	85

From one to four tablets, in divided doses through the day or at bed time; as a tonic and hepatic laxative.

<b>CATHARTIC COMP. (Veg.)</b> .....1-10 gr.....	75
---	----

One every hour up to four or five doses as necessary, to procure free action of the bowels; then one every night for a time.

<b>CATHARTIC COMP. (Veg.)</b> .....1-5 gr.....	75
--	----

Two or three during the day and two at night, with another in the morning if necessary; after the bowels act freely, diminish to one every night.

When ordering or prescribing

<b>CATHARTIC TONIC</b> (Dr. Bovee.)	(Strychnine Sulph. 1-40 gr.)	
	Cascarin.....1-3 gr.	
	Aloin.....1-3 gr.	
	Res. Podophyllin...1-5 gr.	
	Ext. Belladonna...1-4 gr.	
	<b>Plain or Chocolate Coated</b>	2 00

One at night, followed by one after each meal next day, if necessary; for a tonic cathartic in biliousness and constipation.

<b>CAULOPHYLLIN</b> .....1-10 gr.....	1 10
Antispasmodic and diuretic.	

Give from one to three every hour, to relieve spasm of the sphincter vesicæ and post-partum pains, and as a diuretic.

<b>CERIUM OXALATE</b> .....1-10 gr.....	75
Stomachic sedative.	

One every fifteen minutes diffused in water on the end of a spoon for nausea or infants. When nausea stops, give one, half an hour before each feeding, for two or three meals.

<b>CERIUM OXALATE</b> .....1 gr.....	1 00
--------------------------------------	------

One every twenty minutes for adult, every half hour for grown child; for nausea. When stomach is settled, one every two hours with hot liquid food as long as necessary.

<b>CERIUM OXALATE</b> .....2 grs.....	1 40
---------------------------------------	------

One every hour for four or five doses, for sick stomach and nausea.

<b>CERIUM OXALATE</b> .....5 grs.....	1 75
---------------------------------------	------

One to two tablets three or four times a day, for vomiting in pregnancy.

<b>CERIUM OXALATE COMP.</b> .....	Cerium Oxalate....1 gr.	1 15
(Dr. Guernsey.)	Ipecac Powd....1-20 gr.	

One every hour until nausea stops, followed by one every hour with hot beef tea in small quantities.

<b>CHALK</b> .....	Chalk Prepared....1 gr.	75
Antacid.	Oil Cinnamon. 1-20 min.	

A tablet every half hour until relieved, for a child with dyspepsia and gastric distress; for a baby, give one with each feeding.

<b>CHALK MIXTURE</b> .....1 teaspoonful.....	75
--	----

One tablet every two or three hours, for the summer diarrhoea of children. It is best to precede this treatment with a single dose of castor oil.

<b>CHARCOAL</b> .....2 grs.....	75
Absorbent.	

A tablet as needed to control nausea and sea-sickness, and to correct flatulence.

<b>CHARCOAL</b> ......5 grs.....	75
----------------------------------	----

Give one every three hours or oftener in conditions of gastric dyspepsia with eructations.

<b>CHARCOAL</b> .....10 grs.....	75
----------------------------------	----

A tablet after each meal for indigestion with the production of gas, nausea, and palpitation.

<b>CHLORALAMID, Schering's</b> ...5 grs.....	13 00
--	-------

Per 100.....1 30

A tablet every three hours in chorea and to avert epileptic seizures, and as an hypnotic to replace chloral and sulphonal.

<b>CHLORALAMID, Schering's</b> ...10 grs.....	25 00
---	-------

Per 100.....2 50

One tablet repeated in an hour to produce sleep; for insomnia of senility, neurasthenia and alcoholism.

<b>CHLORALAMID</b> , Schering's...15 grs. ....	37 00
Per 100.....	3 75

A tablet as necessary to avert epileptic and choreic attacks; to produce sleep in old age, hysteria, cardiac disease and morphinism.

<b>CHLORODYNE</b> .....	{ Morphine Mur....1-6 gr. Ext. Cannabis Ind.1-4 gr. Nitroglycerin...1-300 gr. Ext. Hyoscyam....1-2 gr. Oleoresin Capsic.1-10 gr. Oil Peppermint.1-10 min.	
	<b>Plain or Chocolate Coated</b>	3 75

One, and repeat in an hour if necessary for pain. After taking three doses, stop three hours before resuming. For diarrhœa, one every three hours until relieved.

<b>CHLORODYNE No. 2</b> .....	{ Morphine Mur....1-12 gr. Ext. Cannabis Ind.1-8 gr. Nitroglycerin...1-600 gr. Ext. Hyoscyamus.1-4 gr. Oleoresin Capsic.1-20 gr. Oil Peppermint.1-20 min.	
	<b>Plain or Chocolate Coated</b>	2 00

One, and repeat in an hour if necessary for pain. After taking three doses, stop three hours before resuming. For diarrhœa, one every three hours until relieved.

<b>CHLOROSIS</b> .....	{ Quinine Sulph.....1 gr. Iron Powd.....1 1-2 gr. Ext. Ergot.....1 gr. Ext. Nux Vom....1-5 gr.	
(Dr. Bishop.)		
	<b>Plain or Chocolate Coated</b>	2 50

One tablet, three times a day, in anæmia, chlorosis and amenorrhœa.

<b>CHLOROSIS No. 2</b> .....	{ Iron Protochlor....1-4 gr. Merc. Bichlor....1-120 gr. Arsenic Chlor...1-200 gr. Quinine Chlor....1-8 gr.	
		1 40

One or two tablets after each meal where a vigorous iron tonic is required, and during convalescence.

<b>CHOCOLATE LAXATIVE</b> .....	{ Ext. Cascara Sag..1-4 gr. Aloin.....1-40 gr. Podophyllin....1-40 gr. Ext. Tamarinds....2 gr. Sweet Chocolate...9 gr.	
		2 00

Give one to four according to age of patient, as an agreeable laxative for children.

<b>CHOLAGOGUE</b> .....	{ Calomel.....1 gr. Ipecac Powd....1-8 gr. Acid Arsenous...1-15 gr. Corros. Sublim...1-15 gr.	
(Dr. Porter.)		1 25

One every three hours until bowels move more freely; then one every second or third night, or every week, as necessary.

<b>CHOLAGOGUE No. 2</b> .....	{ Calomel.....1-10 gr. Ipecac Powd....1-10 gr. Sodium Bicarb....2 gr.	
(Dr. Kennedy.)		1 00

One every two hours, as a mild cholagogue for adults. For constipation of infants, one with as much sugar of milk as will lie on a silver ten-cent piece.

<b>CHOLERA</b> .....	{ Opium Powd.....1-2 gr. Asafœtida.....1 gr. Black Pepper....1 1-2 gr.	
		1 75

Two every two hours with hot water or tea, for early stages of diarrhœa, particularly where cholera is suspected.

When ordering or prescribing

<b>CHOLERA INFANTUM</b> .....	{ Zinc Sulphocarb. 1-20 gr. Salol.....1-10 gr. Bismuth Subnit. ....1-2 gr. Calomel.....1-60 gr. Lactopeptin.....1-2 gr. }	3 00
(Dr. Hillkowitz.)		

One every ten minutes until five are given, then one every hour; as a remedy for cholera infantum and summer diarrhoea in children from six to twelve months old.

<b>CHOLERA INFANTUM No. 2.</b> .....	{ Zinc Sulphocarb. 1-20 gr. Salol.....1-10 gr. Bismuth Subnit. ....1-2 gr. Calomel.....1-60 gr. Pepsin Pure.....1-4 gr. }	1 50
(Dr. Fenwick.)		

One tablet every ten minutes the first hour, then one every hour until relieved.

<b>CHOLERA MORBUS, Bilious.</b> .....	{ Bismuth Subnit. ....3 gr. Calomel.....1-8 gr. Salol.....1 gr. }	3 50
(Dr. Hitchcock.)		

If there is an irritating substance in the bowels, give a seidlitz powder first, then a tablet after each bad passage; not oftener than every hour, nor more than ten in twenty-four hours.

<b>CIMICIFUGA COMP.</b> .....	{ Tr. Cimicifuga....1 min. } { Tr. Gelsemium....1 min. }	2 00
-------------------------------	---	------

One every one to two hours for menstrual headaches and ovarian congestion.

<b>CIMICIFUGA FL. EXT.</b> .....1 min. ....	1 15
---	------

Anti-rheumatic and cardiac tonic.

Give one to three tablets every two hours, for headache dependent upon pelvic derangement, sex hypochondriasis, and as a cardiac tonic in fatty heart and rheumatism.

<b>CINCHONIDIA SALICYLATE.</b> 1 gr.....	2 00
--	------

One every hour, for rheumatism and rheumatic pharyngitis.

<b>CINCHONIDIA SALICYLATE.</b> 2 grs.....	3 00
---	------

One to three tablets, three times a day according to age, for rheumatism and rheumatic pharyngitis.

<b>CINCHONIDIA SALICYLATE.</b> 2 1-2 grs.....	3 50
---	------

Plain or Chocolate Coated

One every two hours for rheumatism and rheumatic pharyngitis.

<b>CINCHONIDIA SALICYLATE.</b> 3 grs.....	4 00
---	------

One every two hours for rheumatism and rheumatic pharyngitis.

<b>CINCHONIDIA SALICYLATE.</b> 5 grs.....	5 00
---	------

One, three or four times a day in rheumatism and rheumatic pharyngitis.

<b>COCAINE COMP.</b> .....	{ Potass. Chlorate....1 gr. } { Cocaine Mur...1-100 gr. }	1 25
(Dr. F. A. Burrall.)		

One on the tongue every half hour for pharyngitis and colds.

<b>COCAINE HYDROCHLOR</b> ....1-100 gr.....	1 00
---	------

<b>COCAINE HYDROCHLOR</b> ....1-50 gr.....	1 35
--	------

<b>COCAINE HYDROCHLOR</b> .....1-20 gr.....	1 90
---	------

<b>COCAINE HYDROCHLOR</b> .....1-10 gr.....	2 75
---	------

<b>COCAINE HYDROCHLOR</b> ....1-8 gr.....	3 25
---	------

<b>COCAINE HYDROCHLOR</b> ....1-6 gr.....	4 00
---	------

COCAINE HYDROCHLOR.....	1-5 gr.....	4 50
COCAINE HYDROCHLOR.....	1-4 gr.....	5 25
COCAINE HYDROCHLOR.....	1-8 grs.....	20 00
	Per 100.....	2 00

A tablet dissolved in a drachm of distilled water makes a two per cent. solution.

COCAINE HYDROCHLOR.....	2 1-4 grs.....	40 00
	Per 100.....	4 00

A tablet dissolved in one drachm of distilled water makes a four per cent. solution.

COCAINE THROAT.....	<div style="display: inline-block; vertical-align: middle;">           Cocaine Mur.....1-12 gr.            Cubebs Powd.....1 gr.            Acid Benzoic.....1-2 gr.            Potass. Chlorate.....2 gr.            Licorice Powd.....q. s.         </div>	3 00
---------------------	--	------

One on the tongue every two or three hours for chronic cough with tickling and irritation of the throat.

CODEINE.....	1-10 gr.....	2 75
CODEINE.....	1-8 gr.....	3 25
CODEINE.....	1-5 gr.....	4 50
CODEINE.....	1-4 gr.....	5 25
CODEINE.....	1-2 gr.....	10 50
CODEINE.....	1 gr.....	20 00

CODEINE COMP.....	<div style="display: inline-block; vertical-align: middle;">           Codeine.....1-3 gr.            Camphor.....1 gr.            Prepared Chalk.....5 gr.         </div>	7 00
(Dr. M. J. Asch.)	Chocolate Coated only	

One every two or three hours as required, for diarrhoea and intestinal pains.

CODEINE PHOSPHATE.....	1-4 gr.....	5 75
CODEINE PHOSPHATE.....	1-2 gr.....	10 50
CODEINE SULPHATE.....	1-5 gr.....	3 75
CODEINE SULPHATE.....	1-4 gr.....	4 50
CODEINE SULPHATE.....	1-2 gr.....	8 25
COLCHICIN.....	1-100 gr.....	1 20

Anti-rheumatic.

One every three hours until a cathartic effect is produced. For rheumatism, gout and nephritis.

COLCHICIN.....	1-50 gr.....	1 75
----------------	--------------	------

A tablet three times a day, for neuralgia, rheumatism, and chronic inflammation of the kidneys.

COLD, CATARRHAL.....	<div style="display: inline-block; vertical-align: middle;">           Acetanilid.....3 gr.            Camphor Powd.....1-2 gr.            Codeine.....1-6 gr.            Ext. Nux Vom.....1-6 gr.         </div>	4 75
(Dr. Newman.)	Chocolate Coated only	

One every two hours for catarrhal cold.

COLD No. 1.....	<div style="display: inline-block; vertical-align: middle;">           Morph. Sulph.....1-32 gr.            Tartar Emetic.....1-32 gr.            Ext. Licorice.....1-10 gr.            Oil Gaultheria.....q.s.         </div>	1 35
-----------------	--	------

One every two or three hours, for cough with bronchial irritation; lengthen the intervals to one every four hours if nausea occurs.

When ordering or prescribing

COLD No. 2..... (Dr. Satterlee.)	Antim. Sul. Aur. .1-12 gr.	1 00
	Ext. Conium.....1-12 gr.	
	Ipecac Powd.....1-6 gr.	
	Potass. Nitrate...1-6 gr.	
	Ammon. Mur.....1-6 gr.	

Two after each meal for cold of adults.

COLD No. 3..... (Dr. Gage.)	Quinine Sulphate...2 gr.	3 60
	Dover's Powder...2 gr.	
	Capsicum Powd...1-4 gr.	
	Tr. Aconite.....1 min.	
Plain or Chocolate Coated		

One tablet every three or four hours until effect is produced; give two or three at bedtime if necessary.

COLD No. 3, Modified..... (Dr. T. M. Lippitt.)	Quinine Sulphate...2 gr.	3 75
	Dover's Powder...2 gr.	
	Capsicum Powd...1-4 gr.	
	Tr. Aconite.....1 min.	
	Calomel.....1-4 gr.	
Plain or Chocolate Coated		

One tablet every three or four hours until effect is produced; give two or three at bedtime if necessary.

COLD No. 4..... (Dr. H. L. Coit.)	Aconite.....1-10 gr.	75
	Camphor.....1-10 gr.	
	Opium Powd.....1-10 gr.	
	Potass. Nitrate...1-10 gr.	
	Chocolate Coated.....	
		85

Give adults ten tablets in water at bedtime, or one every hour during the day for coryza. This treatment may be supplemented with tonic doses of quinine.

COLD No. 5. Early Stages.....	Tr. Aconite Rad...1 min.	1 40
	Atropine Sulph...1-600 gr.	
	Morph. Sulph.....1-36 gr.	
	Hydrarg. Chl. Mit. 1-6 gr.	

One every hour for three doses, then every two hours for three more. Take for one day after secretion and weeping from nostrils is stopped.

COLD No. 6..... (Dr. Curtis.)	Camphor.....1-4 gr.	1 15
	Atropine Sulph...1-500 gr.	
	Strych. Sulph...1-100 gr.	
	Aloin.....1-100 gr.	
	Tr. Aconite.....1-2 min.	
Chocolate Coated only		

One every hour, for fresh cold, until mouth becomes dry; then one every two or three hours.

COLD No. 7..... (Dr. Brooks.)	Calomel.....1-5 gr.	1 35
	Podophyllin.....1-25 gr.	
	Ext. Belladonna 1-25 gr.	
	Sodium Bicarb.....2 gr.	

One every half hour after seven in the evening while awake, followed next morning by a seidlitz powder, for breaking up colds.

COLD No. 8..... (Dr. Koyle.)	Fl. Ext. Gelsem...1-2 min.	1 00
	Oil Gaultheria...1-20 min.	

One every half hour, for acute congestive colds.

COLIC..... (Dr. Satterlee.)	Morph. Sulph....1-80 gr.	1 60
	Bismuth Subnit.....1 gr.	
	Calomel.....1-40 gr.	

One every fifteen minutes until four are taken; rest an hour and repeat if necessary in bad cases; give an adult four at the first dose, and four every hour afterwards, as long as necessary.

<b>COLIC No. 2, INFANTILE</b> .....	<div>Paregoric.....2 min.</div> <div>Sodium Bicarb.....1 gr.</div> <div>Oil Fennel....1-10 min.</div>	1 10
Dissolve eight in eight teaspoonfuls of hot water; give ten to twenty drops every ten or fifteen minutes.		
<b>CONIUM COMP.</b> .....	<div>Ext. Conium.....1-30 gr.</div> <div>Cubeb's Powd....1-10 gr.</div> <div>Ext. Licorice....1-10 gr.</div>	75
Expectorant.		
One every half hour or every hour, for cough with tickling in the throat.		
<b>CONIUM COMP. with Codeine.</b> .....	<div>Ext. Conium.....1-30 gr.</div> <div>Cubeb's Powd....1-10 gr.</div> <div>Ext. Licorice....1-10 gr.</div> <div>Codeine.....1-25 gr.</div>	1 15
Anodyne expectorant.		
One every hour for tickling in the throat, with persistent cough.		
<b>CONSTIPATION. Aloin, Strych-</b>	<div>Aloin.....1-5 gr.</div> <div>Ext. Belladonna.1-16 gr.</div> <div>Strychnine Sulph.1-60 gr.</div> <div>Pepsin.....1-4 gr.</div> <div>Ipecac Powd....1-15 gr.</div>	1 50
<b>nine Comp. with Pepsin</b> .....		
	Chocolate Coated.....	1 60
One before supper and another at bedtime, followed by one every two hours next day, if necessary; as a tonic to the liver and secretions.		
<b>CONSTIPATION. CHRONIC</b> ...	<div>Ext. Colocy. Co.....2 gr.</div> <div>Ext. Nux Vomica.1-4 gr.</div> <div>Ext. Belladonna..1-4 gr.</div> <div>Ext. Gentian.....1 gr.</div> <div>Oil Caraway....1-2 min.</div>	4 00
(Dr. Feltwell.)		
One or two at night for habitual constipation; gradually lessen the dose, or increase the interval as the constipation is relieved.		
<b>CONVALLARIA MAJAL. Fl.Ex</b> 1 min.....		1 25
Cardiac tonic.		
Two or three tablets after meals to diminish the frequency and increase the power of the heart; to remove dropsy, and to replace digitalis where that is unsafe.		
<b>CONVALLAMARIN</b> .....1-100 gr.....		1 75
One tablet three times a day, increased with great caution; for cardiac weakness and rapidity, and dropsy.		
<b>COPAIBA BALSAM</b> .....1-100 gr.....		75
Diuretic.		
One to three tablets every two hours, increasing slowly according to the severity of the case; for vesical and urethral inflammation, and to increase the flow of urine.		
<b>COPPER ARSENITE</b> .....1-5000 gr.....		75
Put five tablets in a glass of water, and give a teaspoonful every half hour, for diarrhœa in nursing infants.		
<b>COPPER ARSENITE</b> .....1-1000 gr.....		75
A tablet in a little water every two hours, for the summer diarrhœa of children.		
<b>COPPER ARSENITE</b> .....1-500 gr.....		75
Take one every hour or two until the discharges become natural.		
<b>COPPER ARSENITE</b> .....1-200 gr.....		75
A tablet three times a day for diarrhœa in children.		
<b>COPPER ARSENITE</b> .....1-100 gr.....		75
One tablet in one-half glass of water. One teaspoonful of this every half-hour until relieved of the diarrhœa.		

When ordering or prescribing

**COPPER ARSENITE**.....1-50 gr..... 75

One in one-half glass of water. One teaspoonful of this every half-hour until relieved of the diarrhœa.

**CORRECTIVE, INFANTS** ..... { Calomel .....1-20 gr. } 1 50  
(Dr. Ives.) { Ipecac Powd.....1-50 gr. }  
{ Sodium Bicarb.....1-2 gr. }  
{ Bismuth Subnit.....1 gr. }

A tablet, crushed and mixed with an equal bulk of sugar and placed on the child's tongue every one or two hours, to correct morbid conditions of the alimentary tract.

**CORRECTIVE, INFANTS No. 2** ..... { Calomel.....1-20 gr. } 1 50  
(Dr. Ives.) { Ipecac Powd.....1-50 gr. }  
{ Sodium Bicarb.....1-2 gr. }  
{ Bismuth Subnit.....1 gr. }  
{ Oil Anise ..... 1-20 min. }

A tablet, crushed and mixed with an equal bulk of sugar and placed on the child's tongue every one or two hours, to correct morbid conditions of the alimentary tract.

**CORRECTIVE TABLETS**..... { Rhubarb Powd...1-2 gr. } 2 25  
(Dr. W. B. Culbert) { Ipecac Powd.....1-12 gr. }  
{ Sodium Bicarb.....1 gr. }  
{ Sodium Salicylate...3 gr. }  
{ Ext. Nux Vom...1-20 gr. }  
{ Ext. Cascara Sag.1 1-2 gr. }  
{ Ess. Ment. Pip..1-2 min. }  
**Plain or Chocolate Coated**

One or two tablets, with plenty of water three times a day before eating, for torpid liver or disordered digestion. For laxative, one or two at night.

**CORROSIVE SUBLIMATE** ....1-10,000 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-5000 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-1000 gr..... 75  
**CORROSIVESUBLIMATE** ....1-500 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-200 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-100 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-60 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-50 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-40 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-32 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-30 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-20 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-16 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-12 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-10 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-8 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-4 gr..... 75  
**CORROSIVE SUBLIMATE** ....1-2 gr..... 90  
**CORROSIVE SUBLIMATE** ....1 gr..... 1 00

A tablet dissolved in four tablespoonfuls of water makes a solution of about 1 to 1,000; for external use.

**CORROSIVE SUB. (colored blue)** 1 gr..... 1 00  
**Antiseptic.**

A tablet dissolved in four tablespoonfuls of water makes a solution of about 1 to 1,000; for external use.

<b>CORROSIVE SUBLIMATE</b>	(Corrosive Sub.....1 gr.)	1 00
<b>SOLUBLE.</b> (For Hair Wash.)	(Tartaric Acid.....1 gr.)	

Dissolve a tablet in eight tablespoonfuls of water, and apply twice a week for dandruff.

<b>CORROSIVE SUBLIM. COMP..</b>	{ Corrosive Sub....1-32 gr. }	75
(Dr. Cushman.)	{ Ipecac Powd.....1-8 gr. }	
Alterative.		

One tablet after meals as an alimentary stimulant and alterative, in chronic dyspepsia, rheumatism and skin diseases.

<b>CORROSIVE SUBLIMATE &amp;</b>	(Corrosive Sub....1 3-4 gr.)	1 25
<b>CITRIC ACID</b> .....	(Citric Acid....87-100 gr.)	
Antiseptic.		

Dissolve a tablet in four ounces of water; the strength of corrosive sublimate in the resulting solution is 1 to 1,000. The citric acid retains in solution the mercury albuminate which is formed, and thus prolongs the antiseptic action.

<b>CORROSIVE SUBLIMATE &amp;</b>	(Corrosive Sub....1 3-4 gr.)	1 25
<b>CITRIC ACID</b> colored blue)	(Citric Acid....87-100 gr.)	

Dissolve a tablet in four ounces of water; the strength of corrosive sublimate in the resulting solution is 1 to 1,000. The citric acid retains in solution the mercury albuminate which is formed, and thus prolongs the antiseptic action.

<b>CORYZA</b> .....	(Quinine Sulph....1-2 gr.)	2 00
(Dr. Richards.)	(Ammon. Mur....1-2 gr.)	
	(Camphor.....1-2 gr.)	
	(Opium Powd....1-10 gr.)	
	(Ext. Belladonna.1-10 gr.)	
	(Ext. Aconite....1-10 gr.)	
<b>Chocolate Coated only</b>		

For a fresh cold, give two tablets for first dose; followed by one every hour until cold is broken up.

<b>CORYZA No. 2</b> .....	(Quinine Sulph....1-3 gr.)	1 50
	(Ammon. Mur....1-3 gr.)	
	(Camphor.....1-3 gr.)	
	(Opium Powd....1-15 gr.)	
	(Ext. Belladonna.1-15 gr.)	
	(Ext. Aconite....1-15 gr.)	
<b>Chocolate Coated only</b>		

For a fresh cold, give two tablets for first dose; followed by one every hour until cold is broken up.

<b>CORYZA No. 3</b> .....	(Camphor.....1-2 gr.)	2 00
(Dr. Henry J. Kenyon.)	(Quinine Sulph....1-2 gr.)	
	(Morph. Sulph....1-64 gr.)	
	(Atropine Sulph.1-2000 gr.)	
	(Sugar of Milk.....q. s.)	
<b>Plain or Chocolate Coated</b>		

Four tablets at bedtime, or one every two to four hours—but preferably at bedtime.

Through the vaso-motor nerves, this tablet acts as a specific stimulant to the congested mucous membrane of the naso-pharynx, promotes secretion and favors the return of the diseased parts to their normal condition in from twenty-four to forty-eight hours.

<b>CORYZA No. 4</b> .....	(Atropine Sulph..1-600 gr.)	1 15
(Dr. S. MacCuen Smith.)	(Acid Arsenous...1-240 gr.)	
	(Strych. Sulph....1-240 gr.)	
	(Morph. Sulph....1-60 gr.)	
	(Quinine Sulph....1-10 gr.)	

One or two every hour for four doses, then one every three or four hours if necessary, to control the discharge and headache of acute coryza.

<b>CORYZA No. 5</b> ..... (Dr. S. MacCuen Smith.)	Atropine Sulph..1-600 gr.	1 15
	Acid Arsenous...1-240 gr.	
	Strych. Sulph....1-240 gr.	
	Morph. Sulph....1-100 gr.	
	Quinine Sulph....1-10 gr.	
	Camphor.....1-4 gr.	

**Chocolate Coated**..... 1 25

One every hour for cold in the head with weeping nostrils, until the throat feels dry; then one every two or three hours, continuing a day or two after the cold has disappeared.

<b>COUGH, ADULTS</b> ..... (Dr. Riggs.)	Ammon. Mur.....3-4 gr.	2 00
	Ext. Opium.....1-15 gr.	
	Fld. Ext. Ipecac..1-4 min.	
	Ext. Bellad.....1-30 gr.	
	Ext. Licorice.....1 gr.	
	Oil Anise.....1-20 min.	

One every hour or two for coughs and bronchial irritation.

<b>COUGH, CHERRY</b> ..... (Drs. S. & W. Hunt.)	Ammon. Mur.....1 gr.	1 75
	Wine Antimony..4 min.	
	Wine Ipecac.....4 min.	
	Syr. Wild Cherry.15 min.	

One every hour until cough becomes loose, then every two hours. For all forms of bronchitis, particularly those accompanied by an irritable cough.

<b>COUGH, INFANTS</b> ..... (Dr. Riggs.)	Ammon. Mur.....1-4 gr.	1 25
	Tr. Opium.....1-4 min.	
	Fld. Ext. Ipecac..1-5 min.	
	Ext. Licorice.....1 gr.	
	Tr. Belladonna..1-5 min.	
	Oil Anise.....1-40 min.	

Dissolve fifteen tablets in fifteen teaspoonfuls of water and give an infant one teaspoonful every hour or two, for bronchial cough and irritation.

<b>COUGH MIXTURE</b> ..... (Dr. Satterlee.)	Ammon. Mur.....1-4 gr.	1 00
	Paregoric.....5 min.	
	Corrosive Sub....1-96 gr.	

Give one every hour, or two every two hours during the early stages of cough, and increase the intervals as the symptoms improve.

<b>COUGH No. 2</b> .....	Ammon. Carb.....1 gr.	1 30
	Squill Powd.....3-4 gr.	
	Senega Powd....3-4 gr.	
	Paregoric.....5 min.	

One every hour or two, for persistent cough.

<b>COUGH No. 3</b> ..... (Dr. E. M. Lyon.)	Tr. Sanguinaria...3 min.	2 00
	Morph. Acetate...1-16 gr.	
	Wine Antimony...5 min.	
	Syr. Wild Cherry....q. s.	

One every two or three hours, as a sedative, diaphoretic and expectorant.

<b>COUGH No. 4</b> ..... (Dr. H. B. Delatour.)	Tr. Aconite.....1-2 min.	1 15
	Wine Antimony...2 min.	
	Paregoric.....5 min.	

One or two tablets every hour during the acute stages of laryngeal inflammation; especially useful in croup of children.

<b>COUGH, PERSISTENT</b> ..... (Dr. Brewer.)	Ammon. Mur.....2 gr.	1 60
	Ipecac Powd....1-4 gr.	
	Ext. Squill Acet...1-4 gr.	
	Fl. Ext. Senega...1 min.	

One every two or three hours, for cough.

**CREOSOTE (Beechwood.)**.....1-10 gr..... 75

A tablet three times a day, gradually increasing the dose, for diarrhœa, flatulence, and tuberculosis.

**CREOSOTE (Beechwood.)**.....1-2 min..... 1 15  
**Chocolate Coated**..... 1 25

One or two after each meal, for consumption, tubercular diarrhœa, and as an intestinal disinfectant.

**CREOSOTE (Beechwood.)**.....1 min..... 1 40  
**Chocolate Coated**..... 1 50

One to three after meals, for consumption, tubercular diarrhœa, and as an intestinal disinfectant; useful in the abdominal torpidity of epileptics.

**CREOSOTE (Beechwood)**.....2 min..... 1 75  
**Chocolate Coated**..... 1 85

One to two after meals, for consumption, tubercular diarrhœa, and as an intestinal disinfectant; useful in the abdominal torpidity of epileptics.

**CREOSOTE COMP**..... { Creosote.....1-8 min.  
 Cocaine Hydroch.....1-20 gr.  
 Cerium Oxalate.....2 gr.  
 Pepsin Pure.....1-4 gr.  
 Tinct. Nux Vom.....1-4 min.  
**Plain or Chocolate Coated**..... 2 00

One or two after meals, repeated if necessary in two hours, for nausea and vomiting. One or two every two or three hours for seasickness; to relieve the nausea of travelling, and the nausea and vomiting of pregnancy.

**CROUP, (Spasmodic.)**..... { Ext. Can. Indica.....1-20 gr.  
 Ext. Hyoscyam.....1-20 gr.  
 Paregoric.....5 min.  
 Syr. Ipecac.....5 min.  
 Ext. Licorice.....2 gr.  
 1 75

Dissolve six tablets in eighteen teaspoonfuls of hot water, stir and give a teaspoonful every fifteen minutes to a child six months old, until relieved.

**CROUP No. 2**..... { Dover's Powder...1-2 gr.  
 Ipecac Powd....1-100 gr.  
 Spongia.....1-100 gr.  
 1 00

One hourly until symptoms are relieved. Give every half-hour for three doses, if symptoms are urgent; then one hourly.

**CROUP No. 3**..... { Kermes Mineral...1-6 gr.  
 Ipecac Powd....1-12 gr.  
 1 00

One every one-half to one hour as an expectorant; for croup use three to four tablets as an emetic.

**CUBEBS COMP**..... { Rhubarb Powd...1-2 gr.  
 Cubebs Powd....1-2 gr.  
 Calc. Magnesia.....1 gr.  
 Oil Peppermint,1-20 min.  
 1 10

A tablet half an hour before meals with a little water, to stimulate the peptic glands and induce appetite.

**CYSTITIS**..... { Acid Boracic.....2 gr.  
 Potass. Bicarb.....2 gr.  
 Ext. Buchu.....1 gr.  
 Ext. Triticum Rep..1 gr.  
 Ext. Corn Silk....1-2 gr.  
 Ext. Hydrangea..1-2 gr.  
 Atropine Sulph..1-500 gr.  
 3 00

One every three hours, with a large glass of water; for inflammation of the bladder and urinary tract, as well as suppressed renal secretion.

When ordering or prescribing

CYSTITIS No. 2, Improved.....	Acid Benzoic.....1 gr.	1 25
	Sodium Borate.....1 gr.	
	Oil Gaultheria.....q. s.	

Two every two hours, generally used for about ten days, for cystitis, gravel and some types of rheumatism.

CYSTITIS No. 3 .....	Acid Benzoic.....3 gr.	3 40
	Sodium Biborate....2 gr.	
	Ext. Buchu.....1 gr.	
	Ext. Tritic. Repens..1 gr.	
	Ext. Corn Silk.....1-2 gr.	
	Ext. Hydrangea...1-2 gr.	
	Atropine Sulph..1-500 gr.	

One to two every three hours, with large glass of water, for alkaline urine.

DAMIANA COMP.....	Ext. Damiana.....2 gr.	2 65
	Phosphorus.....1-30 gr.	
	Ext. Nux Vom.....1-4 gr.	
	Plain or Chocolate Coated	

One morning and night for a week, then one every night, as a stimulant to the generative organs.

DAMIANA CO. NO. 2..... (Dr. La Madrid.)	Zinc Phosphide...1-10 gr.	2 75
	Ext. Cannab. Ind...1-4 gr.	
	Strych. Sulph.....1-40 gr.	
	Ext. Damiana...1-2 gr.	
	Plain or Chocolate Coated	

One after meals, as a nerve tonic.

DAMIANA COMP. No. 3.....	Ext. Damiana.....2 gr.	2 65
	Ext. Nux Vom.....1-5 gr.	
	Cantharides...1-20 gr.	
	Plain or Chocolate Coated	

One tablet three times a day, for impotence.

DANDELION COMP.....	Fl. Ex. Dandelion..5 min.	2 00
	Fl. Ext. Cascara...5 min.	
	Plain or Chocolate Coated	

Two at night, or night and morning if necessary; as a laxative in chronic constipation.

DIABETIC..... (Dr. Harley.)	Croton Chloral....1-2 gr.	3 00
	Ext. Opium.....1-2 gr.	
	Ext. Aloes Barb...1-6 gr.	
	Plain or Chocolate Coated	

One tablet three times a day.

DIAPHORETIC ANODYNE.... (Dr. S. Lovigett.)	Morph. Acetate...1-20 gr.	1 65
	Acetanilid.....1-2 gr.	
	Tr. Aconite Rad..1-2 min.	
	Ipecac Powd.....1-8 gr.	

One tablet every hour or two hours as needed.

DIARRHŒA.....	Calomel.....1-8 gr.	1 40
	Morph. Sulph....1-16 gr.	
	Capsicum Powd...1-16 gr.	
	Ipecac Powd.....1-32 gr.	
	Camphor.....1-16 gr.	
	Chocolate Coated.....	1 50

One tablet every hour for four doses, then every two hours until relieved.

DIARRHŒA, INFANTS.. . . (Dr. Liggett.)	Sodium Salicylate...1 gr.	1 75
	Bismuth Subcarb...1 gr.	
	Beta Naphtol....1-5 gr.	

One every hour or two for fermentative diarrhœa in infants.

**DIARRHŒA, INFANTS No. 2.** { Copper Arsen....1-100 gr. }  
 { Merc. Bichlor....1-100 gr. } 1 00  
 { Morph. Sulph....1-150 gr. }  
 One every one or two hours for infants diarrhœa.

**DIARRHŒA No. 2.** { Pepsin Pure.....1-2 gr. }  
 { Bismuth Subnit....6 gr. } 4 50  
 (Dr. Hinz.) { Fl. Ext. Nux Vom. 2 min. }  
 { Fl. Ext. Hydrast. 5 min. }  
 One every two hours for three doses, then every three hours, for diarrhœa caused by indigestion.

**DIARRHŒA No. 3.** { Opium Powd.....1-4 gr. }  
 { Camphor.....1-4 gr. } 1 15  
 (Dr. Sullivan.) { Ipecac Powd....1-8 gr. }  
 { Lead Acetate....1-6 gr. }  
 Chocolate Coated ..... 1 25

Give two at the first dose, then one every hour for six hours, afterwards one after each loose movement, for congestive, irritative and inflammatory diarrhœas.

**DIARRHŒA No. 4.** { Bismuth Subnit.... 2 gr. }  
 { Salol.....2 gr. } 4 25  
 (Dr. Tinker.)  
 One every two hours in acute cases; one every four hours in chronic cases. For gastro-intestinal catarrh in adults and children.

**DIARRHŒA No. 5.** { Bismuth Subnit.... 3 gr. }  
 { Sacch. Pepsin.....2 gr. } 3 50  
 (Board of Health.) { Arom. Chalk Powd. 2 gr. }  
 One tablet after each movement; for diarrhœa, accompanied by irritation from undigested food.

**DIARRHŒA No. 6.** { Bismuth Subnit....1 gr. }  
 { Ipecac Powd....1-20 gr. } 1 50  
 (Dr. Elliott Riggs.) { Opium Powd....1-40 gr. }  
 For Children.

One every two or three hours to a child about one year old, as may be necessary to check diarrhœa; preceded by eight to ten doses Bismuth and Ipecac Comp. No. 2, taken one every three hours.

**DIARRHŒA No. 8.** { Opium Powd.....1-8 gr. }  
 { Ipecac Powd....1-15 gr. } 1 00  
 (Dr. Drumheller.) { Lead Acetate....1-4 gr. }  
 One tablet every two or three hours, for diarrhœa.

**DIARRHŒA No. 9, Improved.** { Comp. Chalk Powd. 1 gr. }  
 { Acid Tannic.....1-20 gr. } 1 00  
 { Opium Powd....1-40 gr. }  
 { Camphor.....1-40 gr. }  
 { Oil Cinnamon.....q. s. }

One crushed and taken on the tongue or diffused in water on the end of a spoon, every twenty minutes, for a baby with diarrhœa, pain or flatulence; after the sixth dose, give every hour.

**DIARRHŒA No. 10.** { Camphor Water...4 min. }  
 { Paregoric .....4 min. } 3 75  
 (Dr. Foster.) { Bismuth Subnit....4 gr. }  
 { Syr. Rhub. Arom. 20 min. }  
 { Oil Peppermint. 1-8 min. }

Two tablets every one, two or three hours, as case demands.

**DIARRHŒA No. 11.** { Salol.....1-2 gr. }  
 { Bismuth Salicylate. 1 gr. }  
 { Calomel.....1-40 gr. } 2 75  
 (Dr. Prendergast.) { Paregoric.....2 min. }  
 { Mist. Cret. Arom....q. s. }  
 { Oil Cinnamon.....q. s. }

One or two tablets every hour until patient is better; for diarrhœa, accompanied by colicky pains, and where an intestinal antiseptic is indicated.

When ordering or prescribing

DIARRHŒA No. 12. (Dr. J. D. Kennedy.)	Copper Sulph. ....	1-12 gr.	2 00
	Zinc Sulphate. ....	1-12 gr.	
	Ext. Nux Vom. ....	1-6 gr.	
	Ipecac Powd. ....	1-10 gr.	
	Opium Powd. ....	1-2 gr.	

One or two as occasion requires; in diarrhœa with tenesmus, pain and watery stools. For children with green, pasty stools and acidity of stomach, give one-half to one tablet according to age.

DIARRHŒA No. 13 (Dr. E. Chapin.)	Tr. Opium Deod. ....	4 min.	7 00
	Tr. Ginger. ....	4 min.	
	Tr. Catechu. ....	8 min.	
	Syr. Rhub. Ar. ....	3-4 teasp.	

A tablet every two hours, or one after each loose movement, if necessary.

DIARRHŒA No. 14. (Drs. Lewis & Coffman.)	Opium Powd. ....	1-4 gr.	1 35
	Camphor Powd. ....	1-4 gr.	
	Ipecac Powd. ....	1-4 gr.	
	Lead Acetate ....	1-4 gr.	
Chocolate Coated			1 45

One every hour for six doses, then one after each loose movement until symptoms improve.

DIARRHŒA, FERMENTA- TIVE. (Dr. J. W. Hyde.)	Bismuth Subgall. ....	1 gr.	2 00
	Salol. ....	1 gr.	
	Cerium Oxalate ....	1 gr.	
	Dover's Powd. ....	2 gr.	
	Oil Anise. ....	1-10 min.	

For adults, one tablet every two or three hours; for children, mix one with four to six teaspoonfuls of water and give one teaspoonful every two or three hours.

DIFFUSIVE MALARIAL	Corrosive Sub. ....	1-50 gr.	1 25
	Ammon. Mur. ....	2 gr.	
	Cinchonia Sulph. ....	1-10 gr.	
	Cinchonid. Sulph. ....	1-10 gr.	
Quinine Sulph. ....			1-10 gr.

One to three tablets after meals, preceded by three one grain calomel tablets at intervals of an hour; for dumb ague, headache and cachexia of malarial origin.

DIGESTIVE	Pepsin. ....	2 gr.	3 50
	Pancreatin. ....	1 gr.	
	Ipecac Powd. ....	1-10 gr.	
	Jamaica Ginger. ....	4-5 gr.	
	Bismuth Subnit. ....	2 gr.	
	Ext. Nux Vom. ....	1-10 gr.	
Oil Wintergreen. ....			q. s.

One or two after meals, repeated in an hour, for cases of impaired or slow digestion requiring a stomachic and carminative in addition to the ordinary solvents for food.

DIGESTIVE No. 2. (Dr. Howard.)	Pepsin. ....	1-3 gr.	2 25
	Bismuth Subnit. ....	2-3 gr.	
	Sodium Bicarb. ....	1-2 gr.	
	Pancreatin. ....	2-3 gr.	
	Ginger Powd. ....	1-3 gr.	
Charcoal. ....			1 1-3 gr.

One or two after eating, followed by another in an hour if necessary.

DIGESTIVE TONIC (Dr. Satterlee.)	Iron Lactate. ....	1 gr.	11 00
	Quinine Sulph. ....	1 gr.	
	Acid Arsenous. ....	1-30 gr.	
	Strych. Sulph. ....	1-30 gr.	
	Lactopeptine. ....	2 gr.	
Per 100. ....			1 25

One tablet before each meal, as a general tonic and appetizer, as well as digestive. Is well borne by the most delicate stomach.

DIGITALIA.....	1-1000 gr.....	1 00
DIGITALIA.....	1-500 gr.....	1 10
DIGITALIA.....	1-400 gr.....	1 25
DIGITALIA.....	1-300 gr.....	1 35
DIGITALIA.....	1-200 gr.....	1 50
DIGITALIA.....	1-150 gr.....	1 75
DIGITALIA.....	1-100 gr.....	2 25
DIGITALIA.....	1-60 gr.....	3 00
DIGITALIA.....	1-50 gr.....	3 50
DIGITALIS COMP.....	{ Tr. Digitalis..... 2 min. }	1 00
(Dr. Wood.)	{ Brucia..... 1-100 gr. }	
Give a tablet every three hours, as a heart stimulant.		
DIGITALIS COMP. No. 2.....	{ Tr. Digitalis... 2 min. }	1 00
(Dr. Wood.)	{ Strych. Sulph... 1-100 gr. }	
One every three hours, as a heart stimulant and tonic.		
DIGITALIS FL. EXT.....	1-2 min.....	75
One every three or four hours, for mitral disease, partial suppression of urine, and dropsy.		
DIGITALIS FL. EXT.....	1 min.....	1 00
Give one after meals to control excessive and irregular heart action and stimulate the kidneys; reduce the dose if headache and flushing occurs.		
DIGITALIS POWD.....	1-20 gr.....	75
Heart tonic and diuretic.		
One every three hours for a child, to regulate the heart and increase the flow of urine.		
DIGITALIS POWD.....	1-10 gr.....	75
A tablet every three hours, for a child as a diuretic and to strengthen cardiac systole.		
DIGITALIS POWD.....	1-5 gr.....	75
Give one every three hours to assist absorption of serous exudations and as a heart tonic.		
DIGITALIS POWD.....	1-4 gr.....	75
One every three or four hours, as a heart tonic.		
DIGITALIS POWD.....	1-2 gr.....	75
One tablet three times a day, as a renal and cardiac tonic.		
DIGITALIS & IRON COMP.....	{ Ext. Digitalis... 1-10 gr. }	1 00
Cardiac tonic and hæmatic.	{ Iron Phosphate... 1-4 gr. }	
	{ Strych. Sulph... 1-60 gr. }	1 10
Chocolate Coated.....		
A tablet after meals, for anæmia with palpitation of the heart, and in convalescence from exhausting illness.		
DINNER.....	{ Aloe Purif..... 2 gr. }	1 50
(Lady Webster.)	{ Mastic..... 1-2 gr. }	
	{ Red Rose..... 1-2 gr. }	1 60
Chocolate Coated.....		
One to three at night, or one or two after the principal meal of the day; for the relief of indigestion with constipation.		

When ordering or prescribing

<b>DIPHTHERIA &amp; GRIP</b> ..... (Dr. Briggs.)	(Hydrarg. Bichlor 1-50 gr.)	2 00
	Guaiac.....1 gr.	
	Tr. Sanguinaria...3 min.	
	Tr. Colchicum.....6 min.	
	Tolu.....1-5 gr.	
	Rhubarb Powd...1-2 gr.	
	Cubeb's Powd...1 1-5 gr.)	

Dissolve a tablet slowly in the mouth every two or three hours, supplement with a heart tonic and quinine, also with a salicylate of sodium tablet, if needed, for relief of pain.

<b>DIPSOMANIA</b> .....	{ Gold & Sod. Chlor 1-20 gr. }	2 00
	{ Strych. Nitrate... 1-60 gr. }	

One tablet every two or three hours, reducing to three a day, to control the craving for alcohol and as a tonic after a debauch.

<b>DIPSOMANIA No. 2</b> .....	(Gold & Sod. Chl...1-24 gr.)	2 75
	Strych. Nitrate...1-60 gr.	
	Nitroglycerine .1-400 gr.	
	Atropine Sulph. 1-200 gr.	
	Tr. Digitalis.....3 min.	
	Oleoresin Capsic...1-8 gr.)	
	<b>Plain or Chocolate Coated</b>	

One or two every two hours cautiously increased; used in nervous debility following overwork, excesses or debauches, and especially in controlling opium habit and chronic alcoholism.

<b>DIURETIC</b> .....	(Digitalis Powd....1 gr.)	1 75
	Ext. Buchu.....1 gr.	
	Potass. Nitrate ....1 gr.	
	Squill Powd.....1 gr.)	
	<b>Chocolate Coated</b> .....	1 85

One, three times a day, followed by a large glass of water.

<b>DIURETIC No. 2</b> ..... (Dr. Waldstein.)	{ Caffeine.....1-4 gr. }	1 05
	{ Potass. Nitrate...1-2 gr. }	
	{ Lithium Carb ....1-4 gr. }	

Three to six daily as a diuretic.

<b>DIURETIC No. 4</b> ..... (Dr. Christy.)	(Digitalis Powd.....1 gr.)	2 50
	Potass. Nitrate....2 gr.	
	Ext. Buchu .....1-2 gr.	
	Ext. Scopolarius...1-2 gr.	
	Oil Juniper.....1 gtt.)	

One every two hours as a diuretic; after the third dose give every four hours, for suppression of urine, dropsy and chronic Bright's disease; omit the medicine one or two days each week.

<b>DIURETIC No. 5</b> .....	(Digitalis Powd....1 2 gr.)	1 50
	Squills Powd.....1-2 gr.	
	Calomel.....1-2 gr.	
	Caffeine Alk.....1-4 gr.)	

One three times a day, followed each time by a large glass of water.

<b>DOBELL'S SOLUTION</b> .....	2 00
--------------------------------	------

Each tablet represents one ounce.

Dissolve one tablet in one to four fluid ounces of water, add half teaspoonful of glycerine, to be used as a nasal douche or in atomizer as spray, or for gargle.

<b>DOVER'S COMP</b> ..... (Dr. Terriberry.)	{ Quinine Sulph....1-2 gr. }	1 75
	{ Dover's Powder..1-2 gr. }	

To an infant under six months, one every two hours at first, in syrup; over six months and up to two years, give one every hour at first, then diminish the dose as the effect is felt. Used in acute bronchitis and pneumonia of children.



<b>DYSPEPTIC No. 2</b> .....	{ Bismuth Subgall.....5 gr. }	4 25
Laxative.	{ Ext. Cascara.....1 gr. }	

One three times a day, for flatulent or fermentative dyspepsia.

<b>DYSPEPTIC No. 3</b> .....	{ Codeine Sulph.....1-6 gr. }	6 35
(Dr. C. F. Halstead.)	{ Ext. Nux Vom.....1-4 gr. }	
	{ Ext. Pancreatis.....3 gr. }	
	{ Ox Gall Purif.....1 gr. }	
	{ Ext. Belladonna...1-4 gr. }	

One tablet immediately after meals.

<b>DYSPEPTIC COMP</b> .....	{ Pepsin.....1 gr. }	1 85
(Dr. Outerbridge.)	{ Charcoal.....3 gr. }	
	{ Sodium Bicarb. 2 1-2 gr. }	

One or two after eating, followed by another in one hour, for dyspepsia and acid stomach.

<b>DYSPEPTIC FERMENTATIVE</b> .....	{ Bismuth Subnit.....2 gr. }	2 25
(Dr. Drumheller.)	{ Paregoric.....10 min. }	
	{ Ammon. Carb.....1 gr. }	

One tablet every two or three hours, for fermentative or functional dyspepsia.

<b>EISENZUCKER</b> .....	3 grs.	1 00
Children's iron tonic.		

One to three tablets, three times a day after meals, as an iron tonic for children.

<b>EISENZUCKER</b> ....	5 grs.	1 25
-------------------------	--------	------

One to three tablets three times a day after meals, as an iron tonic.

<b>ELATERIN CRYSTAL</b> .....	1-10 gr.	7 50
Purgative.		

One every three hours for three doses if necessary, where a strong purgative effect is required; best combined with inspissated ox gall, about three grains to each dose of elaterin.

<b>ELATERIUM, Clutterbuck's</b> .....	1-10 gr.	1 25
---------------------------------------	----------	------

One at night as a purge; one, repeated in an hour, followed by a saline, for suppression of urine and where sure action is necessary.

<b>EMMENAGOGUE</b> .....	{ Ergotin.....1 gr. }	4 00
	{ Ext. Gossypium....1 gr. }	
	{ Iron Sulph Dry....1 gr. }	
	{ Ext. Bl'k Hellebore.1 gr. }	
	{ Aloes Socotrine....1 gr. }	
	{ Oil Savine.....1-4 min. }	

Plain or Chocolate Coated

One every two hours as an emmenagogue.

<b>EMMENAGOGUE No. 2</b> .....	{ Pulv. Aloes Soc....1 gr. }	1 65
	{ Ferri Sulph.Exsic.1-2 gr. }	
	{ Ext. Helleb. Nig..1-4 gr. }	
	{ Pulv. Myrrh.....1-4 gr. }	
	{ Pulv. Sapo.....1-4 gr. }	
	{ Pulv. Canella.....1-8 gr. }	
	{ Pulv. Zingib.Jam..1-8 gr. }	

Chocolate Coated.....

One to three tablets three times daily.

<b>EMMENAGOGUE, 1-2 strength</b> .....	{ Ergotin.....1-2 gr. }	3 00
	{ Ext. Gossypium....1-2 gr. }	
	{ Iron Sulph. Dry...1-2 gr. }	
	{ Ext. Bl'k Helleb..1-2 gr. }	
	{ Aloes Socotrine...1-2 gr. }	
	{ Oil Savine.....1-8 min. }	

Chocolate Coated only

One or two tablets every two or three hours, as an emmenagogue.

**EMETIA.** ..... { Calomel.....1-10 gr. } 1 50  
 { Pimenta Powd.....5 gr. }  
 One repeated every ten minutes for four doses, or until vomiting is checked.

**ENTERIC TONIC.** ..... { Strych. Sulph.....1-60 gr. }  
 { Leptandrin.....1-5 gr. }  
 { Cascarin.....1-5 gr. } 2 25  
 (Dr. North.) { Hydrastin.....1-5 gr. }  
 { Ext. Hyoscyam.....1-5 gr. }  
 { Oil Anise.....1-10 min. }

For habitual constipation, take one tablet at night and go to stool at a definite time of day. In case of failure, take morning and night.

**EPILEPTIC**..... { Sodium Bromide....3 gr. }  
 { Ammon. Bromide...3 gr. }  
 { Potass. Bromide....3 gr. } 3 00  
 (Dr. Batton.) { Tr. Calab. Bean 2 1-2 min. }  
 { Ext. Hyoscyam....1-2 gr. }

Two tablets three times daily with a swallow of water; may be given for a year if required, for epilepsy.

**EPILEPTIC No. 2**..... { Sodium Bromide....3 gr. }  
 { Ammon. Bromide...3 gr. } 2 75  
 (Dr. Batton.) { Potass. Bromide....3 gr. }  
 { Tr. Calab. Bean 2 1-2 min. }

Two tablets three times daily with a swallow of water; may be used continuously for a long period.

**ERGOTIN**.....1-50 gr..... 75

One every two hours to a child, as a general nerve tonic and vaso-motor stimulant.

**ERGOTIN**.....1-10 gr..... 1 00

One three times a day for a general vaso-motor tonic in chorea and epilepsy of children.

**ERGOTIN**.....1-6 gr..... 1 15

One three times a day as a general nerve tonic and to counteract tendency to internal hemorrhage.

**ERGOTIN**.....1-4 gr..... 1 25

**Chocolate Coated** 1 35

One to four, repeated as required for nerve tonic.

**ERGOTIN**.....1-2 gr..... 1 60

**Chocolate Coated** 1 70

One every hour or two as an oxytocic and to control internal bleeding.

**ERGOTIN**.....1 gr..... 2 00

**Plain or Chocolate Coated**

A tablet every hour, as necessary, to control hæmoptysis, and hemorrhage from the bowel; also after parturition, to secure uterine contraction.

**ERGOTIN**.....2 gr..... 3 25

**Plain or Chocolate Coated**

One every three hours to control internal hemorrhage, and as a general nerve tonic.

**ERGOTIN COMP.**..... { Ergotin.....1 1-2 gr. }  
 { Tr. Digitalis.....1-2 min. } 3 25  
 General tonic. { Quinine Sulph.....1-2 gr. }

One to three tablets every three hours for nerve and heart tonic.

**ERGOTIN COMP. No. 2**..... { Ergotin.....1-2 gr. } 2 00  
 { Ext. Can. Indica...1-4 gr. }

One every three hours.

<b>ESERINE SULPHATE</b> .....	I-100 gr.....	1 75
Cerebral tonic.		
A tablet three times a day as a cerebral tonic in tetanoid and choreic conditions.		
<b>ESERINE SULPHATE</b> .....	I-50 gr.....	2 50
A tablet one to three times a day in convulsive affections and intestinal atony.		
<b>EUONYMIN</b> .....	I-10 gr.....	1 00
Hepatic.		
One every hour for five doses, in hepatic torpor.		
<b>EUONYMIN</b> .....	I-8 gr.....	1 15
A tablet every hour for three doses, followed by an aperient, for torpidity of the liver.		
<b>EUONYMIN</b> .....	I-4 gr.....	1 50
One three times a day, with half an ounce of castor oil at night, for torpidity of the liver and duodenal catarrh.		
<b>EUONYMIN</b> .....	I-2 gr.....	2 25
One to three a day, in divided doses, followed by an aperient, for deficiency of bile in the intestines.		
<b>EUONYMIN COMP.</b> .....	{ Euonymin.....I-4 gr. Podophyllin.....I-8 gr. Aloin.....I-8 gr. }	1 75
(Dr. Welch.)		
	Chocolate Coated.....	1 85
One every night for a week, and one after meals every second day, if necessary, in cases of hepatic torpor with biliousness, headache and light colored feces.		
<b>EXPECTORANT</b> .....	{ Bellad. Fl. Ext.. I-8 min. Ipecac Powd....I-10 gr. Ext. Licorice....I-8 gr. Codeine.....I-16 gr. Ext. Senega....I-10 gr. }	2 25
(Dr. Rankin.)		
One every hour or two, for an irritative bronchial cough.		
<b>EXPECTORANT, Anodyne No. 2</b>	{ Ammon. Mur.....I gr. Tartar Emetic....I-30 gr. Sanguinar. Nit..I-100 gr. Morphine Sulph..I-50 gr. }	1 50
(Dr. Bolton.)		
One or two tablets every two hours, as an anodyne in coughs with respiratory inflammation.		
<b>EXPECTORANT No. 4</b> .....	{ Ammon. Carb.....I gr. Ammon. Muriate...I gr. Ext. Licorice.....I-2 gr. Squill Powd. ....I-4 gr. Oil Cinnamon.....q. s. }	1 25
(Dr. Bierhoff.)		
One every hour or two, as a stimulant and expectorant in laryngitis and bronchitis of children; best given dissolved in one or two teaspoonfuls of water.		
<b>EXPECTORANT No. 5</b> .....	{ Ammon. Mur.....2 gr. Morph. Sulph....I-25 gr. Tartar Emetic....I-25 gr. Senega Fl. Ext..I-2 min. Oil Wintergr'n.I-10 min. }	1 50
A tablet every hour or two, as an expectorant in cough.		
<b>EXPECTORANT TONIC</b> .....	{ Strych. Sulph....I-60 gr. Tr. Digitalis.....2 min. Terpin Hydrate...3 gr. Opium Powd.....I-24 gr. Ext. Hyoscyam...I-4 gr. }	2 00
One or two tablets three or four times daily, for winter cough of aged or infirm.		

<b>FEVER No. 2</b> .....	{	Tr. Aconite.....1-5 min.	1 00
		Tr. Bryonia.....1-10 min.	
		Tr. Belladonna..1-10 min.	

(Dr. Davis.)

One tablet every two or three hours, with plenty of water, for the fevers and ordinary colds of infancy and childhood. Give adults two to four tablets at a dose.

<b>FEVER No. 3</b> .....	{	Tr. Aconite.....1-2 min.	1 40
		Tr. Bryonia.....1-10 min.	
		Codeine Sulph....1-40 gr.	
		Tartar Emetic....1-80 gr.	

(Dr. J. C. Applegate.)

One every two hours for a child and every hour for an adult, for fever, or colds with fever.

<b>FEVER No. 4</b> .....	{	Tr. Aconite.....1 min.	1 15
		Tr. Gelsemium....1 min.	
		Tr. Belladonna....1 min.	

(Dr. F. W. Koehler.)

One every hour to adults; for children, dissolve two tablets in ten teaspoonfuls of water and give a teaspoonful every hour. Used in fevers as an antipyretic and anodyne, also used in many forms of neuralgia.

<b>FEVER, IMPROVED</b> .....	{	Tr. Aconite.....1 min.	1 50
		Morph. Sulph....1-50 gr.	
		Tartar Emetic....1-50 gr.	
		Ipecac Powd....1-4 gr.	

(Dr. Kenyon.)

One every hour for six doses, then every two to four hours; for cold with fever, to relieve the sensation of tightness or fulness in the bronchial tubes, and in the early stage of bronchitis.

<b>FEVER, INFANTS</b> .....	{	Aconite.....1-100 gr.	75
		Ipecac Powd....1-100 gr.	

One every hour for ten doses, afterwards at intervals of two to four hours.

<b>FEVER &amp; AGUE</b> .....	{	Copper Sulph....1-10 gr.	2 00
		Opium Powd....1-8 gr.	
		Quinine Sulph....1-2 gr.	
		Podophyllin....1-12 gr.	

One to six a day for fevers due to malarial infections; occasionally as a preventative in malarious districts.

<b>FOLLICULAR TONSILLITIS</b> .....	{	Tr. Aconite....1-5 min.	1 50
		Tr. Belladonna..1-10 min.	
		Tr. Bryonia....1-10 min.	
		Merc. Iod. Red..1-100 gr.	
		Morphine Sulph.1-100 gr.	
		Sodium Salicylate..1 gr.	
		Oil Wintergr...1-20 min.	

One every fifteen minutes dissolved on back of tongue for four doses; then one every hour.

<b>FOUR CHLORIDES</b> .....	{	Iron Chloride....1-2 gr.	3 00
		Arsenic Chlor....1-64 gr.	
		Merc. Bichlor...1-48 gr.	
		Quinine Chloride...1 gr.	

Tonic.

One just after each meal for tonic in anæmia, and as an alterative.

<b>FOUR CHLORIDES No. 2</b> .....	{	Iron Chloride....1-2 gr.	2 00
		Arsenic Chlor....1-100 gr.	
		Merc. Bichlor...1-100 gr.	
		Quinine Chloride.1-2 gr.	

One after each meal. A useful tonic in all kinds of anæmia.

<b>GALLIC ACID COMP.</b> .....	{	Gallie Acid.....3 gr.	1 85
		Ext. Belladonna..1-5 gr.	

General astringent.

One tablet as necessary for hæmatemesis, menorrhagia and as a general astringent.

<b>GARGLE</b> .....	<div> <div> Sodium Bicarb. .3 1-3 gr.  Sodium Salicyl. .3 1-3 gr.  Acid Boracic. ....3 1-3 gr.  Oil Wintergreen. ....q. s. </div> </div>	2 00
---------------------	--	------

Three tablets to a cupful of hot water; gargle every hour until relieved, using it hot if possible.

<b>GASTRITIS</b> .....	<div> <div> Ext. Nux Vomica. 1-4 gr.  Ext. Belladonna 1-8 gr.  Ext. Opium .....1-8 gr.  Potass. Nitrate...1-2 gr.  Silver Nitrate....1-4 gr. </div> </div>	2 50
<b>Plain or Chocolate Coated</b>		

One an hour before eating in gastric trouble; one tablet one or two hours after eating in enteritis duodinal and all intestinal troubles.

<b>GENTIAN, ALKALINE</b> .....	<div> <div> Infus. Gent. Co...1 teasp. }  Sodium Bicarb. ....5 gr. } </div> </div>	1 25
(Dr. J. J. Moran.)		

Two or three tablets dissolved in water before each meal. A tonic stomachic where there is catarrhal gastric disorder, and ordinary atonic dyspepsia.

<b>GELSEMIN</b> .....	1-16 gr.	1 25
<b>Antispasmodic and relaxant.</b>		

One every hour for three doses, for neuralgia, toothache and nervous cough.

<b>GELSEMIN</b> .....	1-10 gr.	1 50
-----------------------	----------	------

A tablet, repeated in an hour, together with hot baths, for dental neuralgia; and for hepatic and renal colic.

<b>GELSEMIN</b> .....	1-8 gr.	1 65
-----------------------	---------	------

A tablet three times a day as a depressant in pleurisy and pneumonia, and for cough with a strong nervous element.

<b>GELSEMIN</b> .....	1-4 gr.	2 35
-----------------------	---------	------

One or two doses at intervals of two hours, for dental neuralgia, headache, and to control the paroxysms of asthma.

<b>GELSEMINE SULPH</b> .....	1-100 gr.	4 50
------------------------------	-----------	------

Three to five a day in divided doses, for spermatic and neuralgic affections and as a sedative to the spinal cord.

<b>GELSEMINE SULPH</b> .....	1-50 gr.	8 00
------------------------------	----------	------

A tablet every three or four hours, for neuralgia; also as an antidote to Strychnine.

<b>GELSEMIUM FL. EXT.</b> .....	1 min.	1 00
---------------------------------	--------	------

One every hour to six doses, in the onset of fevers, especially of malarial origin, and for various neuralgias.

<b>GOLD &amp; SODIUM CHLORIDE</b> .....	1-30 gr.	2 00
---	----------	------

**Tonic.**

A tablet after meals as a general tonic, for anæmia and after wasting disease.

<b>GOLD &amp; SODIUM CHLORIDE</b> .....	1-20 gr.	2 50
---	----------	------

A tablet after meals as a general tonic, for anæmia and after wasting disease.

<b>GOLD &amp; SODIUM CHLORIDE</b> .....	1-10 gr.	4 25
---	----------	------

One with meals, and one at night, to combat alcohol and morphine craving, to hasten the elimination of syphilitic poison, and as a hæmal tonic.

GONORRHOEA.....	{	Cubebs Powd.....	1 gr.
		Copaiba Solid.....	1 gr.
		Iron Sulph. Dry...	1-4 gr.
		Oil Sandal.....	1-4 min.
		Oil Wintergr.....	1-10 min.
		Venice Turpen..	1-4 min.

Plain or Chocolate Coated 2 25

Two tablets, three or four times daily, usually supplemented with ten grains of bicarbonate of sodium or potassium.

GONORRHŒA No. 2..... (Dr. Burd.)	{	Cubebs Powd....	1-4 gr.	I 50
		Copaiba Solid...	1-4 gr.	
		Iron Sulph. Dry...	1-4 gr.	
		Venice Turpent....	1-2 gr.	
		Chocolate Coated.....		

1 50

1 60

One tablet every two hours, or two tablets every three hours; used in catarrhal or subacute stage of gonorrhœa.

GONORRHŒA No. 4..... (Dr. C. E. Campbell.)	{	Tr. Aconite.....	1 min.	I 25
		Tr. Gelsemium....	1 min.	
		Tr. Cubebs.....	1 min.	
		Tr. Cantharides...1	min.	

1 25

One every three hours with a half teaspoonful of Spts. Nitre, in gonorrhœa after the acute stage.

GRAY POWDER COMP..... (Dr. Anderton.)	(	Merc. with Chalk...1	gr.	} 3 75
		Salol.....1	gr.	
		Bismuth Subnit....3	gr.	

3 75

One every hour until ten doses have been taken; for torpid liver, loose, colorless, mal-odorous stools, with or without vomiting.

GRAY POWDER CO. No. 2... (Dr. Hutchinson.)	{	Merc. with Chalk...1	gr.	1 35
		Dover's Powder....1	gr.	

1 35

Give one tablet every hour or two until symptoms improve; then every three hours, for a child with cough, fever and constipation.

GRIP.....	{	Ammon. Salicyl....	3 gr.	} 6 50
		Phenacetine.....	1 gr.	
		Caffeine.....	1-2 gr.	

6 50

A tablet every two or three hours in the febrile stages of influenza, until the cardiac depression and the fever improve.

GRIP No. 3..... (Dr. Searles.)	{	Phenacetine.....	1 gr.	} 7 00
		Salicin.....	1 gr.	
		Quinine Sulph....	1-2 gr.	

7 00

One, cautiously, preceded by a cathartic, every hour for three doses, then one every two hours, for grip.

GRIP No. 4.....	{	Acetanilid.....	2 gr.	} 3 65
		Caffeine Cit.....	1-2 gr.	
		Salol.....	2 gr.	

3 65

One every two to three hours, for grip with violent pains in head, with fever. After first day give one three times a day with two grains of quinine until pain and fever subside.

GRIP No. 5.....	{	Acetanilid.....	2 gr.	} 5 00
		Quinine Sulph.....	2 gr.	
		Salol.....	2 gr.	

5 00

One every two to three hours in the early stages of grip, until the pains are relieved, then one three times a day until the fever disappears.

GUAIAIC COMP..... Throat.	(	Guaiac.....	I 2-3 gr.	} 1 10
		Ammon. Mur....	I 2-3 gr.	
		Ext. Licorice.....	I-2 gr.	

1 10

One slowly dissolved on the tongue, for astringent and antiseptic effects upon the throat.

<b>GUAIAAC &amp; LICORICE</b> .....		{ Resin Guaiac.....1-2 gr. }	1 20
(Dr. Smith.)		{ Ext. Licorice.....1-2 gr. }	
One every two hours, for tonsillitis.			
<b>GUAIACOL</b> .....	1-2 gr.	<b>Chocolate Coated only</b>	1 75
One to three every three or four hours.			
<b>GUAIACOL</b> .....	1 gr.	<b>Chocolate Coated only</b>	2 50
One to two every three or four hours.			
<b>HÆMATIC</b> .....		{ Acid Arsen.....1-120 gr. }	75
(Dr. Clymer.)		{ Iron Powd.....1-10 gr. }	
		{ Merc. Sub. Cor..1-200 gr. }	85
		{ Nux Vomica Po..1-20 gr. }	
		<b>Chocolate Coated</b> .....	
Give one about an hour after meals, as an hæmatic tonic in anæmia.			
<b>HÆMATIC TONIC</b> .....		{ Strychnine Sulph.1-30 gr. }	3 00
(Dr. Seymour.)		{ Acid Arsenous...1-30 gr. }	
		{ Ext. Aloes.....1-5 gr. }	85
		{ Ext. Hyoscyamus..1-4 gr. }	
		{ Iron Sulph. Dry....2 gr. }	
		{ Quinine Bisulph....1 gr. }	
One after each meal with a swallow of water, in anæmia and chlorosis, and as a general tonic after acute illness; especially those maladies of a depleting and exhaustive nature.			
<b>HÆMOGALLOL</b> .....	3 grs.	<b>Per 100</b> .....	16 50
			1 75
Begin with one after meals; the second week, give two at a dose; if well tolerated, give three for the third week and after; for anæmia, chlorosis and menstrual disorders dependent upon them.			
<b>HEADACHE</b> .....		{ Acetanilid.....1 gr. }	1 15
(Dr. Chamberlain.)		{ Ext. Belladonna..1-8 gr. }	
		{ Ext. Gelsemium..1-4 gr. }	
One or two at first and repeat every two to four hours, as required; for congestive or neuralgic headache, with fever.			
<b>HEADACHE, Improved</b> .....		{ Acetanilid.....1 gr. }	1 85
(Dr. Chamberlain.)		{ Ext. Belladonna..1-8 gr. }	
		{ Ext. Gelsemium..1-4 gr. }	
		{ Caffeine Citrate..1-2 gr. }	
One or two at first, and repeat one every two to four hours, as required.			
<b>HEART STIMULANT</b> ....		{ Tr. Digitalis.....2 min. }	1 75
		{ Tr. Strophanthus..2 min. }	
		{ Tr. Belladonna..1-4 min. }	1 85
		{ Nitroglycerin...1-100 gr. }	
		<b>Chocolate Coated</b> .....	
One every three hours for smoker's heart, palpitation, and angina pectoris.			
<b>HEART STIMULANT No. 2</b> ....		{ Tr. Digitalis.....2 min. }	1 75
		{ Tr. Strophanthus..2 min. }	
		{ Tr. Belladonna..1-4 min. }	1 85
		{ Nitroglycerin...1-200 gr. }	
		<b>Chocolate Coated</b> .....	
One every three hours for smoker's heart, palpitation, and angina pectoris.			
<b>HEART STIMULANT No. 3</b> ....		{ Atropine Sulph...1-75 gr. }	1 25
(Dr. B. F. Horner.)		{ Strych. Sulph....1-30 gr. }	
One tablet every three hours for certain cases of slow heart, attended with valvular lesions at aortic or pulmonary orifices, and of some particular forms at mitral valve.			

HEART STIMULANT No. 4.... (Dr. Durant.)	{ Fl. Ext. Digitalis. . . 1 min.	1 35
	{ Nitroglycerin. . . 1-100 gr.	
	{ Strych. Sulph. . . . 1-50 gr.	
	<b>Chocolate Coated</b>	1 45

One tablet, three or four times a day, when digitalis is indicated in chronic valvular diseases.

HEART STIMULANT No. 5.... (Dr. L. L. Seaman.)	{ Tr. Cact. Grand. 2 1-2 min.	2 50
	{ Tr. Strophanth. 2 1-2 min.	
	{ Tr. Digitalis. . . 2 1-2 min.	
	<b>Plain or Chocolate Coated</b>	

One every three hours, "specially indicated in mitral stenosis or regurgitation."

HEART STIMULANT No. 6.... (Dr. Wood.)	{ Tr. Digitalis. . . . . 1 min.	1 25
	{ Tr. Strophanth. . . . 1 min.	
	{ Strych. Sulph. . . . 1-30 gr.	

One tablet three times a day as a heart stimulant.

HEART STIMULANT No. 7....	{ Digitalis. . . . . 1-100 gr.	2 50
	{ Nitroglycerine. . 1-200 gr.	
	{ Strych. Sulph. . . . 1-60 gr.	

One tablet three or four times a day, as a heart stimulant.

HEART TONIC .....	{ Fl. Ext. Cac. Grand. 5 min.	6 50
	{ Tinct. Digitalis. . . 2 min.	
	{ Caffeine Citrate. . . 2 gr.	

One every three hours for weak heart or feeble heart of aged.

HEART TONIC No. 2..... (Dr. Gordinier.)	{ Nitroglycerin. . . 1-200 gr.	2 00
	{ Tr. Strophanthus. . 3 min.	
	{ Fl. Ext. Digitalis. . 1 min.	
	{ Strych. Sulph. . . . 1-60 gr.	
	{ Iron Powd. . . . . 1 gr.	
	<b>Plain or Chocolate Coated</b>	

Give one three times daily, or if occasion requires, every three or four hours, as a general heart tonic, and tonic in anæmia.

HEART TONIC No. 3..... (Dr. Pultz.)	{ Tr. Digitalis. . . . . 1 min.	1 50
	{ Tr. Strophanthus. . 1 min.	
	{ Tr. Cactus Grand. . 1 min.	
	{ Sparteine Sulph. . 1-100 gr.	
	{ Nitroglycerin . . . 1-100 gr.	
	<b>Chocolate Coated</b>	1 60

One tablet every three hours for irregular and weak heart.

HEART TONIC No. 5.....	{ Digitalis. . . . . 1 1-2 gr.	3 00
	{ Scammony Po. . . 1 1-2 gr.	
	{ Squill Powd. . . . 1 1-2 gr.	

One every day for four days.

HELLEBOREIN.....	1-100 gr. ....	1 50
------------------	----------------	------

Cardiac tonic.

One every two hours, until the pulse is under control, then one every four to six hours, as a heart tonic where digitalis disagrees.

HELLEBOREIN.....	1-50 gr. ....	2 60
------------------	---------------	------

A tablet two or three times a day, to strengthen cardiac systole and as a general vaso-motor stimulant; increase with caution.

HELONIN .....	1-10 gr. ....	1 30
---------------	---------------	------

Diuretic and tonic.

One three times a day to increase the flow of urine, and as an emmenagogue.

When ordering or prescribing

HELONIAS COMP.....	Ext. Helonias.....	1-2 gr.	4 00
	Acid Salicylic.....	1 gr.	
	Acid Boric.....	3 gr.	
	Acid Tannic.....	2 gr.	
	Ext. Opium.....	1-2 gr.	
	Ext. Hyoscyamus...	1 gr.	
	Alum.....	3 gr.	
	Eucalyptol.....	q. s.	
	Thymol.....	q. s.	

Insert one tablet into the vagina at night, after hot vaginal douche. For inflammation of the vagina, uterus and appendages, and catarrhal conditions of vaginal mucous membrane.

HEMICRANIA .....	Croton Chloral.....	1 gr.	3 50
	Ext. Can. Indica....	1-4 gr.	

Give one tablet and another in half an hour; for nervous or sick headache.

HEPATIC .....	Euonymin.....	1-8 gr.	1 15
	Podophyllin.....	1-20 gr.	
	Ipecac Powd.....	1-8 gr.	
	Calomel.....	1-8 gr.	
	Aloin.....	1-12 gr.	
	Chocolate Coated.....		1 25

HEPATIC No. 2.....	Ext. Physostigma...	1-2 gr.	10 50
	Podophyllin.....	1-8 gr.	
	Euonymin.....	2 gr.	
	Ext. Colocy Co....	1 gr.	
	Ext. Hyoscyam...	1-4 gr.	

One or two at night for habitual constipation, and inaction of the liver and large intestine.

HEPATIC No. 3 .....	Ext. Physostig...	1-12 gr.	2 50
	Podophyllin.....	1-8 gr.	
	Mass. Hydrarg...1-2 gr.		
	Ext. Colocy. Co....	1 gr.	
	Ext. Hyoscyam...1-2 gr.		

One or two at a dose, at bedtime, three or four times a week, for congested liver and hepatic torpidity or inaction.

HEPATIC ALTERATIVE.....	Bismuth Subnit.....	1 gr.	2 50
	Salol.....	1 gr.	
	Calomel.....	1-10 gr.	

One tablet every hour until laxative effect is produced, then two or three times a day for a week or more; for bilious dyspepsia with flatulence.

HEPATIC COMPOUND No. 4..	Hydrastin.....	1-4 gr.	3 00
	Asclepin.....	1-4 gr.	
	Podophyllin.....	1-4 gr.	
	Leptandrin.....	1-4 gr.	

One or two tablets as a mild laxative; three to five as a purge in chronic or acute constipation, hepatitis and lumbago.

HEPATO HYDRARG .....	Podophyllin.....	1-8 gr.	2 25
	Ext. Belladonna...	1-4 gr.	
	Blue Mass.....	3 gr.	

One or two at night, followed by a saline next morning. When two are given, one should be taken an hour after the other. Used when a decided action of the liver and bowels is desired.

HEROIN .....	1-20 gr.	1 85
--------------	----------	------

One tablet for the aged, three times a day, and at bedtime, to allay coughs and pains in the chest in bronchitis, pharyngitis and angina.

<b>HEROIN</b> .....	1-12 gr.	2 50
One to two tablets three times a day, and at bedtime, to allay coughs and pains in the chest in bronchitis, pharyngitis and angina.		
<b>HEROIN</b> .....	1-6 gr.	4 00
One tablet three times a day, and at bedtime, to allay coughs and pains in the chest in bronchitis, pharyngitis and angina.		
<b>HEROIN</b> .....	1-5 gr.	4 75
One tablet three times a day, and at bedtime, to allay coughs and pains in the chest in bronchitis, pharyngitis and angina.		
<b>HEROIN</b> .....	1-2 gr.	10 25
One tablet at a dose; repeat cautiously to avoid nausea.		
<b>HORSE CHESTNUT TINCT.</b> ...	10 min	1 50
One every three hours, continuing for three or four weeks, for piles.		
<b>HYDRARG. COMP. No. 1.</b> .....	<div> <div>Calomel.....1-24 gr.</div> <div>Tartar Emetic...1-120 gr.</div> <div>Nitr. Potass....2 1-2 gr.</div> <div>Ext. Licorice... ..</div> <div>Oil Sassafras.....</div> <div>Oil Wintergreen..aa. q.s.</div> </div>	1 10
(Dr. Leaming.)		
Bronchial expectorant.		
One every three hours for cough after grip.		
<b>HYDRARG. COMP. No. 2</b> .....	<div> <div>Calomel.....1-12 gr.</div> <div>Tartar Emetic...1-60 gr.</div> <div>Nitr. Potass....5 gr.</div> <div>Ext. Licorice.....</div> <div>Oil Sassafras.....</div> <div>Oil Wintergreen..aa. q.s.</div> </div>	1 50
(Dr. Leaming.)		
Give a tablet every four hours as an expectorant in bronchitis and influenza.		
<b>HYDRARG. COMP. No. 3</b> .....	<div> <div>Calomel.....1-6 gr.</div> <div>Tartar Emetic...1-30 gr.</div> <div>Nitr. Potass....10 gr.</div> <div>Ext. Licorice.....</div> <div>Oil Sassafras.....</div> <div>Oil Wintergreen..aa. q.s.</div> </div>	2 00
(Dr. Leaming.)		
One after meals, for chronic inflammation of the bronchial tubes with thick and scanty sputa, bronchial dilatation and emphysema.		
<b>HYDRARG. &amp; IPECAC COMP.</b> ..	<div> <div>Mass. Hydrarg.....1 gr.</div> <div>Ipecac Powd....1-8 gr.</div> <div>Opium Powd ....1-4 gr.</div> </div>	1 50
Alterative and hepatic.		
One after meals with an occasional saline in the morning, for loss of function in the liver, with biliousness, headache and constipation.		
<b>HYDRARG. &amp; PODOPHYLLIN</b> {	<div> <div>Mass. Hydrarg.....1 gr.</div> <div>Podophyllin .....1-4 gr.</div> </div>	1 10
Hepatic.		
One to four at bedtime, as an hepatic laxative in duodenal catarrh and intestinal dyspepsia.		
<b>HYDRASTIA, (White Alkaloid.)</b>	1-20 gr.	1 75
Stimulant.		
One or two after meals for dyspepsia, for a stimulant effect on the involuntary muscular fibres of intestines.		
<b>HYDRASTIN.</b> .....	1-100 gr.	75
One or two after meals for dyspepsia, for a stimulant effect on the involuntary muscular fibres of intestines.		

When ordering or prescribing

<b>HYDRASTIN</b> .....	1-20 gr.....	1 10
One or two after meals for dyspepsia, for a stimulant effect on the involuntary muscular fibres of intestines.		
<b>HYDRASTIN MUR</b> .....	1-100 gr.....	75
One or two after meals for dyspepsia, for a stimulant effect on the involuntary muscular fibres of intestines.		
<b>HYDRASTIN CAPSICUM CO.</b> ..	{ Hydrastin..... 1-4 gr. } { Podophyllin..... 1-10 gr. } { Capsicum Powd... 1-4 gr. }	2 00
(Dr. Postle.)		
One to three tablets a day, between meals, as a stomachic tonic and laxative, and also as an appetizer.		
<b>HYDRASTIN COMP</b> .....	{ Podophyllin..... 1-20 gr. } { Hydrastin..... 1-10 gr. }	1 25
One every three hours, for not more than three doses, to regulate the intestines in atonic conditions, decreasing to one or two at night.		
<b>HYOSCINE HYDROBROM</b> ....	1-200 gr.....	3 40
Anodyne.		
One every hour or two, as necessary to quiet the nerves and induce sleep, and to supplement the action of other hypnotics.		
<b>HYOSCINE HYDROBROM</b> ....	1-100 gr.....	6 00
One tablet every hour for three doses, to control the paroxysms of asthma and for neuralgia. Must be given with caution in the case of a weak heart.		
<b>HYOSCINE HYDROBROM</b> ....	1-50 gr.....	11 00
One as a nerve sedative, for mania, various neurotic conditions, asthma, and as an anodyne in pain; repeated with caution.		
<b>HYOSCYAMIA, (White Cryst.)</b> ..	1-500 gr.....	2 00
A tablet every two hours, for cough with pain and restlessness.		
<b>HYOSCYAMIA, (White Cryst.)</b> ..	1-100 gr.....	2 40
One tablet repeated in an hour, as an hypnotic and sedative.		
<b>HYOSCYAMIA, (White Cryst.)</b> ..	1-300 gr.....	3 00
<b>HYOSCYAMIA, (White Cryst.)</b> ..	1-200 gr.....	3 75
<b>HYOSCYAMIA, (White Cryst.)</b> ..	1-100 gr.....	7 00
<b>HYOSCYAMIA, (White Cryst.)</b> ..	1-50 gr.....	12 50
<b>HYOSCYAMUS COUGH</b> .....	{ Codeine..... 1-16 gr. } { Ext. Hyoseyam... 1-8 gr. }	2 25
One every three hours in cough accompanied by pain or a strong nervous element.		
<b>HYOSCYAMUS EXTRACT</b> ....	1-4 gr.....	75
Sedative.		
One tablet every four hours, for vesical tenesmus and dysuria, and cough with pain.		
<b>HYOSCYAMUS EXTRACT</b> ....	1-2 gr.....	1 15
A tablet three times a day to control the pain of inflammation in the respiratory and uro-poietic systems and as a general nerve sedative.		
<b>HYPNOTIC</b> .....	{ Sodium Bromide.... 5 gr. } { Acetanilid..... 2 gr. } { Hyoscyamia.... 1-400 gr. } { Digitalia..... 1-400 gr. }	3 50
One every two hours to produce sleep in threatened delirium and mania, and for the pain of various inflammations.		

<b>HYPOPHOSPHITES &amp; QUININE</b> .....		Quinine Hypo.....1 gr.	}	3 50
Tissue tonic.		Ferrum Hypo.....1-2 gr.		
		Calcium Hypo....1-2 gr.		
		Sodium Hypo....1-4 gr.		
		Potass. Hypo....1-4 gr.		
		Mangan. Hypo....1-4 gr.		
		Strych. Hypo....1-64 gr.		
<b>Plain or Chocolate Coated</b>				
A tablet after meals for anæmia, and in convalescence from wasting disease.				
<b>HYPOPHOSPHITES &amp; QUININE</b> .....		Quinine Hypo.....1-2 gr.	}	2 25
One-half strength.		Ferrum Hypo.....1-4 gr.		
		Calcium Hypo....1-4 gr.		
		Sodium Hypo....1-8 gr.		
		Potass. Hypo....1-8 gr.		
		Mangan. Hypo....1-8 gr.		
		Strych. Hypo....1-128 gr.		
<b>Plain or Chocolate Coated</b>				
For a child, one after meals, for anæmia and loss of flesh.				
<b>HYPOPHOSPHITES &amp; QUININE WITH CREOSOTE</b> .....		Quinine Hypo.....1 gr.	}	3 75
		Ferrum Hypo.....1-2 gr.		
		Calcium Hypo....1-2 gr.		
		Sodium Hypo....1-4 gr.		
		Potass. Hypo....1-4 gr.		
		Mangan. Hypo....1-4 gr.		
		Strych. Hypo....1-64 gr.		
<b>Plain or Chocolate Coated</b>				
One after meals, for beginning phthisis, to prevent emaciation and loss of strength, or as a general chemical food and intestinal antiseptic.				
<b>HYSTERIA</b> .....		Iron Sulph. Dry....1 gr.	}	3 00
		Ext. Sumbul.....1 gr.		
		Asafœtida.....2 gr.		
		Acid Arsenous...1-100 gr.		
		Apomorph. Mur..1-50 gr.		
<b>Chocolate Coated only</b>				
One repeated every hour for three or four doses for hysteria and nervousness. Better taken with a draught of water.				
<b>IGNATIA POWD.</b> .....		1-100 gr		75
Tonic.				
One to a child every two hours as a stomachic and general tonic.				
<b>IGNATIA POWD.</b> .....		1-20 gr		75
One tablet every two hours, as a gastric stimulant and nerve tonic.				
<b>IGNATIA POWD.</b> .....		1-10 gr		75
One tablet every two hours, as a gastric stimulant and nerve tonic.				
<b>IGNATIA COMP.</b> .....		Ignatia Powd...1-100 gr.	}	75
(Dr. Gilbert.)		Ipecac Powd....1-10 gr.		
One an hour after meals, for nervous dyspepsia.				
<b>INCONTINENCE</b> .....		Tr. Belladonna....2 min.	}	2 00
		Tr. Cubebs.....2 min.		
		Tr. Nux Vomica..1 min.		
		Tr. Rhub. Arom..1 min.		
		Tr. Cascarilla..2 min.		
<b>Plain or Chocolate Coated</b>				
One at bed time for a child six to seven years old.				
<b>INDIGESTION</b> .....		Nux Vomica....1-50 gr.	}	75
(Dr. Gilbert.)		Carbo Veg.....q. s.		
One tablet every two hours for indigestion from excessive eating or drinking.				

When ordering or prescribing

INDIGESTION No. 3. . . . .	{ Alpha Naphtol. . . 2 1-2 gr. }	5 00
	{ Bism. Subcarb. . . 1 1-2 gr. }	
	{ Rhubarb Powd. . . . 1 gr. }	
	{ Codeine . . . . . 1-20 gr. }	
	{ Ext. Bellad. . . . . 1-20 gr. }	
	{ Ext. Nux Vom. . . 1-20 gr. }	

Two tablets immediately after each meal followed by one an hour after, and two more at bedtime; for intestinal indigestion.

INDIGESTION POWD. No. 1. . . . .	{ Pepsin Sacch. . . . 1-40 gr. }	75
	{ Carbo Veg. . . . . 1-40 gr. }	
	{ Bismuth Subnit. . . 1-40 gr. }	

A tablet crushed in a teaspoonful of sweetened water, for an infant with dyspepsia and vomiting; give fifteen minutes before and directly after feeding.

INDIGESTION POWD. No. 2. . . . .	{ Carbo Veg. . . . . 1-400 gr. }	75
	{ Rhubarb Powd. . . 1-40 gr. }	
	{ Bismuth Subnit. . . 1-40 gr. }	
	{ Pepsin. . . . . 1-40 gr. }	

One after feeding, for the fermentative and acid dyspepsia of nursing infants; crush the tablet in a teaspoonful of water, which may be sweetened if desired.

INFLAMMATUS. . . . . (Dr. Seward.)	{ Tr. Verat Vir. . . 1-12 min. }	1 75
	{ (Norwood's) }	
	{ Tr. Aconite Rt. . 1-12 min. }	
	{ Tr. Opium . . . . . 2-3 min. }	
	{ Tr. Bryonia . . . . 2-3 min. }	
	{ Tr. Belladonna. . 2-3 min. }	
	{ Tr. Cimicifuga. . 2-3 min. }	
	{ Tr. Guaiac. . . . . 2-3 min. }	
	{ Tr. Colch. Seed. . 2-3 min. }	
	{ Tr. Colocynth. . . 2-3 min. }	

One every two to four hours, or one dissolved in fifteen teaspoonfuls of water, of this a teaspoonful every fifteen minutes; for acute stages of rheumatism or neuralgia from exposure, also for early stages of tonsillitis.

INFLUENZA . . . . . (Dr. S. S. Jones.)	{ Atropine Sulph. . 1-1000 gr. }	1 35
	{ Morphine Sulph. . 1-32 gr. }	
	{ Camphor. . . . . 1-2 gr. }	

One every hour for four hours, or three every two or three hours.

INFLUENZA No. 2. . . . . (Dr. Curtis.)	{ Strych. Arsen. . . 1-132 gr. }	1 15
	{ Atropine Sulph. . 1-300 gr. }	
	{ Camphor. . . . . 1-4 gr. }	
	{ Ipecac Powd. . . . 1-20 gr. }	

One every hour, for fresh colds, until mouth becomes dry.

INFUSION BUCHU COMP. . . . . Each tablet represents	{ Infusion Buchu . . . . 3 ii. }	1 75
	{ Tr. Hyoscyamus. . . 20 min. }	
	{ in combination with }	
	{ Potass. Bicarb. . . . grs. vi. }	

One tablet every four hours, dissolved in a glass of hot water, for cystitis and catarrh of the bladder.

INFUSION GENTIAN COMP. . . . .	1 25
--------------------------------	------

Each tablet represents one drachm Compound Infusion Gentian, U. S. P., 1870.

From two to four tablets after each meal as a stomachic and carminative.

INJECTION..... (Dr. Briggs.)	{ Hydrastin... ..1-5 gr. }	2 75
	{ Zinc Sulph... ..3 gr. }	
	{ Morph. Sulph... 1-20 gr. }	
	{ Atropine Sulph... 1-50 gr. }	

Dissolve one tablet in half a glass of warm water and make three injections carefully after each urination; warm the solution for each using, for best results.

INJECTION, CUPRIC.....	{ Lead Acetate.....2 gr. }	2 00
	{ Acid Tannic.....2 gr. }	
	{ Zinc Sulphate.....2 gr. }	
	{ Copper Sulphate....2 gr. }	

One in two to four ounces of water, for urethral injection.

INSOMNIA.....	{ Morph. Brom....1-8 gr. }	6 65
	{ Hyoscin Hydrb. 1-150 gr. }	
	{ Camphor Monobr...2 gr. }	

One at bedtime, and repeat in half an hour if necessary, for insomnia and restlessness.

IPECAC POWD.....	1-1000 gr..	75
------------------	-------------	----

One every three hours for an infant, as a gentle stimulant to the gastric and intestinal mucous membrane.

IPECAC POWD.....	1-200 gr.....	75
------------------	---------------	----

One every three hours, for the atonic dyspepsia of anæmic children.

IPECAC POWD.....	1-100 gr.....	75
------------------	---------------	----

A tablet three or four times a day, for cough and bronchitis in children, as a stimulating expectorant.

IPECAC POWD.....	1-50 gr.....	75
------------------	--------------	----

One every hour, for diarrhœa and dysentery in children.

IPECAC POWD.....	1-10 gr.....	75
------------------	--------------	----

One every three hours for a child, as a vascular and hepatic stimulant; every hour as a sudorific and expectorant in diseases of the respiratory organs.

IPECAC POWD....	1-4 gr.....	90
-----------------	-------------	----

Emetic and expectorant.

As an emetic for an infant, one repeated if necessary; one every four hours as an expectorant in cough.

IPECAC POWD.....	1-2 gr.....	1 15
------------------	-------------	------

Give a tablet, repeated in thirty to sixty minutes, as an emetic in croup, with draughts of warm water.

IPECAC POWD.....	5 grs.....	3 00
------------------	------------	------

Emetic.

One tablet with warm water, as an emetic; if it fails to act promptly repeat in an hour, with the addition of a teaspoonful of mustard.

IPECAC FL. EXT.....	1 min.....	1 50
---------------------	------------	------

One every two hours, for colds and to promote sweating in fevers; one after meals as an hepatic stimulant; useful in bilious headache.

IPECAC WINE.....	5 min.....	1 00
------------------	------------	------

One, repeated in two hours, for sick-headache and migraine, and as a stimulant to the alimentary canal.

IRON ALBUMINATE.....	3 grs.....	2 25
----------------------	------------	------

Nutritive tonic.

One after meals, for anæmia in children.

When ordering or prescribing



IRON, ARSENIC & STRYCH- NINE No. 5. ....	{ Iron Powd. .... 1 gr. } Acid Arsenous. .... 1-20 gr. } Strych. Sulph. .... 1-30 gr. } Chocolate Coated. ....	75 85
One after each meal and at bed hour, for general tonic, loss of appetite and anæmia.		
IRON, ARSENIC & STRYCH- NINE No. 6. ....	{ Iron Pyrophos. .... 1 gr. } Acid Arsen. .... 1-100 gr. } Strych. Sulph. .... 1-60 gr. } Chocolate Coated. ....	75 85
One tablet three times a day, ten minutes before meals; a useful tonic in all kinds of anæmia, with loss of appetite.		
IRON, ARSENIC & STRYCH- NINE No. 7. ....	{ Iron Pyrophos. .... 1 gr. } Acid Arsen. .... 1-50 gr. } Strych. Sulph. .... 1-60 gr. } Chocolate Coated. ....	75 85
One after each meal and at bed hour, as a general tonic and for loss of appetite.		
IRON CARBON. FRASER'S No. 2. ....	{ Ferri Carb. .... 5 gr. } Sacch. Alb. .... 2 1-2 gr. }	1 50
One every three hours, as a tonic, and after wasting diseases.		
IRON CITRATE. ....	1-2 gr. ....	90
A tablet three times a day where a mild hæmatic tonic, with a slightly diuretic effect is required.		
IRON CITRATE. ....	1 gr. ....	1 10
One after meals, as a gentle tonic and hæmatic.		
IRON LACTATE. ....	1-2 gr. ....	75
A tablet two or three times a day, for a child with sensitive stomach, as a tonic; increase gradually as tolerance is secured.		
IRON LACTATE. ....	1 gr. ....	1 00
One after meals where an iron tonic is required, with the least distress to the digestive organs.		
IRON LACTATE. ....	2 grs. ....	1 25
One three times a day, for anæmia, dysmenorrhœa and amenorrhœa; gradually increase the dose.		
IRON LACTATE COMP. .... (Dr. Bermann.)	{ Ext. Trifolium. .... 0.05 } Lactate Iron. .... 0.05 }	1 25
One tablet a quarter of an hour before eating, as a mild bitter tonic and hæmatic; increase to three after meals, by adding one a day.		
IRON & MANGANESE COMP..	{ Iron Powd. .... 1 gr. } Acid Arsen. .... 1-50 gr. } Strych. Sulph. .... 1-60 gr. } Mangan. Binox. .... 1 gr. } Chocolate Coated. ....	1 25 1 35
One or two three times a day after meals in amenorrhœa, anæmia and chlorosis.		
IRON, MERCURY COMP.. .... Alterative, tonic.	{ Iron Powd. .... 1 gr. } Corrosive Sub. .... 1-60 gr. } Strych. Sulph. .... 1-60 gr. } Acid Arsenous. .... 1-60 gr. } Chocolate Coated. ....	75 85
One, three times a day as an alterative, hepatic and general tonic.		
IRON, MERCURY COMP. No. 2	{ Iron Powd. .... 1 gr. } Corrosive Sub. .... 1-50 gr. } Acid Arsen. .... 1-100 gr. }	75
One after eating, for chronic disease with anæmia and emaciation.		

When ordering or prescribing

IRON MERCURY CO. No. 4.....	{ Iron Pyrophos.....1 gr.	75
	{ Acid Arsenous...1-50 gr.	
	{ Strych. Sulph....1-60 gr.	
	{ Corrosive Sub....1-60 gr.	
	Chocolate Coated.....	85

One, three times a day as an alterative, hepatic and general tonic.

IRON & MERCURY COMP. WITH QUININE.....	{ Iron Powd.....1 gr.	1 50
	{ Mercury Bichlor.1-60 gr.	
	{ Acid Arsenous...1-60 gr.	
	{ Strych. Sulph....1-60 gr.	
	Quinine Sulph....1-2 gr.	
	Chocolate Coated.....	1 60

One to four daily, for malarial cachexia, dumb ague, anæmia and general debility.

IRON PHOSPHORUS COMP.. Nerve tonic.	{ Iron Powd.....1 gr.	1 10
	{ Acid Arsenous...1-40 gr.	
	{ Strych. Sulph....1-60 gr.	
	{ Phosphorus....1-100 gr.	
	Chocolate Coated.....	1 20

One, three times a day after meals, for nerve and general tonic.

IRON POWDER.....	1-1000 gr.....	75
------------------	----------------	----

One after each nursing, crushed in a little sweetened milk, for an infant, as a tonic in anæmia and after wasting diarrhoea.

IRON POWDER.....	1-500 gr.....	75
------------------	---------------	----

One after each nursing, crushed in a little sweetened milk, for an infant.

IRON POWDER.....	1-200 gr.....	75
------------------	---------------	----

A tablet after each nursing, for an infant with anæmia and a tendency to diarrhoea.

IRON POWDER.....	1-100 gr.....	75
------------------	---------------	----

A tablet every three hours for a child.

IRON POWDER.....	1-50 gr.....	75
------------------	--------------	----

One every three hours, increasing every day by one until three or four are taken at a dose.

IRON POWDER.....	1-10 gr.....	75
------------------	--------------	----

One every three hours for anæmia in childhood.

IRON POWDER.....	1-5 gr.....	75
------------------	-------------	----

One after meals, for pale and sickly children.

IRON POWDER.....	1-2 gr.....	75
------------------	-------------	----

One, three or four times a day, for anæmia, amenorrhœa and allied disorders.

IRON POWDER.....	1 gr.....	75
------------------	-----------	----

A tablet three or four times a day, for anæmia, anæmic headache and in convalescence.

IRON POWDER.....	2 grs.....	1 10
------------------	------------	------

Three to four daily, for anæmia and various cachexia.

IRON PYROPHOSPHATE.....	1-10 gr.....	75
-------------------------	--------------	----

One every two hours for anæmic diseases of childhood, especially rhachitis.

IRON PYROPHOSPHATE.....	1-2 gr.....	75
-------------------------	-------------	----

One or two after meals, for rhachitic and tubercular disease in childhood.

IRON PYROPHOSPHATE.....	1 gr.....	75
-------------------------	-----------	----

A tablet three times a day, for a tonic in malnutrition and catamenial disorders.

IRON PYROPHOS COMP..... Modified iron tonic.	{ Iron Pyrophos.....1 gr. }	1 60
	{ Quinine Sulph.....1-2 gr. }	
	{ Strych. Sulph.....1-60 gr. }	
	{ Acid Arsenous...1-30 gr. }	
	Chocolate Coated.....	1 70

One after meals, as an hæmatic and alterative tonic.

IRON, QUININE & ALOES CO. (Dr. Duncan.)	{ Iron Powd.....2 gr. }	3 50
	{ Quinine Sulph.....2 gr. }	
	{ Strych. Sulph.....1-40 gr. }	
	{ Acid Arsenous...1-40 gr. }	
	Aloes Powd.....1-8 gr. }	
	Plain or Chocolate Coated	

One every three hours in chronic malaria.

IRON, QUININE & STRYCH- NINE..... Tonic, antiperiodic.	{ Iron Powd.....1-2 gr. }	1 50
	{ Quinine Sulph....1-2 gr. }	
	{ Strych. Sulph....1-120 gr. }	
	Chocolate Coated.....	

One or two after meals, as an antimalarial and general tonic.

IRON, QUININE & STRYCH- NINE No. 2.....	{ Iron Reduced.....1 gr. }	2 25
	{ Quin. Sulph.....1 gr. }	
	{ Strych. Sulph....1-60 gr. }	
	Plain or Chocolate Coated	

One, three times a day, after meals and at bedtime, in anæmia, chlorosis, and as a general tonic.

IRON & STRYCHNINE..... Tonic.	{ Iron Powd.....1 gr. }	75
	{ Strych. Sulph....1-60 gr. }	

Two or three tablets a day, in nervous and anæmic diseases of adolescence.

IRON & STRYCHNINE COMP (Dr. Morrill.) Tonic, antiperiodic.	{ Iron Powd.....1-2 gr. }	1 50
	{ Quinine Sulph....1-2 gr. }	
	{ Acid Arsen....1-100 gr. }	
	{ Strych. Sulph...1-120 gr. }	
	Chocolate Coated.....	1 60

One or two, according to age, three times a day, after meals. Used in chronic malarial poisoning, malarial anæmia, neuralgias, chlorosis and the amenorrhœa of anæmia.

IRON & STRYCHNINE CIT...1-2 gr.....	00
-------------------------------------	----

General nerve tonic.

One after meals, as an hæmatic and nerve tonic in chorea, epilepsy, cerebral and spinal anæmia, and hysteria.

IRON & STRYCHNINE CIT...1 gr.....	1 15
-----------------------------------	------

One after meals, as an hæmatic and nerve tonic in chorea, epilepsy, cerebral and spinal anæmia, and hysteria.

IRON SULPH. Dried.....1 gr.....	1 00
Chocolate Coated only	

One after each meal for an iron tonic.

JALAP.....1-2 gr.....	75
-----------------------	----

One every half hour for five doses, then every hour until laxative effect is produced.

JALAP.....1 gr.....	1 10
---------------------	------

One every half hour for five doses, then every hour until laxative effect is produced.

KALI COMP.....	{ Kali Bichrom...1-100 gr. }	75
	{ Mercury Biniod.1-100 gr. }	

One every two hours, for sore throat from glandular irritation.

When ordering or prescribing

<b>KRAMERIA COMP.</b> ..... (Dr. J. C. Applegate.) Diarrhœa.	{ Fl. Ext. Krameria . . 1 min. Bismuth Subnit. . . . 1 gr. Opium Powd. . . . . 1-4 gr. Licorice R't Po. . . . 1-4 gr. }	1 90
--	--	------

One every hour, for diarrhœa and dysentery; decrease to one every three or four hours as symptoms ameliorate; valuable also in intestinal inflammation, used in conjunction with, or following salines.

<b>KRAMERIA COMP. No. 2.</b> .....	{ Fl. Ext. Krameria 1-2 min. Bismuth Subnit. . . . 1 gr. Opium Powd. . . . . 1-25 gr. }	1 6c
------------------------------------	---	------

One tablet every two or three hours powdered in sugar, for the summer diarrhœa of children; allow only boiled milk until the movements become normal.

<b>LACTO FERRIC</b> ..... Tonic.	{ Iron Lactate. . . . . 1 gr. Cinchonia Sulph. . . . 1 gr. Ext. Hyoscyam. . . . 1-5 gr. Acid Arsenous. . . . 1-30 gr. Strych. Sulph. . . . . 1-60 gr. }	1 50
	<b>Chocolate Coated</b> .....	1 60

A tablet three or four times a day, as a tonic in malnutrition and anæmia with neuroses and ataxia.

<b>LARYNGITIS</b> .....	{ Potass. Brom. . . . 1 1-4 gr. Potass. Chlorate. . . 5-8 gr. Ipecac Powd. . . . . 1-12 gr. Ext. Licorice. . . . . 1 1-4 gr. Squill Powd. . . . . 1-2 gr. }	1 15
-------------------------	---	------

One every hour, for continued cough, with tickling in the throat.

<b>LAXATIVE</b> .....	{ Aloin . . . . . 1-4 gr. Ext. Belladonna. . . 1-8 gr. Strych. Sulph. . . . . 1-60 gr. Ipecac Powd. . . . . 1-16 gr. }	1 15
	<b>Chocolate Coated</b> .....	1 25

One or two at night, as a stimulant to the alimentary canal and to increase peristalsis.

<b>LAXATIVE No. 2</b> ..... (Dr. Hauck.)	{ Quinine Sulph. . . . . 1 gr. Ext. Rhubarb. . . . . 1 gr. Ipecac Powd. . . . . 1-8 gr. Capsicum Powd. . . . 1-4 gr. }	3 00
	<b>Plain or Chocolate Coated</b>	

One or two at a dose every two or three hours, as required for catarrh of the stomach, and dyspepsia, combined with mild forms of constipation.

<b>LAXATIVE No. 3</b> ..... (Dr. Burleigh.)	{ Aloes. . . . . 1 gr. Ext. Belladonna. . . 3-20 gr. Ext. Physostig. . . . 3-20 gr. Ext. Nux Vom. . . . . 1-5 gr. Ext. Cascara. . . . . 1-5 gr. }	2 00
	<b>Plain or Chocolate Coated</b>	

One or two tablets at night, repeated the next morning if necessary, for constipation.

<b>LAXATIVE No. 4</b> ..... (Dr. A. M. Thomas.)	{ Ext. Cascara Sag. . . 1 1-2 gr. Sodium Bicarb. . . . 1 1-4 gr. Hydrastin. . . . . 1-200 gr. Ext. Nux Vom. . . . . 1-8 gr. Ext. Belladonna. . . 1-16 gr. Oil Peppermint. . . 1-40 min. }	2 50
	<b>Plain or Chocolate Coated</b>	

One to four at night, as required as a laxative. Can be taken indefinitely without producing habit.

<b>LAXATIVE No. 5</b> .....	{ Aloin.....1-4 gr. } { Ext. Belladon....1-16 gr. } { Strych. Sulph....1-60 gr. } { Ipecac Powd....1-16 gr. }	1 15
(Dr. W. B. Pierson.)	<b>Chocolate Coated</b> .....	1 25

One or two tablets at night, repeated the next morning if necessary, for constipation.

<b>LAXATIVE No 6</b> .....	{ Ext. Colocy. Co.1 1-2 gr. } { Jalapin.....1-5 gr. } { Pulv. Scammony...3-4 gr. } { Ext. Hyoscyam....1 gr. } { Podophyllin.....1-30 gr. }	3 35
----------------------------	--	------

One or two tablets at night for constipation; take one the next morning if necessary.

<b>LAXATIVE, CHILDREN'S</b> .....	{ Rhubarb Powd..1-10 gr. } { Sulphur.....1-5 gr. } { Sodium Phos.Dried.1 gr. } { Oil Peppermint.1-20 min. }	1 20
(Dr. E. Chapin.)		

One to three tablets three times daily for children over one year. Infants, one or two tablets a day.

<b>LEAD ACETATE</b> .....	1 gr.....	75
---------------------------	-----------	----

For solution.

One tablet dissolved in a tablespoonful of water makes a solution of about one part in 250; for external use and as an injection.

<b>LEAD &amp; BISMUTH COMP</b> .....	{ Lead Acetate.....1 gr. } { Bismuth Subnit....2 gr. } { Camphor.....1-8 gr. } { Opium Powd.....1-4 gr. }	2 25
--------------------------------------	--	------

Give a tablet every hour for three or four doses, in bilious and flatulent diarrhœa; one after meals for gastric catarrh.

<b>LEAD AND OPIUM WASH</b> ...	{ Lead Acetate.....5 gr. } { Ext. Opium.....1-4 gr. } { Ammon. Chlor.....4 gr. }	2 25
--------------------------------	--	------

For solution. For external use.

For making lead and opium wash, dissolve one tablet in two tablespoonfuls, or in one ounce of water.

<b>LEPTANDRA COMP.</b> .....	{ Cascara Fl.Ext.7 1-2 min. } { Leptan. Fl. Ext.7 1-2 min. }	
	<b>Plain or Chocolate Coated</b>	2 50

One or two every other night until constipation is cured.

<b>LEUCORRHŒA</b> .....	{ Ext. Witch Hazel.1-2 gr. } { Ext. Hyoscyamus...1 gr. } { Ext. Hydras. Color.1 gr. } { Ext. Helonias....1-2 gr. } { Alum.....1 gr. } { Acid Tannic.....1 gr. } { Acid Boric.....8 gr. } { Eucalyptol } { Thymol } { Menthol.....aa q. s. }	3 25
-------------------------	--	------

For external use.

Use one tablet night and morning, follow with douche of hot water.

<b>LIME &amp; IRON PHOSPHATES</b> { Lime Phosphate....2 gr. } (Dr. Hubbard.) { Iron Phosphate....2 gr. } { Sugar.....1 gr. } { Oil Wintergr'n.1-10 min. }	1 50
--	------

One tablet to a child; two to an adult, after each meal, as a reconstructive tonic.

**LIME WATER..... 1 00**

For the preparation of Lime Water, add six tablets to a half pint of boiled water in a clean bottle, and allow to stand with occasional agitation until the tablets disintegrate. Shake again and set aside until the excess of lime has settled and the mixture becomes clear. Decant the clear liquid if desired and keep well bottled for use.

**LITHIUM BENZOATE.....1 gr..... 1 75**

One every three hours, for secondary rheumatic affections.

**LITHIUM CARBONATE.....1 gr..... 1 75**

Antilithic.

One every hour, for acute rheumatism; one to three after meals to hasten the excretion of uric acid.

**LITHIUM CARBONATE.....5 grs..... 3 50**

Antilithic.

One or two after each meal, for gout, arthritis deformans, gravel and vesical calculus.

**LITHIUM CARBONATE..... 3 50**

**EFFERVESCENT.**

Each tablet represents three grains lithium salt.

A tablet three or four times a day, in a glass of water, for rheumatism and gout and the uric acid diathesis.

**LITHIUM SALICYLATE .....5 grs..... 4 00**

Antiseptic alkaline.

One, three or four times a day, for the uric acid diathesis, muscular rheumatism and dyspepsia of gouty subjects.

<b>LIVER.....</b>	{	Calomel.....1-8 gr.	} <b>1 35</b>
		Podophyllin.....1-10 gr.	
		Ext. Hyoscyam...1-10 gr.	
		Ipecac Powd.....1-15 gr.	
		Aromatics.....q. s.	

One every hour until laxative action is produced; for a stimulant to the liver and secretions.

<b>LUPULIN &amp; BROMIDE CO....</b>	{	Lupulin.....1-2 gr.	} <b>2 00</b>
		Scutellarin.....1-2 gr.	
		Ergotin.....1-4 gr.	
		Atropine Sulph.1-2000 gr.	
		Zinc Bromide....1-10 gr.	

One or two; three or four times a day, as a general nerve sedative.

<b>MACROTIN COMP.....</b>	{	Macrotin.....1-10 gr.	} <b>1 50</b>
(Dr. John Stevens.)		Bryonia.....1-20 gr.	
		Strych. Arsen...1-125 gr.	
		Fl.Ex.Rhus Tox.1-20 min.	

One every two hours in muscular and sciatic rheumatism. One every hour in severe cases.

<b>MALARIA.....</b>	{	Quinine Sulph.....3 gr.	} <b>4 50</b>
(Dr. Clarke.)		Capsicum Powd....1 gr.	
		Opium Powd.....1-4 gr.	
		<b>Plain or Chocolate Coated</b>	

Two to four tablets at nine in the forenoon, one at three in the afternoon, and one at bedtime with plenty of drinking water; for remittent and malarial fevers.

<b>MALARIAL TONIC.....</b>	{	Quinine Bisulph....2 gr.	} <b>3 50</b>
(Dr. B. F. Hawley.)		Podophyllin.....1-24 gr.	
		Acid Arsenous...1-30 gr.	
		Oleores. Capsic...1-4 gr.	
		<b>Plain or Chocolate Coated</b>	

One tablet at eight, twelve, four and eight o'clock daily, with plenty of drinking water; used in chronic malaria with hepatic complication, and as a tonic in malarial districts.

<b>MANGANESE BINOXIDE</b> . . . 1-10 gr. . . . .	75
One tablet, crushed in a teaspoonful of sweetened water, three or four times a day, for an infant with anæmia.	
<b>MANGANESE BINOXIDE</b> . . . . .1 gr. . . . .	75
<b>Chocolate Coated</b> . . . . .	85
One after meals, for reconstructive tonic, where iron does not agree.	
<b>MANGANESE BINOXIDE</b> . . . . .2 grs. . . . .	1 25
<b>Chocolate Coated</b> . . . . .	1 35
One tablet every three hours, for a general tonic, and as a mild emmenagogue.	
<b>MANGANESE BINOXIDE</b> . . . . .3 grs. . . . .	1 60
One every three hours, beginning a few days before the period, for suppressed or scanty menstruation.	
<b>MENORRHAGIC</b> . . . . . { Acid Gallic. . . . .3 gr. }	
(Dr. Hirst.)	{ Ergotin. . . . .1 gr. }
	{ Hydrastin. . . . .1-2 gr. }
<b>Plain or Chocolate Coated</b>	5 60
One or two, every two or three hours as required.	
<b>MENTHOL COMP.</b> . . . . . { Menthol. . . . .1-20 gr. }	
(Dr. Ayres.)	{ Acid Carbolic. . . . .1-10 gr. }
One tablet every fifteen minutes until relieved, for nausea or vomiting in pregnancy, and as a gastric antiseptic.	
<b>MERCURIAL TONIC</b> . . . . . { Mercury Protiod. . . . .1-8 gr. }	
	{ Iron Citrate. . . . .1-2 gr. }
One after meals, as an hæmatic and alterative tonic for chronic diseases of the blood and skin.	
<b>MERCURIUS COMP.</b> . . . . . { Merc. Iod. Red. . . . .1-100 gr. }	
Throat troubles.	{ Belladonna Po. . . . .1-100 gr. }
	{ Kali Bichrom. . . . .1-100 gr. }
One every hour or two, for red and inflamed throat.	
<b>MERCURY, BELLADONNA CO</b> . . . . . { Merc. Iod. Red. . . . .1-200 gr. }	
Useful in throat troubles.	{ Ipecac Powd. . . . .1-20 gr. }
	{ Belladonna Po. . . . .1-100 gr. }
A tablet every hour or two, for sore throat with swollen glands.	
<b>MERCURY &amp; CHARCOAL</b> . . . . . { Mercury Protiod. . . . .1-4 gr. }	
(Dr. R. Tilley.)	{ Carbo Veg. . . . .1-10 gr. }
<b>Alterative.</b>	
One, half an hour after meals, for chronic skin and visceral disease. The Carbo Veg is added to change the color.	
<b>MERCURY &amp; CHARCOAL No. 2</b> { Mercury Protiod. . . . .1-6 gr. }	
	{ Carbo Veg. . . . .1-10 gr. }
One after meals, as a general tonic and alterative.	
<b>MERCURY &amp; HYOSCYAMUS</b> . . . . . { Mercury Protiod. . . . .1-4 gr. }	
	{ Ext. Hyoscyam. . . . .1-4 gr. }
One every three hours, as a general alterative.	
<b>MERCURY IODIDE COMP.</b> . . . . { Merc. Iod. Red. . . . .1-32 gr. }	
(Dr. Curtis.)	{ Ipecac Powd. . . . .1-8 gr. }
One or two after meals, for the continued treatment of syphilis.	
<b>MERCURY IODIDE CO. No. 2</b> . . { Merc. Iod. Red. . . . .1-16 gr. }	
(Dr. Curtis.)	{ Ipecac Powd. . . . .1-8 gr. }
One or two after meals, for the continued treatment of syphilis.	
<b>MERCURY IODIDE, RED</b> . . . . .1-1000 gr. . . . .	75
For an infant, one every hour, to control vomiting, diarrhœa and dysentery.	

When ordering or prescribing

<b>MERCURY IODIDE, RED.....</b>	<b>1-500 gr.....</b>	<b>75</b>
A tablet three or four times a day, as an intestinal antiseptic for infants.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-200 gr.....</b>	<b>75</b>
One every three or four hours, for croup and serous inflammations during infancy.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-100 gr.....</b>	<b>75</b>
One every two hours, for sore throat; one every two hours for biliousness and as an intestinal antiseptic.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-50 gr.....</b>	<b>75</b>
Give one three times a day, in congenital and acquired cachexia of children.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-32 gr.....</b>	<b>75</b>
One after meals, for lupus, scrofula and lymphatic enlargement and degeneration.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-25 gr.....</b>	<b>75</b>
	<b>Chocolate Coated.....</b>	<b>85</b>
A tablet every three or four hours, in phthisis and chronic bronchitis, and chronic skin diseases.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-20 gr.....</b>	<b>75</b>
One after meals, as an intestinal disinfectant and hepatic stimulant		
<b>MERCURY IODIDE, RED.....</b>	<b>1-10 gr.....</b>	<b>75</b>
	<b>Chocolate Coated.....</b>	<b>85</b>
One every three hours, for biliousness.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-12 gr.....</b>	<b>75</b>
One, three or four times daily, for chronic rheumatism and rheumatoid affections, and as a diuretic in dropsy.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-10 gr.....</b>	<b>75</b>
One after meals and at bedtime, for the treatment of syphilis.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-8 gr.....</b>	<b>75</b>
	<b>Chocolate Coated.....</b>	<b>85</b>
One after meals and at bedtime, for the treatment of syphilis.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-6 gr.....</b>	<b>90</b>
	<b>Chocolate Coated.....</b>	<b>1 00</b>
One every two or three hours, to produce rapid mercurialization, decreasing as the effect appears.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-5 gr.....</b>	<b>90</b>
One every two or three hours, for ascites, and to obtain the constitutional effect in the shortest time; decreasing the dose as soon as possible.		
<b>MERCURY IODIDE, RED.....</b>	<b>1-4 gr.....</b>	<b>90</b>
A tablet every three hours, to control the cerebral effects of syphilis, reducing the dose as the results are obtained.		
<b>MERCURY &amp; IPECAC.....</b>	( Mercury Protiod...1-3 gr.) ( Ipecac Powd.....1-8 gr.)	<b>1 10</b>
One, three or four times a day, for tonic syphilitic treatment.		
<b>MERCURY, IPECAC COMP....</b>	{ Mass. Hydrarg...1-2 gr.} { Ext. Hyoscyam...1-2 gr. { Ipecac Powd.....1-4 gr.}	<b>1 35</b>
One every three or four hours, for cirrhosis and functional disease of the liver.		
<b>MERCURY &amp; IRON.....</b>	( Mercury Protiod...1-4 gr.) ( Ferruginous alterative, } Iron Powd.....1 gr.)	<b>1 10</b>
A tablet with meals, as an alterative and hæmatic tonic, for chronic diseases and anæmia.		

<b>MERCURY &amp; OPIUM</b> .....	{ Mercury Protiod...1-5 gr. } { Opium Powd....1-12 gr. }	1 00
One three times a day, for tonic syphilitic treatment.		
<b>MERCURY &amp; OPIUM COMP.</b> ...	{ Mercury Protiod...1-4 gr. } { Opium Powd....1-12 gr. } { Charcoal.....1-10 gr. }	1 00
One, three times a day, for tonic syphilitic treatment.		
<b>MERCURY &amp; POTASS.IODIDE</b> }	{ Merc. Iod. Red....1 gr. } { Potass. Iodide.....1 gr. }	3 00
(For antiseptic use.)		
One tablet dissolved in four tablespoonfuls of water makes a solution of about 1 to 1000.		
<b>MERCURY PROTIODIDE</b> .....	1-1000 gr.....	75
One every hour, for the vomiting and diarrhoea of infantile gastro-enteritis.		
<b>MERCURY PROTIODIDE</b> .....	1-500 gr.....	75
Give one tablet every three hours, to an infant, to promote intestinal antiseptis.		
<b>MERCURY PROTIODIDE</b> .....	1-200 gr.....	75
One every two hours, to an infant, for hepatic stimulant and aid to digestion.		
<b>MERCURY PROTIODIDE</b> .....	1-100 gr.....	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
<b>MERCURY PROTIODIDE</b> .....	1-50 gr.....	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
<b>MERCURY PROTIODIDE</b> .....	1-25 gr.....	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
<b>MERCURY PROTIODIDE</b> .....	1-20 gr.....	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
<b>MERCURY PROTIODIDE</b> .....	1-16 gr.....	75
One every two hours, to stimulate secretions and as an intestinal antiseptic.		
<b>MERCURY PROTIODIDE</b> .....	1-10 gr.....	75
	<b>Chocolate Coated</b> .....	85
One every three hours, as an hepatic stimulant; also used in continued constitutional treatment.		
<b>MERCURY PROTIODIDE</b> .....	1-8 gr.....	75
	<b>Chocolate Coated</b> .....	85
One or two every three hours, in the tonic treatment of syphilis.		
<b>MERCURY PROTIODIDE</b> .....	1-6 gr.....	75
	<b>Chocolate Coated</b> .....	85
One or two every three hours, in the tonic treatment of syphilis.		
<b>MERCURY PROTIODIDE</b> .....	1-5 gr.....	75
	<b>Chocolate Coated</b> .....	85
One or two every three hours, in the tonic treatment of syphilis.		
<b>MERCURY PROTIODIDE</b> .....	1-4 gr.....	75
	<b>Chocolate Coated</b> .....	85
One every three hours, to produce rapid mercurialization; decrease the dose as soon as possible.		

When ordering or prescribing

<b>MERCURY PROTIODIDE</b> .....1-3 gr.....	95
<b>Chocolate Coated</b> .....	1 05
One every three hours, to produce rapid mercurialization; decrease the dose as soon as possible.	
<b>MERCURY PROTIODIDE</b> .....1-2 gr.....	1 20
<b>Chocolate Coated</b> .....	1 30
A tablet every three hours, with large doses of potass. iodide, to control cerebral symptoms in syphilis; decrease the mercury as symptoms improve.	
<b>MERCURY PROTIODIDE</b> .....1 gr.....	1 85
<b>Chocolate Coated</b> .....	1 95
A tablet every three hours, with large doses of potass. iodide, to control cerebral symptoms in syphilis; decrease the mercury as symptoms improve.	
<b>MERCURY SALICYLATE</b> .....1-4 gr.....	1 40
One every hour, for a rapid effect without severe irritation; lessen the dose as the result is obtained.	
<b>MERCURY SALICYLATE</b> .....1-3 gr.....	1 60
One every two hours, for a rapid effect without severe irritation; lessen the dose as the result is obtained.	
<b>MERCURY SPECIAL</b> .....	1 75
(Dr. Shimp.)	
{ Mercury Protiod...1-4 gr.	
{ Opium Powd.....1-6 gr.	
{ Ext. Aconite.....1-4 gr.	1 75
{ Ext. Conium.....1-3 gr.	
One tablet three times a day in the continued tonic treatment of syphilis.	
<b>MERCURY, STRYCHNINE CO.</b> { Merc. with Chalk...2 gr. }	1 40
(Dr. R. W. Stewart.) { Iron Sulph. Dry...1-2 gr. }	
{ Strych. Sulph....1-100 gr. }	
One to two tablets after meals where a tonic is indicated, with sluggish condition of the liver, and as an alternative in diseases of the blood.	
<b>MERCURY TANNATE</b> .....1-2 gr.....	1 50
One every two hours, for a rapid effect without severe irritation; lessen the dose as the result is obtained.	
<b>MERCURY TANNATE</b> .....1 gr.....	2 00
<b>Chocolate Coated only</b>	
One three times a day; seldom produces salivation, stomatitis, or diarrhœa. Contains about 40 per cent. mercury, and is very rapidly absorbed.	
<b>MERCURY WITH CHALK</b> .....1-100 gr.....	75
One, in a little water on the end of a spoon, repeated every hour for infantile colic with profuse diarrhœa.	
<b>MERCURY WITH CHALK</b> .....1-40 gr.....	75
One every three hours to a child, for colic with profuse diarrhœa.	
<b>MERCURY WITH CHALK</b> .....1-20 gr.....	75
One every hour for an adult, for hepatic corrective, until ten or fifteen have been taken.	
<b>MERCURY WITH CHALK</b> .....1-10 gr.....	75
One every hour, for hepatic corrective, until effect is produced.	
<b>MERCURY WITH CHALK</b> .....1-3 gr.....	75
One every two hours, for tonic to liver, and corrective in bilious diarrhœa until effect is produced; as an alternative, one three times a day.	

<b>MERCURY WITH CHALK</b> .....1-2 gr.....	75
One every two hours, for tonic to liver, and corrective in bilious diarrhoea; as an alterative, one three times a day.	
<b>MERCURY WITH CHALK</b> .....1 gr.....	75
One tablet repeated every three hours until effect is produced, as hepatic alterative and laxative.	
<b>MERCURY YELLOW OXIDE</b> ..1-1000 gr.....	75
One every hour, for a child with indigestion and diarrhoea, to control septic conditions of the intestine.	
<b>MERCURY YELLOW OXIDE</b> ..1-200 gr.....	75
One tablet every two or three hours, for dysentery; one after meals as an alterative in chronic disease.	
<b>MERCURY YELLOW OXIDE</b> ..1-100 gr.....	75
A tablet after meals, for chronic diseases of the skin and lymphatics; one every hour to secure immediate constitutional effects.	
<b>MIGRAINE</b> .....	{ Acetanilid.....2 gr. } { Camph. Monobr...1-2 gr. } { Caffeine Citrate...1-2 gr. } <b>Plain or Chocolate Coated</b> 2 00
Two tablets, repeated every hour, for four or five doses, for neuralgic and nervous headache.	
<b>MIGRAINE No. 2</b> .....	{ Acetanilid.....3 gr. } { Camph. Monobrom.2 gr. } { Caffeine Citrate....1 gr. } 3 00
One, repeated every hour, for three doses, then every two hours; as a relief to migraine, surpraorbital and frontal.	
<b>MIGRAINE No. 3</b> .....	{ Acetanilid.....3 1-2 gr. } { Caffeine.....1-2 gr. } 2 00 { Sodium Bicarb.....1 gr. }
One, repeated in an hour, for sick headache, and pains in the head with fever; may be continued for three doses, then stop and begin again in three hours if not relieved.	
<b>MIGRAINE No. 4</b> .....	{ Acetanilid.....1 2-5 gr. } { Caffeine Alk.....1-5 gr. } 1 40 { Sodium Bicarb...2-5 gr. }
From one to three tablets, according to age, repeated every hour for four to five doses, for neuralgic and nervous headache.	
<b>MIGRAINE No. 5</b> .....	{ Acetanilid.....3 gr. } { Caffeine Citrate....1 gr. } { Camphor Monobr...1 gr. } 3 00 { Sodium Bicarb....1 gr. } (Dr. Boggus.)
One tablet, repeated every hour for three or four doses, for neuralgic and nervous headache.	
<b>MIGRAINE No. 6</b> .....	{ Acetanilid.....2 1-2 gr. } { Sodium Bicarb.....1 gr. } { Camphor Monobr..1-2 gr. } 1 85 { Caffeine Alk.....1-4 gr. } { Capsicum Powd..1-10 gr. } <b>Chocolate Coated</b> ..... 1 95
Two tablets for first dose; repeat one every hour with a swallow of water, for nervous, neuralgic, and sick headache.	
<b>MIGRAINE (Ammoniated) No. 1</b> .....	{ Acetanilid.....1 3-4 gr. } { Caffeine Alkaloid..8 gr. } { Sodium Bicarb....1-2 gr. } 1 25 { Camphor Monobr..1-4 gr. } { Ammon. Muriate..1-2 gr. } <b>Chocolate Coated</b> ..... 1 35
Two tablets every hour for four or five doses; for neuralgic and nervous headache, or pains in the head with fever. If not relieved, stop for three hours then begin again.	

When ordering or prescribing

MIGRAINE (Ammoniated) No. 2	{ Acetanilid.....3 gr. }	1 90
	{ Caffeine Alkaloid...1-4 gr. }	
	{ Sodium Bicarb....1-2 gr. }	
	{ Camphor Monobr...1-2 gr. }	
	{ Ammon. Muriate...1 gr. }	
	Chocolate Coated	2 00

One tablet every hour for four or five doses; for neuralgic and nervous headache, or pains in the head with fever. If not relieved, stop for three hours then begin again.

MIGRAINE IMPROVED.....	{ Acetanilid.....2 1-2 gr. }	2 00
	{ Camphor Monobr...1-2 gr. }	
	{ Sodium Salicylate...1 gr. }	
	{ Ext. Hyoscyam...1-8 gr. }	
	{ Tr. Gelsemium....2 min. }	

One to three tablets; repeat in half hour if necessary; then one tablet every hour for migraine, neuralgia, and rheumatic pains.

MIGRAINE No. 3, Modified.....	{ Acetanilid.....3 1-2 gr. }	2 75
	{ Caffeine Alk.....1-2 gr. }	
	{ Sodium Bicarb.....1 gr. }	
	{ Tully's Powder.....3 gr. }	

One every two to four hours, for the fever of pneumonia, bronchitis and typhoid; and as an analgesic in rheumatism.

MILK.....	{ Sugar of Milk.....26 gr. }	2 25
	{ Calc. Lactophos...1-6 gr. }	
	{ Calc. Carbonate...1 1-2 gr. }	
	{ Sodium Bicarb....1-2 gr. }	
	{ Potass. Bicarb....1-12 gr. }	
	{ Sodium Chloride...1-6 gr. }	

Four tablets to a bottle of milk, as a reconstructive tonic for bottle-fed babies.

MIST. GUNTHER COMP. One-half Size..... (Dr. C. Eugene Gunther.)	{ Tr. Capsicum...1 1-4 min. }	2 50
	{ Magnesia Sulph....5 gr. }	
	{ Quin. Sulph.....5-8 gr. }	
	{ F. E. Buckthorn...5 min. }	

Two tablets four times daily, or more if necessary, for paludal cachexia.

MIXED TREATMENT..... (Dr. Vidal.) Alterative and antisyphilitic.	{ Potass. Iodide.....5 gr. }	5 65
	{ Corrosive Sub....1-30 gr. }	
	{ Syr. Sars. Co....30 min. }	

Begin with three a day, and increase according to the necessities of the case.

MORPHINE & ACONITE.....	{ Morph. Sulph....1-32 gr. }	1 10
	{ Ext. Aconite....1-100 gr. }	
MORPHINE & ACONITE No. 2	{ Morph. Sulph....1-8 gr. }	2 00
	{ Ext. Aconite....1-25 gr. }	
MORPHINE & BELLADONNA	{ Morph. Sulph....1-6 gr. }	2 40
	{ Ext. Belladonna...1-8 gr. }	
MORPHINE COMP No. 1.....	{ Morph. Sulph....1-8 gr. }	2 00
	{ Atropine Sulph...1-150 gr. }	
MORPHINE COMP. No. 1..... Colored Pink.	{ Morph. Sulph....1-8 gr. }	2 00
	{ Atropine Sulph...1-150 gr. }	
MORPHINE COMP. No. 2.....	{ Morph. Sulph....1-4 gr. }	3 35
	{ Atropine Sulph...1-120 gr. }	
MORPHINE COMP. No. 2..... Colored Pink.	{ Morph. Sulph....1-4 gr. }	3 35
	{ Atropine Sulph...1-120 gr. }	
MORPHINE COMP. No. 3.....	{ Morph. Sulph....1-8 gr. }	2 00
	{ Atropine Sulph...1-200 gr. }	

Please specify Fraser's Tablets

MORPHINE COMP. No. 3.....	{ Morph. Sulph.....1-8 gr. }	2 00
Colored Pink,	{ Atropine Sulph.1-200 gr. }	
MORPHINE COMP. No. 4.....	{ Morph. Sulph.....1-4 gr. }	3 35
	{ Atropine Sulph.1-150 gr. }	
MORPHINE COMP. No. 4.....	{ Morph. Sulph.....1-4 gr. }	3 35
Colored Pink,	{ Atropine Sulph.1-150 gr. }	
MORPHINE, ALOIN COMP.....	{ Morphine Sulph.. 1-6 gr. }	2 50
Non-Constipating.	{ Aloin.....1-6 gr. }	
(Dr. Langstaff.)	{ Atropine Sulph. 1-200 gr. }	

One at a dose as directed.

MORPHINE SULPHATE.....	1-100 gr.....	75
MORPHINE SULPHATE.....	1-50 gr.....	95
MORPHINE SULPHATE.....	1-20 gr.....	1 00
MORPHINE SULPHATE.....	1-16 gr.....	1 30
MORPHINE SULPHATE.....	1-10 gr.....	1 50
MORPHINE SULPHATE.....	1-10 gr.....Colored Pink	1 50
MORPHINE SULPHATE.....	1-8 gr.....	1 85
MORPHINE SULPHATE.....	1-8 gr.....Colored Pink	1 85
MORPHINE SULPHATE.....	1-6 gr.....	2 25
MORPHINE SULPHATE.....	1-6 gr.....Colored Pink	2 25
MORPHINE SULPHATE.....	1-4 gr.....	3 20
MORPHINE SULPHATE.....	1-4 gr.....Colored Pink	3 20
MORPHINE SULPHATE.....	1-3 gr.....	3 75
MORPHINE SULPHATE.....	1-3 gr.....Colored Pink	3 75
MORPHINE SULPHATE.....	1-2 gr.....	5 60
MORPHINE SULPHATE.....	1-2 gr.....Colored Pink	5 60
MUSCARIN SULPHATE.....	1-100 gr.....	7 00

A tablet every three or four hours stimulates the cutaneous, salivary and intestinal glands, and the intra-cardiac inhibitory centres; useful in scarlatina, rheumatism and respiratory diseases.

NAPHTALIN, Pure.....	5 grs.....	2 00
Antiseptic.		

A tablet three to six times a day for the diarrhoea and elevated temperature in typhoid; one to three as an anthelmintic, followed by a dose of castor oil.

NAUSEA No. 2.....	{ Bismuth Subnit.....2 gr. }	4 75
	{ Cerium Oxalate....2 gr. }	
	{ Cocaine Hydro.. 1-12 gr. }	

One every hour until stomach is settled, or ten doses are taken. Useful for car sickness or sea sickness.

NAUSEA No. 3.....	{ Cerium Oxalate.....5 gr. }	2 25
(Dr. Guernsey.)	{ Ipecac Powd.....1-10 gr. }	

One every three or four hours, for vomiting of pregnancy.

NAUSEA No. 4.....	{ Bismuth Subnit.....1 gr. }	
	{ Cerium Oxalate....1 gr. }	
	{ Menthol.....1-10 gr. }	
	{ Acid Carbol.....1-10 gr. }	
	{ Cocaine Mur.... 1-40 gr. }	
	{ Tr. Nux Vom.... 1-2 min. }	

Plain or Chocolate Coated 2 00

Two, and repeat one every hour for three or four doses until relieved. For vomiting in pregnancy, sick stomach and seasickness.

When ordering or prescribing

**NAUSEA No. 5**..... { Bismuth Subnit.....5 gr. }  
 (Dr. Bierwirth.) { Cerium Oxalate.....5 gr. } 3 75  
 One or two every half hour to one hour until relieved.

**NERVINE**..... { Camphor.....1 gr. }  
 { Ext. Hyoscyam...1-2 gr. }  
 { Valerian Powd....1-2 gr. }  
**Chocolate Coated only** 1 50

One every half hour for three doses, then every two hours until nervousness is relieved.

**NERVINE No. 2**..... { Lupulin.....1 gr. }  
 (Dr. Lee.) { Zinc Phosphide...1-16 gr. }  
 { Sodium Bromide...2 1-2 gr. } 2 00  
 { Gelsemin .....1-20 gr. }  
 { Acid Arsenous...1-20 gr. }

Two tablets, three times a day, for irritability of the nervous system and neurasthenia.

**NERVINE No. 3**..... { Fl. Ext. Coca.....10 min. }  
 (Dr. E. M. Lyon.) { Tr. Hyoscyamus...10 min. } 3 00  
 { Elix. Val. Ammon...10 min. }

Two tablets in hot water, repeated in one hour, for sleeplessness, hysteria and nervous disorders.

**NERVINE No. 4**..... { Camphor Powd....1 gr. }  
 { Ext. Hyoscyam...1-2 gr. }  
 { Ext. Valerian.....1-2 gr. }  
**Chocolate Coated only** 1 60

One every half hour for three doses, then every two hours until nervousness is relieved.

**NERVINE, TOBACCO**..... { Tr. Strophant...1-2 min. }  
 (Dr. F. H. Koyle.) { Salicin.....2-3 gr. }  
 { Cinchonid. Sulph...1 gr. } 2 50  
 { Ext. Nux Vom....1-5 gr. }  
 { Hydrastis Powd...1-4 gr. }  
 { Fl. Ext. Sanguin...1-4 min. }

One three times daily immediately after eating, for general tonic and cure of tobacco heart.

**NERVE TONIC No. 1**..... { Zinc Phosphide...1-10 gr. }  
 (Dr. Westbrook.) { Ext. Nux Vom....1-4 gr. } 1 40  
 { Iron Powd.....1 gr. }  
**Chocolate Coated**..... 1 50

One, three times a day, may be given as long as improvement continues or until stomach is nauseated; if nausea ensues, may be discontinued for one week, then repeated.

**NERVE TONIC No. 2**..... { Zinc Phosphide...1-10 gr. }  
 (Dr. Westbrook.) { Ext. Nux Vom....1-4 gr. } 1 30  
 { Acid Arsenous...1-20 gr. }  
**Chocolate Coated**..... 1 40

One, three times a day, may be given as long as improvement continues. If nausea or redness of the eyelids, or pains or fulness of the stomach occur, discontinue the tablet for a week, then continue.

**NERVE TONIC No. 4**..... { Zinc Phosphide...1-10 gr. }  
 (Dr. Westbrook.) { Ext. Nux Vom....1-4 gr. } 1 75  
 { Iron Powd.....2 gr. }  
 { Acid Arsenous...1-20 gr. }  
**Chocolate Coated**..... 1 85

One, three times a day, may be given as long as improvement continues. If nausea or redness of the eyelids, or pains or fulness of the stomach occur, discontinue the tablet for a week, then continue.

NERVE TONIC No. 5.....	{ Zinc Phosphide...1-8 gr. }	1 30
	{ Ext. Nux Vom....1-8 gr. }	
	{ Ext. Can. Indica...1-8 gr. }	
	<b>Chocolate Coated.....</b>	<b>1 40</b>

One with meals, for nervous affections with exalted reflexes and excitability.

NERVE TONIC No. 6.....	{ Ext. Can. Indica...1-8 gr. }	2 25
	{ Hyoscyamia....1-400 gr. }	
	{ Zinc Phosphide...1-10 gr. }	

Three or four tablets a day, for chorea, neurasthenia and hysteria.

NERVE TONIC No. 7.....	{ Hyoscyamia....1-400 gr. }	2 50
	{ Zinc Phos.....1-10 gr. }	
	{ Strych. Nitrate...1-20 gr. }	

One, three times a day, for disorders of the nerves dependent upon anæmia and innutrition of the cord.

NERVE TONIC No. 8..... (Dr. Henry J. Kenyon.)	{ Zinc Phosphide...1-40 gr. }	2 00
	{ Ext. Can. Indica...1-8 gr. }	
	{ Ext. Nux Vomica...1-8 gr. }	
	{ Sodium Arsen....1-64 gr. }	
	{ Quinine Sulph....1-2 gr. }	
	{ Ext. Aconite Rad...1-10 gr. }	
	{ Sugar of Milk.....q. s. }	
	<b>Plain or Chocolate Coated.....</b>	<b>2 00</b>

One, at 10 a.m. 4 and 9 p.m. Increase by one tablet a week, until three are taken, three times daily, for chronic neuralgic headache, malarial neuralgia, and as a general nerve tonic.

NEURALGIA..... (Dr. Gross.)	{ Quinine Sulph.....2 gr. }	4 50
	{ Acid Arsenous...1-20 gr. }	
	{ Strychnine Sulph...1-30 gr. }	
	{ Morphine Sulph...1-20 gr. }	
	{ Ext. Aconite Leaf...1-2 gr. }	
	<b>Chocolate Coated.....</b>	<b>4 50</b>

NEURALGIA without Morphine. (Dr. Gross.)	{ Quinine Sulph.....2 gr. }	4 00
	{ Acid Arsenous...1-20 gr. }	
	{ Strychnine Sulph...1-30 gr. }	
	{ Ext. Aconite Leaf...1-2 gr. }	
	<b>Chocolate Coated.....</b>	<b>4 00</b>

NEURALGIA..... (Brown-Sequard.)	{ Ext. Hyoscyamus...2-3 gr. }	4 50
	{ Ext. Conium.....2-3 gr. }	
	{ Ext. Ignatia.....1-2 gr. }	
	{ Ext. Opium.....1-2 gr. }	
	{ Ext. Aconite.....1-3 gr. }	
	{ Ext. Cannab. Ind...1-4 gr. }	
	{ Ext. Stramonium...1-5 gr. }	
	{ Ext. Belladonna...1-6 gr. }	
	<b>Plain or Chocolate Coated.....</b>	<b>4 50</b>

NEURALGIA, One-half strength (Brown-Sequard.)	{ Ext. Hyoscyamus...1-3 gr. }	2 75
	{ Ext. Conium.....1-3 gr. }	
	{ Ext. Ignatia.....1-4 gr. }	
	{ Ext. Opium.....1-4 gr. }	
	{ Ext. Aconite.....1-6 gr. }	
	{ Ext. Cannab. Ind...1-8 gr. }	
	{ Ext. Stramonium...1-10 gr. }	
	{ Ext. Belladonna...1-12 gr. }	
	<b>Plain or Chocolate Coated.....</b>	<b>2 75</b>

When ordering or prescribing

NEURALGIC..... (Dr. Dunlap.)	Quinine Sulph.....1-2 gr.	} 1 75
	Morph. Sulph.....1-80 gr.	
	Strych. Sulph....1-120 gr.	
	Acid Arsenous...1-80 gr.	
	Ext. Aconite.....1-8 gr.	
	Chocolate Coated.....	1 85

One every hour until eight are taken, if necessary. For neuralgia and neuralgic headache.

NEURALGIC No. 2..... (Dr. Henry J. Kenyon.)	Zinc Phosphide...1-16 gr.	} 1 50
	Strych. Sulph....1-60 gr.	
	Ext. Can. Indica...1-8 gr.	
	Sodium Arsen...1-20 gr.	
	Aconitia.....1-400 gr.	
	Chocolate Coated.....	1 60

NEURALGIC No. 4..... (Dr. Gustavus Elliot.)	Fl. Ext. Bellad...1-6 min.	} 1 50
	Fl. Ext. Aconite...1-2 min.	
	Fl. Ext. Gelsem...1-3 min.	

One every hour or every second hour, for painful affections of the nerves, not due to an organic lesion.

NEURALGIC No. 5.....	Aconitia Amor...1-400 gr.	} 2 50
	Acetanilid.....2 gr.	
	Quinine Muriate...1 gr.	
	Arsenic Chloro...1-100 gr.	
	Strych. Muriate...1-120 gr.	

One every two hours, for three doses; then one every three or four hours until relieved.

NEURALGIC HEADACHE..... (Dr. Myers.)	Sodium Bromide....5 gr.	} 2 10
	Caffeine Citrate...1-2 gr.	
	Acetanilid.....1 gr.	
	Ext. Hyoscyam...1-2 gr.	
	Morph. Sulph....1-50 gr.	

A tablet, repeated in half an hour, for migraine, headache and facial neuralgia.

NEURALGIC HEADACHE No. 2	Sodium Bromide....5 gr.	} 3 00
	Caffeine Citrate...1-2 gr.	
	Acetanilid.....1 gr.	
	Ext. Hyoscyam...1-2 gr.	
	Morphine Sulph...1-50 gr.	
	Ext. Can. Indica...1-4 gr.	

A tablet, repeated in half hour, for migraine, headache and facial neuralgia.

NEURALGIC, IMPROVED.....	Quinine Sulph.....2 gr.	} 4 00
	Acetanilid.....2 gr.	
	Ext. Hyoscyamus...1-2 gr.	
	Ext. Can. Ind....1-4 gr.	
	Acid Arsenous...1-100 gr.	
	Strych. Sulph....1-60 gr.	

One tablet; repeat in half hour if necessary. Then one tablet every one or two hours for neuralgia and migraine.

NEURASTHENIA..... (Dr. Krauss.)	Lithium Carb.....3 gr.	} 3 25
	Iron Phosphate...1-3 gr.	
	Ext. Nux Vomica...1-0 gr.	
	Sodium Arsen...1-27 gr.	

One to three tablets after meals, for neurasthenia and lithæmia with predominant nervous symptoms.

<b>NEURASTHENIA No. 2.....</b>		<div> <div> Quin. Valerianate...1 gr.  Strych. Valerian...1-50 gr.  Sumbul Ext.....1 gr. </div> <div> <b>Chocolate Coated only</b> </div> </div>	<b>3 50</b>
One three times daily for neurasthenia and as a general nerve tonic.			
<b>NEUROTIC.....</b> (Dr. North.)		<div> <div> Tr. Benzoin Co....2 min.  Tr. Lavender Co....4 min.  Tr. Valer. Amm...4 min. </div> </div>	<b>2 25</b>
Give one tablet every ten minutes until relieved, for hysteria and nervousness of women.			
<b>NEUTRALIZING .....</b>		<div> <div> Rhubarb Powd. ....1 gr.  Sodium Bicarb.....3 gr.  Magnesium Calc...1 gr.  Pulv. Cardamom...1-8 gr.  Oil Peppermint...1-8 min. </div> </div>	<b>1 50</b>
One or two tablets half an hour after eating, for gastralgia and flatulence, and sour eructations.			
<b>NIGHT SWEAT .....</b>		<div> <div> Zinc Oxide.....1-2 gr.  Salicin .....1 gr.  Hydrastin.....1-10 gr.  Pepsin Lactated...1-2 gr.  Ext. Belladonna.1-25 gr. </div> </div>	<b>2 50</b>
One an hour before retiring, for night sweats.			
<b>NIGHT SWEAT No. 2.....</b>		<div> <div> Agaricin.....1-12 gr.  Picrotoxin.....1-60 gr.  Atropine Sulph.1-120 gr. </div> </div>	<b>1 75</b>
One at bedtime, repeated in two hours if necessary, for night-sweats.			
<b>NITROGLYCERIN .....</b>		<b>1-200 gr.....</b>	<b>75</b>
Heart tonic and antiasthmatic.			
One when face is pale, for fainting, neuralgia of heart, or heart failure. Never give if face is flushed.			
<b>NITROGLYCERIN.....</b>		<b>1-150 gr.....</b>	<b>75</b>
A tablet, repeated in an hour if necessary, for pallor, headache and syncope; one three times a day for epilepsy, catalepsy and angina-pectoris.			
<b>NITROGLYCERIN.....</b>		<b>1-100 gr.....</b>	<b>75</b>
One when face is pale, for fainting, neuralgia of heart, or heart failure. Never give if face is flushed.			
<b>NITROGLYCERIN.....</b>		<b>1-50 gr.....</b>	<b>75</b>
One tablet, repeated in two or three hours if necessary, for cardiac syncope, epileptiform seizure, extreme asthmatic paroxysms, angina pectoris, and parturient eclampsia.			
<b>NITROGLYCERIN CO. No. 1...</b>		<div> <div> Nitroglycerin...1-100 gr.  Tr. Digitalis...2 min. </div> </div>	<b>1 25</b>
One, three times a day for feeble heart and cerebral anæmia; for epilepsy, increase by one tablet every three weeks.			
<b>NITROGLYCERIN CO. No. 2...</b>		<div> <div> Nitroglycerin...1-100 gr.  Tr. Strophanthus...2 min. </div> </div>	<b>1 50</b>
A tablet three times a day; for nephritis with pulse of great tension, cardiac weakness, and the anæmic form of migraine.			
<b>NUCLEIN SOLUTION.....</b>		<b>48-150 min .....</b>	<b>5 00</b>
(Dr. Aulde.)		<b>Per 500.....</b>	<b>2 50</b>
		<b>Per 100 .....</b>	<b>60</b>
One every two to four hours, or two tablets three or four times a day, in all diseased conditions characterized by disorganized state of the blood. From 1-4 to 1-2 a tablet for a child five years old.			

When ordering or prescribing



NUX VOMICA COMP. No. 2....	{ Ext. Nux Vom...1-16 gr. }	75
	{ Acid Arsenous...1-80 gr. }	
	{ Pepsin Pure.....1-32 gr. }	
	{ Aromatics .....q. s. }	

One, three or four times a day, for anorexia, dyspepsia and constipation.

NUX VOMICA COMP. No. 3....	{ Ext. Nux Vom...1-20 gr. }	75
	{ Sulphur.....1-20 gr. }	
	{ Acid Arsenous...1-100 gr. }	

One or two after meals, for chronic skin diseases dependent upon dyspepsia.

NUX & ARSENIC.....	{ Nux Vom. Po....1-10 gr. }	75
	{ Acid Arsenous...1-50 gr. }	
	{ Chocolate Coated..... }	

One after meals, as a tonic in malarial indigestion.

NUX & BRYONIA.....	{ Nux Vom. Po...1-100 gr. }	75
	{ Bryonia Powd...1-100 gr. }	

For a child, give one every two hours, as a tonic to the stomach and an aid to digestion, with slight laxative action.

NUX & BRYONIA No. 2.....	{ Tr. Nux Vom.....1 min. }	1 10
	{ Tr. Bryonia.....1 min. }	

One every two or three hours, as a tonic to the stomach and aid to digestion.

NUX & CANTHARIDES..	{ Nux Vom. Po...1-100 gr. }	75
	{ Cantharides...1-100 gr. }	

A tablet three times a day, for impotence and spermatorrhœa; best combined with iron.

NUX & CARBO VEG.....	{ Nux Vom. Po...1-100 gr. }	75
	{ Carbo Veg.....1-10 gr. }	

One after meals, for atonic gastric disorders in children, and those due to fermentation.

NUX & CARBO VEG. No. 2....	{ Nux Vom. Po....1-2 gr. }	75
	{ Carbo Veg.....1-4 gr. }	

A tablet three times a day, for chronic catarrhal inflammation of the stomach, with flatulence.

NUX & IGNATIA COMP.....	{ Tr. Nux Vomica...1 min. }	1 75
	{ Tr. Ignatia.....1 min. }	
	{ Tr. Cinchona Cal...5 min. }	
	{ Tr. Matricaria....1 min. }	
	{ Tr. Gentian Comp.1 min. }	
	{ Tr. Columbo.....1-2 min. }	
	{ Phosphorus.....1-300 gr. }	
	{ Aromatic Powd...1 gr. }	
	{ Chocolate Coated..... }	1 85

One to two three times a day before meals, for atonic dyspepsia.

NUX & IPECAC COMP.....	{ Tr. Nux Vomica...5 min. }	2 00
	{ Tr. Capsicum.....2 min. }	
	{ Wine Ipecac.....1 min. }	
	{ Quinine Sulph....1-2 gr. }	

(Dr. F. R. Newman.)

One tablet after each meal, for a stomachic tonic and an aid to digestion.

NUX & IPECAC CO. No. 2.....	{ Sodium Bicarb.....5 gr. }	1 25
	{ Tr. Nux Vom.....2 min. }	
	{ Ipecac Powd.....1-20 gr. }	
	{ Oleoresin Ginger.1-6 min. }	
	{ Chocolate Coated..... }	1 35

One or two, half an hour before meals, stimulates appetite and promotes free flow of gastric juice.

NUX & PEPSIN.....	{ Nux Vom. Po....1-100 gr. }	75
(Dr. Skiff.)	{ Pepsin Pure.....1-10 gr. }	
Tonic, antidyspeptic.		
Chocolate Coated.....		85
Two tablets after each meal, for simple indigestion. For deficient gastric secretion producing fermentation and eructation, one tablet every two hours.		
NUX & PEPSIN No. 2.....	{ Nux Vomica.....1-10 gr. }	1 50
	{ Pepsin Pure.....1-2 gr. }	
Two after each meal, followed by one every half hour for three doses, if necessary, as an aid to digestion and a tonic to the stomach.		
NUX & PEPSIN COMP.....	{ Nux Vom. Po....1-100 gr. }	1 75
	{ Pepsin Pure.....1-4 gr. }	
	{ Bismuth Subnit.....1 gr. }	
	{ Calomel.....1-4 gr. }	
Chocolate Coated.....		1 85
One or two after meals, for chronic gastritis, nausea and pain after eating.		
NUX & PHOSPHORUS.....	{ Ext. Nux Vom....1-4 gr. }	1 00
	{ Phosphorus.....1-50 gr. }	
Chocolate Coated.....		1 10
One, three or four times a day, for headache and neuralgia depending upon intestinal dyspepsia.		
NUX & SULPHUR.....	{ Nux Vom. Po....1-100 gr. }	75
	{ Sulphur.....1-100 gr. }	
Two or three after a meal, as an aid to digestion.		
NUX & SULPHUR COMP.....	{ Nux Vom. Po....1-20 gr. }	75
	{ Sulphur.....1-10 gr. }	
	{ Acid Arsenous..1-100 gr. }	
Chocolate Coated.....		85
One after meals, for chronic skin diseases, anæmia, and indigestion.		
OIL SANDALWOOD.....	2 drops.....	4 00
A tablet every two to four hours, for inflammations of the genito-urinary tract.		
OPIUM POWD.....	1-100 gr.....	75
OPIUM POWD.....	1-50 gr.....	75
OPIUM POWD.....	1-20 gr.....	75
OPIUM POWD.....	1-10 gr.....	75
OPIUM POWD.....	1-8 gr.....	75
OPIUM POWD.....	1-4 gr.....	1 00
OPIUM POWD.....	1-2 gr.....	1 40
OPIUM AROMATIC No. 1.....	{ Pulv. Cret. Aromat. cum }	1 10
	{ Opio.....1 gr. }	
A tablet every two hours, to three doses, for diarrhœa and enteritis.		
OPIUM AROMATIC No. 2.....	{ Opium Powd....1-20 gr. }	1 25
(Dr. Davis.)	{ Acid Tannic.....1-20 gr. }	
	{ Pulv. Cret. Aromat. cum }	
	{ Opio.....1 gr. }	
Give one every two or three hours, for intestinal inflammation, discharge and pain.		
OPIUM & CAMPHOR.....	{ Opium Powd....1-4 gr. }	1 10
Sedative.	{ Camphor.....1-4 gr. }	
Chocolate Coated.....		1 20
Three to six a day for painful cough, restlessness, cramps and spasmodic attacks.		

<b>OPIUM &amp; CAMPHOR No. 2....</b>	{ Opium Powd.....1 gr. Camphor Powd.....2 gr. <b>Chocolate Coated only</b>	<b>3 00</b>
One tablet as a dose for restlessness, cramps and spasmodic attacks.		
<b>OPIUM &amp; COPPER SULPH ...</b>	{ Opium Powd.....1-4 gr. Copper Sulph.....1-12 gr. Ipecac Powd.....1-12 gr.}	<b>1 25</b>
One every two or three hours, for diarrhœa.		
<b>OPIUM DENARCOTIZED.....</b>	1-4 gr.....	<b>1 40</b>
<b>OPIUM &amp; LEAD .....</b>	{ Opium Powd.....1-16 gr. Lead Acetate.....1-8 gr.}	<b>1 00</b>
A tablet every two hours, for the summer diarrhœa and dysentery of children.		
<b>OPIUM &amp; LEAD No. 2.....</b>	{ Opium Powd.....1-8 gr. Lead Acetate.....1-4 gr.}	<b>1 00</b>
One every hour for twelve doses, for obstinate diarrhœa and dysentery.		
<b>ORTHODOX.....</b> Dr. Sherwell.)	{ Potass. Iodide.....2 gr. Syr. Ferri Iod.....5 min. Mercury Bichlor.1-64 gr. Donovan's Sol.....2 min. Tr. Nux Vom.....2 min.}	<b>3 50</b>
One to three, three times a day, followed by half a tumbler full of water.		
<b>OXGALL CO. No. 2.. .....</b>	{ Ovgall-Purified.....1 gr. Ext. Pancreatin...1 gr. Ext.Colocynth.Co.1-4 gr. Quinine Muriate..1-2 gr. Ext. Nux Vomica.1-8 gr. Ext. Taraxacum....1 gr. <b>Chocolate Coated only</b>	<b>2 50</b>
One or two before meals to promote intestinal digestion. Used in all cases deficient in biliary secretion.		
<b>PALMETTO COMP .....</b>	{ Ext.Saw Palmetto..1 gr. Ext. Yerba Santa...1 gr. Sugar .....7 gr. Oil Wintergr...1-10 min.}	<b>2 25</b>
Dissolve one in the mouth every three hours, for phthical cough.		
<b>PANCREATIS COMP.....</b>	{ Ext. Pancreatis.....2 gr. Sodium Bicarb.....3 gr.}	<b>3 00</b>
One after eating, for intestinal dyspepsia.		
<b>PAREGORIC 5 minims.....</b> Sedative and carminative.	{ Opium Powd.....1-50 gr. Camphor.....1-50 gr. Acid Benzoic....1-50 gr. Oil Anise.....1-50 gr.}	<b>00</b>
One to three for an infant; best given dissolved in hot water on the end of a spoon.		
<b>PAREGORIC 10 minims.....</b>	{ Opium Powd.....1-25 gr. Camphor.....1-25 gr. Acid Benzoic....1-25 gr. Oil Anise.....1-25 gr.}	<b>1 10</b>
One to six at a dose according to age of patient, given in hot water if practicable.		
<b>PEPPERMINT OIL.....</b>	1-20 min.....	<b>75</b>
One to two, dissolved in hot water on the end of a spoon, for a carminative in infantile colic.		
<b>PEPSIN PURE.....</b>	1-20 gr.....	<b>75</b>
Give one after each nursing to an infant with regurgitation and colic.		

When ordering or prescribing

<b>PEPSIN PURE</b> .....	1-4 gr.....	1 00
One to four tablets after meals, for tardy and incomplete stomach digestion.		
<b>PEPSIN PURE</b> .....	1 gr.....	2 00
One, three times a day, after eating, for enfeebled digestion.		
<b>PEPSIN PURE</b> .....	3 gr.....	3 50
A tablet after eating, for dyspepsia.		
<b>PEPSIN COMP.</b> .....	<div> <div> Pepsin Pure.....1-4 gr.)  Ext. Nux Vom.....1-32 gr.)  Ipecac Powd.....1-16 gr.) </div> </div>	1 25
(Dr. Still) One to two tablets directly after meals, repeated every half hour, if necessary, for three doses; as a tonic and aid to digestion.		
<b>PEPSIN, BISMUTH &amp; CHAR-</b>	<div> <div> Pepsin Conc.....2 gr.)  Bismuth Subnit.....2 gr.)  COAL.....Charcoal.....2 gr.) </div> </div>	2 50
Absorbent stomachic and digestive. A tablet twenty to thirty minutes after meals, for delayed and imperfect digestion with pyrosis and eructation.		
<b>PEPSIN, BISMUTH &amp; GINGER</b>	<div> <div> Pepsin Sacch.....2 gr.)  Bismuth Subnit.....3 gr.)  Ginger.....1 gr.) </div> </div>	2 75
One, three or four times a day, for dyspepsia and flatulence of sedentary life.		
<b>PEPSIN, BISMUTH &amp; STRYCH-</b>	<div> <div> Pepsin Pure.....1 gr.)  Bismuth Subnit.....2 gr.)  NINE.....Strych. Sulph.....1-60 gr.)  Plain or Chocolate Coated </div> </div>	2 25
Give one after meals and one at night, decreasing to two a day as symptoms improve; for atonic dyspepsia.		
<b>PEPSIN &amp; BISMUTH</b> .....	<div> <div> Pepsin Pure.....1-4 gr.)  Antidyspeptic.....Bismuth Subnit.....1 gr.) </div> </div>	1 50
One or two at a time, for functional disorders of the stomach.		
<b>PEPSIN &amp; CALOMEL</b> .....	<div> <div> Pepsin Pure.....1-4 gr.)  Bismuth Subnit.....1 gr.)  Nux Vomica.....1-100 gr.)  Calomel.....1-40 gr.) </div> </div>	1 50
Three to six a day, as a stomachic tonic and intestinal and hepatic stimulant, for chronic dyspepsia with constipation.		
<b>PEPSIN &amp; CHARCOAL</b> .....	<div> <div> Pepsin Pure.....1-4 gr.)  Carbo Veg.....1-4 gr.) </div> </div>	1 00
One every hour or two, for flatulent dyspepsia.		
<b>PEPSIN &amp; PANCREATIN</b> .....	<div> <div> Pepsin Pure.....1-4 gr.)  Digestive.....Pancreatin.....1-4 gr.) </div> </div>	1 50
One to three after meals, for incomplete and tardy digestion, inability to digest oils and fats, and in the treatment of anæmia and emaciation.		
<b>PEPSIN, PANCREATIN COMP.</b>	<div> <div> Pepsin Pure.....1 gr.)  Pancreatin Pure.....1 gr.)  Celery Seed.....1-4 gr.)  Ac. Lactoph. Calc.1-4 gr.)  Plain or Chocolate Coated </div> </div>	3 00
Two after each meal; repeated every two hours, if necessary.		

PEPSIN, PANCREATIN COMP. No. 2.....	{ Pepsin Pure.....3-4 gr. Pancreatin Pure...3-4 gr. Bismuth Subnit.....1 gr. Ext. Nux Vom...1-10 gr. Sodium Bicarb.....2 gr. Ginger Powd.....1-2 gr. Plain or Chocolate Coated	2 50
One to two after meals, for heartburn and flatulence due to indigestion.		
PEPSIN, PANCREATIN COMP. No. 3.....	{ Pepsin Pure.....1 gr. Pancreatin.....1 gr. Nux Vomica.....1-4 gr. Gingerine.....1-16 gr. Plain or Chocolate Coated	3 00
One or two after eating or as desired, for stomachic or intestinal indigestion in all forms dependent upon deficiency in the activity of the secretions, or upon indiscretion in eating.		
PEPSIN, PANCREATIN COMP. Double Strength .....	{ Pepsin Pure.....2 gr. Pancreatin Pure....2 gr. Celery Seed.....1-2 gr. Ac. Lactoph. Calc.1-2 gr.	4 00
One after each meal, repeated in two hours if necessary; two as an aid to digestion.		
PEPSIN SACCHARATED .....	1 gr.	1 10
One every two hours with some nutriment, in the forced nutrition of consumption and during convalescence.		
PEPSIN SACCHARATED .....	5 grs.	1 75
A tablet after meals, as an aid to stomach digestion.		
PEPTIC .....	{ Pepsin Pure.....1-4 gr. Bismuth Subnit....1 gr. Nux Vom. Po...1-100 gr. Tinct. Ginger.....1 min. Spts. Menth Pip...1 min.	1 75
(Dr. North.)		
One to four after meals, for indigestion.		
PHENACETINE.....	1 gr.	5 00
One every two hours for a child; for fever with pain and restlessness. Five at a dose, repeated as necessary, for neuralgia, rheumatism and headache.		
PHENACETINE .....	2 grs.	9 00
Analgesic and antipyretic.	Per 100.....	90
One every two hours for migraine; to quiet restlessness and to reduce fever		
PHENACETINE.....	3 grs.	12 50
	Per 100.....	1 25
One every hour, for the fever and pain of acute rheumatism.		
PHENACETINE.....	4 grs.	15 50
	Per 100.....	1 00
One or two at a dose, repeated as necessary, to depress temperature and quiet pain and restlessness, in fever and rheumatism.		
PHENACETINE.....	5 grs.	18 50
	Per 100.....	1 85
One or two at a dose, repeated as necessary, to depress temperature and quiet pain and restlessness, in fever and rheumatism.		
PHENACETINE, CAFFEINE COMP .....	{ Phenacetine.....3 1-2 gr. Caffeine.....1 gr. Sodium Bromide.7 1-2 gr.	17 75
(Dr. Hubbard.)		
For adults, two for the first dose, and two more after an interval of four hours, repeated if necessary; for neuralgia and headache.		

When ordering or prescribing

PHENACETINE COMP.....	{ Antipyrine.....2 gr. }	13 00
	{ Phenacetine.....2 gr. }	
	{ Caffeine Cit.....1 gr. }	
	{ Sugar of Milk.....1 gr. }	

One to four, according to the height of the temperature, as an antipyretic in all fevers; with analgesic and diuretic secondary effects.

PHENACETINE COMP, No. 2.. (Dr. J. W. Malone.)	{ Phenacetine.....3 gr. }	15 00
	{ Sodium Bicarb.....1 gr. }	

One tablet every two or three hours.

PHOSPHATES..... Tonic.	{ Iron Phosphate.....1 gr. }	2 75
	{ Quinine Phosphate.....1 gr. }	
	{ Strych. Phos.....1-60 gr. }	
Plain or Chocolate Coated		

One three or four times a day as a general tonic and antimalarial; decrease the dose if ringing in the ears and headache occur.

PICROTOXIN.....	1-50 gr.....	1 10
-----------------	--------------	------

One every four to six hours, for spinal irritation, petit mal and hyperidrosis.

PICROTOXIN.....	1-30 gr.....	1 50
-----------------	--------------	------

One at night, to control the sweating in phthisis and rheumatism.

PIL. DUO..... (Dr. McMaster.)	{ Blue Mass.....1 gr. }	1 25
	{ Iron Sulphate.....2 gr. }	

One after eating, three times a day for one year, in early stages of syphilis.

PILOCARPINE MUR.....	1-100 gr.....	1 50
----------------------	---------------	------

One every hour, for pleurisy and other serous inflammations.

PILOCARPINE MUR.....	1-50 gr.....	1 75
----------------------	--------------	------

Give a tablet three to six times a day, to increase the flow of saliva and urine.

PILOCARPINE MUR.....	1-20 gr.....	3 00
----------------------	--------------	------

One to three a day, as a sialogogue, diuretic and diaphoretic.

PILOCARPINE MUR.....	1-10 gr.....	5 00
----------------------	--------------	------

One tablet, for a diaphoretic and diuretic action, repeated with care after three or four hours.

PINK TABLETS (Sugar of Milk).....		50
-----------------------------------	--	----

PLEURISY, ACUTE..... (Dr. J. Harrigan.)	{ Sodium Salicyl..2 1-2 gr. }	1 85
	{ Morphine Sulph. 1-16 gr. }	
	{ Tr. Bryonia.....1 min. }	

Two every two hours, for adults, to relieve acute pleurisy.

PLUMMER.....	1-1000 gr.....	75
--------------	----------------	----

One to three, morning and night, as an alterative, for a child.

PLUMMER.....	1-100 gr.....	75
--------------	---------------	----

One or two every two hours, as an hepatic stimulant to the liver, and to rouse glandular and lymphatic action.

PLUMMER.....	1-10 gr.....	75
--------------	--------------	----

One every two hours.

PODOPHYLLIN.....	1-100 gr.....	75
------------------	---------------	----

One every hour for children, until effect is produced, for constipation with clay colored fæces.

PODOPHYLLIN.....	1-40 gr.....	75
------------------	--------------	----

One every hour until the bowels move freely, then one after meals; for chronic hepatic atony.

<b>PODOPHYLLIN</b> .....	1-20 gr.....	75
Two or three at night and three during the next day, to stimulate the excretion of bile in chronic constipation.		
<b>PODOPHYLLIN</b> .....	1-10 gr.....	75
	Chocolate Coated.....	85
One every hour or two, for bilious headache; then two at night, for several days.		
<b>PODOPHYLLIN</b> .....	1-8 gr.....	75
	Chocolate Coated.....	85
One after meals for biliousness and beginning jaundice.		
<b>PODOPHYLLIN</b> .....	1-4 gr.....	75
	Chocolate Coated.....	85
One to two at night, and one every three hours next day until a cathartic effect is produced, for biliousness and constipation.		
<b>PODOPHYLLIN</b> .....	1-2 gr.....	1 25
	Chocolate Coated.....	1 35
A tablet at night, for biliousness and habitual headache.		
<b>PODOPHYLLIN, BELLADON- NA COMP.</b> .....	<div> <div>Podophyllin.....1-20 gr.</div> <div>Ext. Colocynth Co...2 gr.</div> <div>Pil. Rhei Comp.....2 gr.</div> <div>Ext. Belladonna...1-10 gr.</div> </div>	3 50
(Dr. Francis H. Stuart.)		
One at bedtime, as a laxative—recommended as a post-partum laxative.		
<b>PODOPHYLLIN COMP.</b> .....	<div> <div>Podophyllin.....1-2 gr.</div> <div>Ext. Nux Vom...1-10 gr.</div> <div>Ext. Hyoscyam...1-8 gr.</div> </div>	1 50
Tonic laxative.		
	Chocolate Coated.....	1 00
One or two at bedtime, for atonic conditions of the intestine, and as a stimulant to the liver.		
<b>PODOPHYLLIN &amp; LEPTAN- DRIN</b> .....	<div> <div>Podophyllin.....1-4 gr.</div> <div>Leptandrin.....1-2 gr.</div> </div>	1 50
Hepatic.		
From one to four at night, or divided through the day, for strong hepatic stimulation in functional disorders of the liver.		
<b>POTASS. ARSENITIS</b> .....	1-5000 gr.....	75
Alterative.		
One tablet to a child, repeated every hour, as a tonic and for skin diseases, also for some kinds of diarrhœa. One tablet equals one-fiftieth minim of Fowler's solution.		
<b>POTASS. ARSENITIS</b> .....	1-1000 gr.....	75
One tablet every two hours for diseases of the skin, chronic diarrhœa, neuralgia, etc. One tablet equals one-tenth minim of Fowler's solution.		
<b>POTASS. ARSENITIS</b> .....	1-500 gr.....	75
One every three hours for tonic, also for skin disease; dose may be slowly increased. One tablet equals one-fifth minim of Fowler's solution.		
<b>POTASS. ARSENITIS</b> .....	1-100 gr.....	75
One tablet equals one minim of Fowler's solution; used in skin diseases, useful also in coryza and some forms of diarrhœa.		
<b>POTASS. ARSENITIS</b> .....	1-50 gr.....	75
One after meals, increased slowly, for anæmia, chorea and chronic skin diseases. One tablet equals two minims of Fowler's solution.		
<b>POTASS. ARSENITIS</b> .....	1-40 gr.....	75
One three times a day for chronic rheumatism, malarial poisoning and as an alterative. One tablet equals two and a half minims of Fowler's solution.		

When ordering or prescribing



<b>POTASS. IODIDE</b> .....	2 grs.....	2 50
One every hour or two, for chronic rheumatism and diseases of the skin; with free use of drinking water.		
<b>POTASS. IODIDE</b> .....	5 grs.....	5 00
One, three times a day, increase gradually to four at a dose, for secondary syphilitic diseases; one every three hours with a swallow of water, for asthma.		
<b>POTASS. IODIDE, Modified</b> ....	{ Potass. Iodide.....5 gr. } { Tr. Belladon....2 1-2 min. }	6 00
One, three times a day, increase gradually to four at a dose, for secondary syphilitic diseases; one every three hours with a swallow of water, for asthma.		
<b>POTASS. IODIDE Modified No. 2</b>	{ Potass. Iodide.....5 gr. } { Tr. Belladon....1 1-4 min. }	6 00
One, three times a day, increase gradually to four at a dose, for secondary syphilitic diseases; one every three hours with a swallow of water, for asthma.		
<b>POTASS. NITRATE</b> .....	5 grs.....	75
One every three hours, as a diuretic and laxative; lessens fever and pulse-rate, and relieves testicular neuralgia.		
<b>POTASS. PERMANGANATE</b> ....	1 gr.....	1 00
One every three hours, for acute rheumatism, diphtheria and zymotic diseases. Also one, three times daily as an emmenagogue; also useful in amenorrhœa.		
<b>POTASS. PERMANGANATE</b> ....	2 grs.....	1 15
One, three times daily as an emmenagogue. Also one to four to a pint of hot water, as an injection in vaginitis.		
<b>PRORHINITIS</b> .....	{ Merc. Iod. Red...1-200 gr. } { Ipecac Powd....1-50 gr. } (Dr. Gilbert.) { Belladonna Po...1-100 gr. }	75
One every two hours, for two or three days, then every four hours as long as necessary. Useful in simple catarrhal conditions and inflammations of a subacute or chronic nature.		
<b>PROSTATIC COMP.</b> .....	{ Ext. Saw Palmetto...2 gr. } (Dr. C. R. P. Fisher.) { Ext. Nux Vom....1-8 gr. }	2 75
<b>Plain or Chocolate Coated</b>		
One or two before meals and on retiring, for irritability and chronic enlargement of the prostate, and loss of power in the bladder.		
<b>PUERPERAL ANTISEPTIC</b> ....	{ Mercury Bichlor.1-48 gr. } (Dr. C. W. Stevens.) { Potass. Chlorate...1-4 gr. }	75
Give two every two hours unless diarrhœa follows, then reduce the dose to half a tablet.		
<b>QUININE ARSENATE</b> .....	1-64 gr.....	90
<b>Chocolate Coated</b> .....		1 00
One every three hours, with gradual increase of the dose, for malarial anæmia and cachexia.		
<b>QUININE ARSENATE</b> .....	1-50 gr.....	90
<b>Chocolate Coated</b> .....		1 00
One every three hours, increasing by one a week until five are taken, for dumb ague and chronic malaria.		
<b>QUININE ARSENATE</b> .....	1-15 gr.....	1 25
One every three or four hours, for malaria and malarial anæmia.		

When ordering or prescribing

QUININE, ARSENIC AND STRYCHNINE.....	{ Quinine Sulph.....1 gr. }	
	{ Acid Arsenous...1-50 gr. }	
	{ Strych. Sulph....1-60 gr. }	

Plain or Chocolate Coated 2 25

One after meals, for a general tonic effect in anæmia and malarial infection.

QUININE & ATROPINE CO... Tonic.	{ Quinine Bisulph....1 gr. }	2 25
	{ Potass. Arsen...1-200 gr. }	
	{ Atropine Sulph.1-600 gr. }	

One every hour as a tonic sedative, to relieve restlessness and irritability; also in the fevers of children as an antipyretic.

QUININE & CAPSICUM . . . . .	{ Quin. Sulph.....2 gr. }	2 25
	{ Capsicum Powd...1-4 gr. }	
	Chocolate Coated only	

One to three tablets, for an antiperiodic and stimulant.

QUININE & CHOCOLATE..... Tonic for children.	{ Quinine Tannate....1 gr. }	3 75
	{ Chocolate.....9 gr. }	

One to three at a time, as an agreeable tonic for children.

QUININE & HYOSCYAMUS CO Sedative, tonic.	{ Morph. Sulph....1-40 gr. }	2 50
	{ Ext. Hyoscyam...1-10 gr. }	
	{ Quinine Sulph.....1 gr. }	

From one to four a day, for the irritability of exhaustion, as a general tonic and sedative.

QUININE & LICORICE . . . . .	{ Quinine Sulph....1-10 gr. }	1 10
	{ Ext. Licorice....1-10 gr. }	

One after meals, for children as a tonic and regulator.

QUININE BISULPHATE.....1-10 gr.....	1 00
-------------------------------------	------

One, three times a day, as an antiperiodic and tonic, for a child.

QUININE BISULPHATE.....1-4 gr.....	1 10
------------------------------------	------

One, half an hour before eating, as a stomachic tonic.

QUININE BISULPHATE.....1-2 gr.....	1 25
	Chocolate Coated... .. 1 35

A tablet three or more times a day, often combined with iron to advantage, as a general tonic.

QUININE BISULPHATE.....1 gr.....	1 50
	Chocolate Coated..... 1 60

One after meals as a tonic and antiperiodic; one every hour or two as an antipyretic in children's fevers.

QUININE BISULPHATE.....2 grs.....	2 25
	Plain or Chocolate Coated.

One after meals to strengthen the pulse, increase muscular force, and improve the tone of the nervous system.

QUININE BISULPHATE.....3 grs.....	3 00
	Plain or Chocolate Coated.

One every two hours, with free use of drinking water, as an antipyretic for coughs and fevers in childhood.

QUININE BISULPHATE.....4 grs.....	4 00
	Plain or Chocolate Coated.
	Per 100..... 50

One every hour to four doses, unless relieved sooner, for migraine and malarial hemicrania; one every three hours as a malarial tonic.

<b>QUININE BISULPHATE</b> .....	5 grs.....	5 00
	<b>Plain or Chocolate Coated.</b>	
	<b>Per 100</b> .....	60
A tablet every two or three hours, as an antipyretic in fever; for chills, and fever, a tablet, at six, four, and two hours before the chill.		
<b>QUININE COMP</b> .....	<div> <div>Quinine Sulph.....1 gr.</div> <div>Acid Arsenous...1-40 gr.</div> <div>Iron Powd.....1 gr.</div> <div>Strych. Sulph.....1-40 gr.</div> <div>Oil Bl'k Pepper. 1-3 min.</div> </div>	
General tonic.	<b>Plain or Chocolate Coated</b>	2 25
One or two after meals, as a general tonic in enfeebled and anæmic conditions, with or without a malarial element.		
<b>QUININE, DIGITALIS COMP.</b> ..	<div> <div>Quinine Sulph....1-2 gr.</div> <div>Strych. Sulph....1-100 gr.</div> <div>Digitalis Powd....1-2 gr.</div> <div>Iron Powd.....1-2 gr.</div> </div>	1 50
(Dr. Fowler.)	<b>Chocolate Coated</b> .....	1 60
Heart and general tonic.		
A tablet three or four times a day, for anæmia, feeble or irregular heart action, and nervous exhaustion.		
<b>QUININE, IODINE COMP</b> .....	<div> <div>Quinine Sulph.....2 gr.</div> <div>Acid Arsen.....1-40 gr.</div> <div>Iodine (Free) ....1-12 gr.</div> </div>	3 25
(Dr. Duncan.)		
Alterative.		
<b>QUININE, IRON &amp; ARSENIC</b> ..	<div> <div>Quinine Sulph....1-2 gr.</div> <div>Iron Powd.....1-2 gr.</div> <div>Acid Arsenous. 1-120 gr.</div> </div>	1 50
Hæmatic.	<b>Chocolate Coated</b> .....	1 60
One every three or four hours, increasing gradually, for simple anæmia and all enfeebled conditions with pallor.		
<b>QUININE SULPHATE</b> .....	1-10 gr.....	95
A tablet three or four times a day, to strengthen the pulse and invigorate the nervous system.		
<b>QUININE SULPHATE</b> .....	1-4 gr.....	1 10
	<b>Chocolate Coated</b> .....	1 20
Three or four a day, for anæmia and during convalescence; best combined with iron.		
<b>QUININE SULPHATE</b> .....	1-2 gr.....	1 25
	<b>Chocolate Coated</b> .....	1 35
One every hour, as a febrifuge and antiperiodic for children, stop if headache and ringing of the ears occur.		
<b>QUININE SULPHATE</b> .....	1 gr.....	1 50
	<b>Chocolate Coated</b> .....	1 60
One every hour, in acute malarial fevers; when symptoms begin to improve, decrease to every three hours, and then to three times a day; one after meals, as a general tonic.		
<b>QUININE SULPHATE</b> .....	2 grs.....	2 25
	<b>Plain or Chocolate Coated.</b>	
One, three or four times a day after meals, as a tonic.		
<b>QUININE SULPHATE</b> .....	3 grs.....	3 00
	<b>Plain or Chocolate Coated.</b>	
One for tonic three times a day—two to three for breaking up a cold, generally given with five to ten grains of Dover's Powder.		
<b>QUININE SULPHATE</b> .....	4 grs.....	4 00
	<b>Plain or Chocolate Coated.</b>	
A tablet three or four times a day as a tonic; one every two hours for neuralgia.		

**QUININE SULPHATE** ..... 5 grs. .... 5 00  
**Plain or Chocolate Coated.**

A tablet at six, four, and two hours before the chill in malarial fever; one or two at night, followed by another next morning, generally combined with Dover's Powder, for recent colds.

**REALGAR**, see Arsen. Red Sulph.

**RHEUMATIC** ..... { Acid Salicylic.....3 gr. }  
 (Dr. J. F. Reilley.) { Wine Colchicum...1 min. } 2 50  
 { Resin Guaiac.....3 gr. }

One every two hours until pain is relieved.

**RHEUMATIC No. 3** ..... { Sodium Salicyl...7 1-2 gr. }  
 { Sodium Bicarb.....2 gr. } 2 50  
 { Wine Colch. S. 2 1-2 min. }  
 { Oil Wintergreen....q. s. }

One to two tablets every three hours, for acute and chronic rheumatism, and rheumatic pharyngitis.

**RHEUMATISM** ..... { Sodium Salicylate...5 gr. }  
 (Dr. Liggett.) { Colchicin.....1-120 gr. } 3 50  
 { Codeine Sulph....1-16 gr. }

One tablet every three hours, until rheumatic pain ceases and a laxative action is obtained.

**RHEUMATISM No. 3** ..... { Sodium Salicylate..5 gr. }  
 { Colchicin.....1-120 gr. } 3 90  
 { Codeine Sulph...1-16 gr. }  
 { Macrotin.....1-10 gr. }  
 { Ext. Poke Root...1-2 gr. }

One tablet every three or four hours until pain is relieved, and a laxative action obtained.

**RHINITIS, One-half strength**... { Camphor.....1-4 gr. }  
 (Dr. Lincoln.) { Fl. Ext. Bellad...1-8 min. } 1 50  
 { Quinine Sulph....1-4 gr. }  
**Chocolate Coated**..... 1 60

One every half hour for four doses, then every hour until throat feels dry; afterwards every three hours until a day after the acute coryza disappears.

**RHINITIS, Full Strength**..... { Camphor.....1-2 gr. }  
 { Fl. Ext. Bellad...1-4 min. }  
 { Quinine Sulph....1-2 gr. }  
**Plain or Chocolate Coated**..... 2 00

One every hour for four doses, then every hour and a half until throat feels dry; afterwards every three hours until a day after acute coryza disappears.

**RHINITIS No. 2**..... { Camphor.....1-4 gr. }  
 (Dr. B. S. Booth.) { F. E. Belladonna.1-8 min. } 1 10  
 { Tr. Euphrasia....1 min. }  
**Chocolate Coated**..... 1 20

One every half hour for eight doses; then one every hour until relieved.

**RHUBARB**..... 1-2 gr. .... 75

One every two hours, for a tonic to the intestinal canal, and mild astringent laxative.

**RHUBARB** ..... 1 gr. .... 75

One every hour until the bowels move, for fecal accumulation; one after meals, repeated in half an hour, for gastric and duodenal catarrh.

**RHUBARB COMP. U.S.P.**..... 1-2 gr. .... 75

One after each meal or two at night, as a laxative; increase for purgative effect.

RHUBARB COMP. U. S. P.....	{ Rhubarb Powd.....2 gr. }	1 50
	{ Myrrh Powd.....1 gr. }	
	{ Aloes Purified...1-2 gr. }	
	{ Oil Peppermint.1-10 min. }	
	Chocolate Coated.....	1 60

One to three at night, for the relief of habitual constipation with flatulence, due to atony of the intestines.

RHUBARB & GINGER CO....	{ Rhubarb Powd...1 1-4 gr. }	1 60
	{ Sodium Bicarb...1 1-4 gr. }	
	{ Ginger Powd...2 1-2 gr. }	
	{ Cardamom Powd.5-8 gr. }	
	Oil Peppermint.1-20 min.	

A tablet after meals, for dyspepsia, flatulence and constipation.

RHUBARB & HYOSCYAMUS COMP.....	{ Rhubarb Powd.....3 gr. }	2 00
	{ Calomel.....1 gr. }	
	{ Ext. Hyoscyamus...1 gr. }	

Alterative and laxative.

A tablet in the morning and one at night, for constipation with collection of mucus in the lower bowel, and dry stools, with flatulence and headache.

RHUBARB & IPECAC.....	{ Rhubarb Powd...1-4 gr. }	75
	{ Ipecac Powd.....1-10 gr. }	

(Dr. Hitchcock.)

Children's laxative.

One at a dose; four to eight daily, as needed in habitual duodenal torpidity, with lack of bile in the passages.

RHUBARB & IPECAC COMP..	{ Rhubarb Powd.....1 gr. }	1 10
	{ Sodium Bicarb.....5 gr. }	
	{ Ipecac Powd.....1-8 gr. }	
	{ Oil Peppermint.1-20 min. }	
	Chocolate Coated.....	1 20

(Roosevelt Hospital)

Alkaline laxative and stomachic

One after each meal, for indigestion with constipation.

RHUBARB & IPECAC COMP.	{ Rhubarb Powd.....2 gr. }	1 50
	{ Sodium Bicarb.....5 gr. }	
	{ Ipecac Powd.....1-8 gr. }	
	{ Oil Peppermint.1-20 min. }	

No. 2.....  
(Roosevelt Hospital.)

One after each meal, for indigestion with constipation.

RHUBARB & IPECAC COMP.	{ Rhubarb Powd.....2 gr. }	1 60
	{ Ipecac Powd.....1-4 gr. }	
	{ Sodium Bicarb.....5 gr. }	
	{ Tr. Nux Vomica...5 min. }	
	Oil Peppermint.1-20 min.	

No. 3.....  
(Roosevelt Hospital.)

One after each meal, for indigestion with constipation.

RHUBARB & IPECAC COMP.	{ Rhubarb Powd.....1 gr. }	1 80
	{ Sodium Bicarb.....5 gr. }	
	{ Ipecac Powd.....1-8 gr. }	
	{ Ext. Nux Vom...1-4 gr. }	
	{ Oil Peppermint.1-10 min. }	
	{ Fl. Ext. Cascara...5 min. }	

No. 4.....  
(Dr. M. L. Goodkind.)

One just before meals for indigestion, either stomachic or intestinal, with sluggishness of hepatic functions and constipation.

RHUBARB & IPECAC CO. No. 4	{ Rhubarb Powder...1-2 gr. }	1 25
	{ Sodium Bicarb...2 1-2 gr. }	
	{ Ipecac Powder...1-16 gr. }	
	{ Ext. Nux Vomica.1-8 gr. }	
	{ Oil Peppermint.1-20 min. }	
	{ Fl. Ext. Cascara.2 1-2 min. }	
	Chocolate Coated only	

Half Size.....  
(Dr. Goodkind.)

One or two before meals for indigestion, either stomachic or intestinal, with sluggishness of hepatic functions and constipation.

When ordering or prescribing



<b>SALICYLATE COMP.</b> .....	{ Sodium Salicyl...2 1-2 gr. }	2 65
Antipyretic.	{ Acetanilid.....5 gr. }	

One or two at the first dose, followed by one every three hours if necessary ; for rheumatism with pain and high temperature.

<b>SALINE INFUSION</b> .....	{ Sodium Chlor...11 2-3 gr. }	2 00
(Dr. Geo. Ryerson Fowler.)	{ Sodium Sulphate....1 gr. }	
	{ Sodium Carb.....2-3 gr. }	
	{ Sodium Phos.....1-3 gr. }	
	{ Magnesia Phos..1 2-3 gr. }	

A combination of the soluble salts of the blood in the proportions in which they exist in the body. Use one tablet to a pint of water previously sterilized by boiling, to wash away chemical antiseptic agents from wounds after irrigation, or as a saline infusion in cases of excessive hemorrhage, shock, etc.

<b>SALINE INFUSION No. 2</b> .....	{ Sodium Chlor...9 4-5 gr. }	1 50
(Dr. Geo. Ryerson Fowler.)	{ Sodium Sulphate..2-5 gr. }	
	{ Sodium Carbon...1-3 gr. }	
	{ Sodium Phosph...1-3 gr. }	
	{ Magnes. Phos...4-11 gr. }	

Use two tablets to a pint of water, previously sterilized by boiling, to wash away chemical antiseptic agents from wounds after irrigation, or as a saline infusion in case of excessive hemorrhage, shock, etc.

<b>SALOL</b> .....	1-10 gr.....	75
--------------------	--------------	----

Intestinal antiseptic.

One every hour, as an intestinal antiseptic for diarrhœa and dysentery of children.

<b>SALOL</b> .....	1-2 gr.....	1 35
--------------------	-------------	------

A tablet every hour or two, as an intestinal antiseptic for diarrhœa and dysentery of children.

<b>SALOL</b> .....	1 gr.....	1 70
--------------------	-----------	------

One every hour, for inflammation of the bladder and urethra, combined with oil sandalwood (2 drop) tablet ; for catheter fever, fermentation of urine in the bladder, and gonorrhœa.

<b>SALOL</b> .....	2 1-2 grs.....	3 00
--------------------	----------------	------

One every two hours, as an intestinal antiseptic, and one every hour for acute rheumatism.

<b>SALOL</b> .....	3 grs.....	3 65
--------------------	------------	------

A tablet three or four times a day, as an intestinal antiseptic.

<b>SALOL</b> .....	5 grs.....	5 50
--------------------	------------	------

One to four tablets every two or three hours, as an antipyretic and intestinal antiseptic.

<b>SALOL COMP. No. 2</b> .....	{ Salol.....1 gr. }	2 50
	{ Bismuth Subnit.....1 gr. }	
	{ Opium Powd.....1-40 gr. }	
	{ Camphor Powd...1-20 gr. }	

Give one every hour for four doses, then every two hours ; for fermentative diarrhœa of children, with pain.

<b>SALOL &amp; COPAIBA CO</b> .....	{ Salol.....2 gr. }	4 50
	{ Balsam Copaiba....2 gr. }	
	{ Oleoresin Cubebs...1 gr. }	
	{ Pepsin Pure.....1-2 gr. }	
	{ Ext. Kava Kava....1 gr. }	
	{ Oil Sandalwood..1-4 min. }	
	{ Oil Wintergreen.1-4 min. }	

Chocolate Coated only

One tablet three or four times daily. Renders urine bland, antiseptic, and non-irritating, diminishes urethral discharge, and relieves pain of urinating.

<b>SALOL &amp; COPAIBA COMP.</b> One-half Strength.....	Salol.....	1 gr.	} 2 50
	Balsam Copaiba.....	1 gr.	
	Oleoresin Cubebs.....	1-2 gr.	
	Pepsin Pure.....	1-4 gr.	
	Ext. Kava Kava.....	1-2 gr.	
	Oil Sandalwood.....	1-8 min.	
	Oil Wintergreen.....	1-8 min.	

Two tablets three or four times daily. Renders urine bland, antiseptic, and non-irritating, diminishes urethral discharge, and relieves pain of urinating.

<b>SALOL &amp; PHENACETINE....</b> Analgesic and antiseptic	Salol.....	2 1-2 gr.	} 14 00
	Phenacetine.....	2 1-2 gr.	
	Per 100.....		
			1 50

One every half hour for three doses, then every two hours, for fever and neuralgia.

<b>SALOL AND PHENACETINE</b> No. 2.....	Salol.....	1 1-4 gr.	} 7 00
	Phenacetine.....	1 1-4 gr.	

One every hour for four or five doses, then every two hours, for fever, with pain and neuralgia.

<b>SANGUINARIA COMP.</b> (Dr. O. A. Gordon.)	Tartar Emetic.....	1-96 gr.	} 2 00
	Codeine.....	1-24 gr.	
	Tr. Sanguinaria.....	2 min.	
	Oil Anise.....	q. s.	

One every hour as an expectorant.

<b>SANGUINARIA &amp; CODEINE..</b>	Tr. Sanguinaria.....	1 min.	} 1 30
	Codeine.....	1-25 gr.	

One every hour, as a sedative and expectorant in laryngitis, bronchitis and whooping cough; stop as symptoms improve.

<b>SANTONIN.....</b>	1-10 gr.....	1 00
Vermifuge.		

One to four tablets every two hours, followed by a cathartic; as a vermifuge for children.

<b>SANTONIN.....</b>	1-2 gr.....	1 25
----------------------	-------------	------

One or two, according to age, every two hours or oftener, followed by a cathartic; as a vermifuge.

<b>SANTONIN.....</b>	1 gr.....	2 00
----------------------	-----------	------

One every three hours for not more than six doses, followed by a cathartic; as a vermifuge for children.

<b>SANTONIN &amp; CALOMEL.....</b> Vermifuge.	Santonin.....	1-2 gr.	} 1 65
	Calomel.....	1-2 gr.	

One to two, according to age, every two hours until cathartic action is produced; as a vermifuge for children.

<b>SANTONIN &amp; CALOMEL No. 2.</b>	Santonin.....	1 gr.	} 2 10
	Calomel.....	1 gr.	

One every three hours until cathartic action is produced, or not more than four taken; as a vermifuge for children.

<b>SANTONIN &amp; PODOPHYLLIN</b> (Dr. Sidney.)	Santonin.....	1-2 gr.	} 1 50
	Podophyllin.....	1-20 gr.	

One every hour until the bowels move, for round worms in children; lessen the dose for the very young.

<b>SCIATICA.</b> (Dr. Jno. T. Metcalf.)	Tr. Aconite.....	3-4 min.	} 1 10
	Tr. Belladonna.....	3-4 min.	
	Tr. Colchicum.....	3-4 min.	
	Tr. Cimicifuga.....	3-4 min.	

Two tablets, every three hours for rheumatism and sciatica.

<b>SEDATIVE</b> .....	{ Sodium Brom....2 1-2 gr. Potass. Brom....2 1-2 gr. Ammon. Brom...2 1-2 gr. Tr. Hyoscyamus...5 min. Tr. Cannabis Ind...5 min. }	2 50
(Dr. Brown.)		

One dissolved in water, repeated every half hour, for three or four doses at night; for wakefulness due to overwork, delirium and nervous neuralgia; also for irritable bladder and emissions.

<b>SEDATIVE No. 3</b> .....	{ Ext. Valerian.....1 gr. Ext. Sumbul.....1 gr. Asafœtida Powd...1 gr. }	2 00
(Dr. Baer.)		
	<b>Chocolate Coated only</b>	

One tablet every three or four hours and one at bedtime, for relief of nervousness, hysteria, and as a general sedative.

<b>SEDATIVE No. 4</b> .....	{ Can. Ind. Fl. Ext...2 min. }	1 75
(Dr. O. W. Sadler.)	{ Gelsem. Fl. Ext...2 min. }	

One or two tablets at a dose, repeated every hour for two doses, if necessary. In headache, trigeminal neuralgias and ovarian and dysmenorrhœal pains.

<b>SEDATIVE No. 5</b> .....	{ Ext. Sumbul.....1-2 gr. Ext. Hyoscyamus. 1-2 gr. Ext. Valerian.....1-2 gr. Ext. Can. Ind. ....1-10 gr. }	2 25
	<b>Chocolate Coated only</b>	

One, and repeat every hour for three doses, then every three hours as needed.

<b>SEDATIVE, Genital</b> .....	{ Gelsemine Sulph. 1-50 gr. }	14 00
(Dr. W. E. Derry.)	{ Hyosc. Hydrobr...1-100 gr. }	
	<b>Per 100</b> .....	1 50

One at bedtime, for nocturnal emissions. If possible, anticipate emissions and give a tablet that night.

<b>SENEGA COMP.</b> .....	{ Tr. Squill.....1 min. }	1 10
(Dr. Cleveland.)	{ Tr. Senega.....1 min. }	
Expectorant.	{ Tr. Ipecac.....1 min. }	

Dissolve six tablets in twelve teaspoonfuls of water, and give a child a teaspoonful every hour; two teaspoonfuls every hour for an adult.

<b>SENEGA COMP. No. 2</b> .....	{ Ammon. Carb.....2 gr. Ex. Senega U.S.P. 70. 1 gr. Ext. Hyoscyamus. 1-8 gr. Ipecac Powd.....1-8 gr. Oil Cinnamon...1-2 min. }	2 25
(Dr. D. E. English.)		

One tablet every one to four hours, followed by a swallow of water. A stimulant expectorant and sedative, indicated in all acute catarrhal diseases of the respiratory tract, as soon as the feverish and acute stage is past; also in chronic troubles of same nature, and especially in bronchitis of the aged.

<b>SICK HEADACHE</b> .....	{ Irisin.....1-10 gr. Podophyllin....1-20 gr. Sanguinarin....1-20 gr. Nux Vomica Po...1-20 gr. Euonymin.....1-4 gr. }	1 50
(Dr. Hale.)		

One, repeated every hour until effect is produced; for sick headache. Used also as an hepatic corrective.

<b>SODA MINT</b> .....	{ Sodium Bicarb.....5 gr. Oil Peppermint. 1-4 min. }	95
Alkaline stomachic.	{ Arom. Spts. Amm. .2 min. }	
	<b>Chocolate Coated</b> .....	1 05

One as needed, for indigestion, flatulence, heartburn, sick headache, sour stomach, and nausea.

<b>SODA MINT &amp; PEPSIN</b> .....	{ Soda Mint.....1-2 gr. }	1 50
	{ Pepsin.....1-2 gr. }	

One to four after eating, repeated at intervals of half an hour, for gastric distress and dyspepsia.

<b>SODIUM ARSENATE</b> .....	1-100 gr.....	75
------------------------------	---------------	----

Alterative.

One every three or four hours, increasing slowly until five are taken; stop when œdema of the face and eyelids and disturbance of digestion occur; for chronic diseases of the skin.

<b>SODIUM ARSENATE</b> .....	1-60 gr.....	75
------------------------------	--------------	----

One three times a day, as a tonic and antimalarial; stop if gastric symptoms occur.

<b>SODIUM ARSENATE</b> .....	1-50 gr.....	75
------------------------------	--------------	----

One after meals, for chronic eczema, psoriasis, and other skin diseases; increase slowly and stop on occurrence of gastric disorder and puffiness of the face.

<b>SODIUM ARSENATE</b> ....	1-25 gr.....	75
-----------------------------	--------------	----

Begin with one and increase as tolerance is established; stop if the stomach should become deranged; for chorea, epilepsy and malarial neuralgia.

<b>SODIUM ARSENATE</b> .....	1-16 gr.....	75
------------------------------	--------------	----

One after meals, for chronic rheumatism, skin diseases and malarial cachexia; stop the treatment on the appearance of facial œdema and gastric distress.

<b>SODIUM ARSENATE</b> .....	1-10 gr.....	75
------------------------------	--------------	----

A tablet two or three times a day, for chorea, epilepsy, and neuralgia, chronic diseases of the joints and skin, and malarial infection; stop the treatment if symptoms of reaction occur.

<b>SODIUM BENZOATE</b> .....	2 grs.....	1 25
------------------------------	------------	------

One to four tablets according to age, every three or four hours, for rheumatism, gout and uric acid diathesis.

<b>SODIUM BENZOATE</b> .....	5 grs .....	1 75
------------------------------	-------------	------

One to two tablets, three times a day, for rheumatism, gout and uric acid diathesis.

<b>SODIUM BICARBONATE</b> .....	5 grs.....	75
---------------------------------	------------	----

Alkaline.

One or two after eating to correct acidity of the stomach, and as an aid to digestion in dyspepsia. May be repeated in half an hour if necessary.

<b>SODIUM BICARBONATE</b> .....	10 grs.....	75
---------------------------------	-------------	----

A tablet every half hour, for two or three doses, to correct acid dyspepsia.

<b>SODIUM BICARBONATE CO.</b> ..	{ Sodium Bicarb.....2 gr. }	1 25
(Dr. C. G. Rancourt.)	{ Pepsin Saccharated.1 gr. }	
	{ Rhubarb Powd...1-2 gr. }	

Four tablets at a dose after meals; for a child, put one in half a glass of water and give during the day. Useful in gastric fever with a heavily coated tongue.

<b>SODIUM BROMIDE</b> .....	5 grs.....	1 30
-----------------------------	------------	------

Sedative.

One dissolved in water every fifteen minutes for three doses. May be repeated in two hours, for nervousness, restlessness and sleeplessness.

<b>SODIUM BROMIDE</b> .....	10 grs.....	1 75
-----------------------------	-------------	------

One every hour for three doses to induce sleep; one, three or four times a day, for nervous irritability.

<b>SODIUM PHOSPHATE</b> .....5 gr.....	1 00
Three at night as a mild saline laxative; one after each meal for duodenal catarrh, and the uric acid diathesis.	
<b>SODIUM SALICYLATE</b> .....1-2 gr.....	1 00
Antirheumatic, analgesic.	
A tablet every hour in acute rheumatism, as an antipyretic and analgesic.	
<b>SODIUM SALICYLATE</b> .....1 gr.....	1 10
One every hour for rheumatism or sour stomach, or five every three hours for rheumatism.	
<b>SODIUM SALICYLATE</b> .....1 gr.....	1 10
Flavored with Wintergreen.	
A tablet every hour, for the fever and pain of acute rheumatism; diminish as symptoms improve.	
<b>SODIUM SALICYLATE</b> .....2 grs.....	1 25
Antirheumatic.	
One every hour or two in acute articular rheumatism and serous inflammations of rheumatic origin; decrease the dose if nausea and vomiting occur.	
<b>SODIUM SALICYLATE</b> .....3 grs.....	1 40
One every two hours for three doses, for the headache and neuralgia of rheumatic patients.	
<b>SODIUM SALICYLATE</b> .....5 grs.....	1 75
One tablet every two to four hours, for gout and rheumatism.	
<b>SODIUM SALICYLATE</b> .....5 grs.....	1 75
Flavored with Wintergreen.	
One tablet every two to four hours for gout and rheumatism.	
<b>SODIUM SULPHITE</b> .....5 grs.....	1 50
Intestinal antiseptic.	
One, three or four times a day, for intestinal stasis and fermentation.	
<b>SODIUM SULPHOCARB</b> .....1 gr.....	1 10
One every hour for five or six doses, then every two hours, for an intestinal antiseptic and to reduce temperature.	
<b>SODIUM SULPHOCARB</b> .....2 1-2 grs.....	1 35
One or two every hour, to reduce the temperature, correct the bowels and promote intestinal antiseptics in typhoid fever and gastro-enteritis.	
<b>SORE THROAT</b> .....	1 50
Diphtheritic.	
One every two hours.	
<b>SORE THROAT No. 2</b> .....	1 50
For tonsillitis, give one every three hours for children of six to twelve years. Two tablets every four hours for patients above twelve years old. For diphtheria, crush and give one every two or three hours.	
<b>SORE THROAT No. 3</b> .....	75
(Dr. E. G. Rankin.)	
One hourly for sore throat tending to ulceration and bronchitis.	
<b>SPARTEINE SULPHATE</b> .....1-50 gr.....	75
A tablet every two or three hours, to regulate the heart and increase the flow of urine.	
<b>SPARTEINE SULPHATE</b> .....1-20 gr.....	1 10
One every three hours, as a diuretic in renal congestion and to control irregular cardiac action.	

When ordering or prescribing

Please specify Fraser's Tablets

<b>STRYCHNINE MURIATE</b> .....	1-60 gr.....	1 10
One every three hours, as a nerve and stomachic tonic.		
<b>STRYCHNINE MURIATE</b> .....	1-30 gr.....	1 10
One every three hours, as a nerve and stomachic tonic.		
<b>STRYCHNINE MURIATE</b> .....	1-20 gr.....	1 10
One, three times a day, as a general nerve tonic.		
<b>STRYCHNINE MURIATE CO.</b> {	Corrosive Sub....1-20 gr. } Ext. Belladonna.1-10 gr. } Strych. Mur.....1-40 gr. }	1 10
One night and morning for the correction of atonic conditions of the ali- mentary canal and consequent reflex disorders.		
<b>STRYCHNINE NITRATE</b> .....	1-100 gr.....	1 00
A tablet three times a day, as a general tonic in neurasthenia and spinal anæmia.		
<b>STRYCHNINE NITRATE</b> .....	1-60 gr.....	1 00
	<b>Chocolate Coated</b>	1 10
One after meals for amenorrhœa, with anæmia and a tendency to melan- cholia.		
<b>STRYCHNINE NITRATE</b> .....	1-50.....	1 00
One after eating, for dyspepsia with anorexia, constipation and epigastric distress.		
<b>STRYCHNINE NITRATE</b> .....	1-40 gr.....	1 00
	<b>Chocolate Coated</b>	1 10
One after eating, for dyspepsia with anorexia, constipation and epigastric distress.		
<b>STRYCHNINE NITRATE</b> .....	1-30 gr.....	1 00
	<b>Chocolate Coated</b>	1 10
One every four hours, for vesical atony and dyspepsia; decrease as symp- toms improve.		
<b>STRYCHNINE NITRATE</b> .....	1-20 gr.....	1 00
	<b>Chocolate Coated</b>	1 10
One tablet for the weakness and nervous depression following excessive use of alcohol or tobacco; may be repeated in two hours.		
<b>STRYCHNINE SULPHATE</b> ....	1-1000 gr.....	75
One every two hours, as a stomachic and general nerve tonic.		
<b>STRYCHNINE SULPHATE</b> ....	1-500 gr.....	75
One every two hours, as a stomachic and general nerve tonic.		
<b>STRYCHNINE SULPHATE</b> ....	1-200 gr.....	75
One every three hours, as a stomachic and general nerve tonic.		
<b>STRYCHNINE SULPHATE</b> ....	1-100 gr.....	75
	<b>Chocolate Coated</b> .....	85
One, three times a day, as a stomachic and general nerve tonic.		
<b>STRYCHNINE SULPHATE</b> ....	1-60 gr.....	75
	<b>Chocolate Coated</b> .....	85
One after each meal, for a nerve tonic and aid to digestion.		
<b>STRYCHNINE SULPHATE</b> ....	1-50 gr.....	75
	<b>Chocolate Coated</b> .....	85
One after each meal, as a nerve tonic and aid to digestion.		

When ordering or prescribing

<b>STRYCHNINE SULPHATE</b> .....1-40 gr.....	75
Chocolate Coated.....	85
One after each meal, as a nerve tonic.	
<b>STRYCHNINE SULPHATE</b> .....1-30 gr.....	75
Chocolate Coated.....	85
One after each meal, as a nerve tonic.	
<b>STRYCHNINE SULPHATE</b> .....1-20 gr.....	75
Chocolate Coated.....	85
One after each meal, as a nerve tonic.	
<b>STRYCHNINE SULPHATE</b> .....1-16 gr.....	75
Chocolate Coated.....	85
<b>STRYCHNINE SULPHATE</b> .....1-10 gr.....	1 00
Chocolate Coated.....	1 10
<b>STRYCHNINE VALER</b> .....1-100 gr.....	1 10
Chocolate Coated.....	1 20
One, three times a day, for a general nerve tonic.	
<b>STRYCHNINE VALER</b> .....1-60 gr.....	1 10
Chocolate Coated.....	1 20
One, three times a day, for a general nerve tonic.	
<b>STRYCHNINE VALER</b> .....1-50 gr.....	1 10
Chocolate Coated.....	1 20
A tablet night and morning, increasing to three or four a day if well tolerated, for nervous diseases and prostration, and as a tonic aphrodisiac.	
<b>SULPHONAL</b> .....3 grs.....	16 90
Per 100.....	1 70
A tablet every hour or two for three or four doses, for insomnia.	
<b>SULPHONAL</b> .....5 grs.....	26 25
Per 100.....	2 65
One every hour or two, as a hypnotic.	
<b>SULPHONAL</b> .....8 grs.....	40 00
Per 100.....	4 00
One or two, one hour before retiring, repeated every hour for two doses if necessary.	
<b>SULPHONAL</b> .....15 grs.....	70 00
Per 100.....	7 00
One an hour before bedtime for insomnia; repeat at bedtime, then another in an hour if necessary.	
<b>SULPHONAL &amp; HYOSCINE</b> ... { Sulphon. ....5 gr. }	37 50
Sedative and hypnotic. { Hyosc. Hydrobr.1-200 gr. }	
Per 100.....	3 75
One, three times a day, or every three hours at night, if necessary, as a sedative and hypnotic.	
<b>SULPHUR</b> .....1-100 gr.....	75
One every two hours, as a laxative for an infant, and in the chronic stage of pertussis.	
<b>SULPHUR</b> .....1-50 gr.....	75
A tablet three or four times a day, with local applications, for subacute and chronic skin diseases of children.	
<b>SULPHUR</b> .....1-10 gr.....	75
One every three hours, for persistent bronchitis after whooping cough.	
<b>SULPHUR</b> .....1-2 gr.....	75
One every three hours for chronic catarrh, rheumatism and skin diseases.	

<b>SULPHUR</b> .....	1 gr .....	75
One or two after meals, for acne, pityriasis, and other skin diseases, and as a laxative for children.		
<b>SULPHUR COMP.</b> .....	{ Sulphur.....5 gr. } { Cream Tartar.....1 gr. }	1 00
One or two at night, as a mild stimulating laxative.		
<b>SULPHUR COMP. No. 2</b> .....	{ Sulphur.....2 1-2 gr. } { Cream Tartar...2 1-2 gr. }	1 00
One after meals and two at night, for hæmorrhoids, and as a laxative in pregnancy.		
<b>SULPHUR COMP. No. 3</b> .....	{ Sulphur.....5 gr. } { Cream Tartar.....2 gr. } { Ext. Ipecac.....1-100 gr. } { Ext. Capsicum..1-500 gr. } { Acid Arsen...1-1000 gr. } { Calcium Sulph...1-8 gr. }	1 50
	<b>Chocolate Coated</b> .....	1 60
Two to four a day, for chronic eczema, pityriasis, acne, and other obstinate cutaneous diseases.		
<b>SULPHUR &amp; IPECAC</b> .....	{ Sulphur.....1-10 gr. } (Dr. Gilbert.) { Ipecac Powd....1-10 gr. }	75
One every three hours, in gastric derangements and as an aid to digestion.		
<b>SUMBUL COMP.</b> .....	{ Iron Sulph. Dry.....1 gr. } (Dr. Goodell.) { Ext. Sumbul.....1 gr. } { Asafoetida.....2 gr. } { Acid Arsenous...1-40 gr. }	3 00
	<b>Chocolate Coated only</b>	
One after each meal as a general nerve tonic, and tonic in anæmia.		
<b>SUN CHOLERA TABLETS</b> ...	{ Tr. Opii.....3 min. } { Tr. Rhei.....3 min. } { Tr. Capsici.....3 min. } { Spts. Camphoræ...3 min. } { Spts. Menth. Pip...3 min. }	1 60
Each tablet represents 15 minims of Sun Cholera Mixture.	<b>Chocolate Coated</b> .....	1 70
Give two tablets at the first dose; then one after each loose movement, for six or eight doses if necessary.		
<b>SUN CHOLERA TABLETS</b> ...	{ Tr. Opii....2 min. } { Tr. Rhei....2 min. } { Tr. Capsici.....2 min. } { Spts. Camphoræ...2 min. } { Spts. Menth. Pip...2 min. }	1 35
<b>No. 2</b> .....	<b>Chocolate Coated</b> .....	1 45
Each tablet represents 10 minims of Sun Cholera Mixture.		
Give two tablets at the first dose; then one after each loose movement, for six or eight doses if necessary.		
<b>SYPHILITIC</b> .....	{ Potass. Iodide...2 1-2 gr. } { Corrosive Sublim.1-40 gr. }	3 00
A tablet three times a day after meals, increasing according to the necessities of the case.		
<b>SYR. SQUILLS COMP.</b> .....	5 min.....	1 50
One every three hours, or oftener, for winter cough, asthma, pneumonia and phthisis.		
<b>TARAXACUM, PODOPHYLL- IN COMP.</b> .....	{ Ext. Taraxacum...1-6 gr. } { Podophyllin.....1-8 gr. } { Ext. Nux Vom...1-10 gr. } (Dr. Barr.) { Ipecac Powd.....1-4 gr. }	1 35
One tablet after meals, as a stomachic.		

---

When ordering or prescribing



<b>THROAT</b> .....	{ Potass. Iodide...1-10 gr. }	1 10
	{ Salt.....1-10 gr. }	
	{ Sugar.....1-2 gr. }	
Allow a tablet to dissolve in the throat every half hour, for hoarseness, dryness and tickling.		

<b>THROAT TABLETS</b> ..... (Dr. F. J. Quinlan.)	{ Acid Benzoic.....1-5 gr. }	1 50
	{ Tr. Belladonna....1 min. }	
	{ Paregoric.....10 min. }	
	{ Ext. Licorice.....q. s. }	

Dissolve one slowly on the tongue, not oftener than every four hours, for dryness of the throat attended with pain. Used also for the relief of acute and subacute inflammation of the fauces.

<b>TINCT. CINCHONA COMP.</b> .....10 minims.....	1 75
--	------

One tablet for a child, three times a day, as a tonic and febrifuge.

<b>TINCT. CINCHONA COMP.</b> ....1 teaspoonful.....	2 25
---	------

One tablet three or four times a day, for tonic.

<b>TINCT. THUJA</b> .....5 minims.....	1 50
--	------

One tablet every three hours, for cystitis.

<b>TONIC</b> ..... (Dr. Hammond.)	{ Iron Pyrophos....1-2 gr. }	1 50
	{ Quinine Sulph....1-2 gr. }	
	{ Strych. Sulph....1-120 gr. }	
	<b>Chocolate Coated</b> .....	1 60

One or two tablets just after each meal, as a nerve or general tonic in anæmia.

<b>TONIC No. 3</b> ..... (Dr. Hoopes.)	{ Ferri Pyrophos.....2 gr. }	1 75
	{ Quinine Sulph.....1-4 gr. }	
	{ Strych. Sulph....1-60 gr. }	
	<b>Chocolate Coated</b> .....	1 85

A tablet three times a day after wasting disease, and as an hæmætic in anæmia and malarial poisoning.

<b>TONIC ALTERNATIVE</b> .....	{ Corrosive Sub....1-50 gr. }	75
	{ Strych. Sulph....1-100 gr. }	
	{ Ipecac Powd....1-20 gr. }	
	<b>Chocolate Coated</b> .....	85

One or two after meals, as a tonic and alternative.

<b>TONIC ALTERNATIVE No. 2</b> .... (Dr. E. Jay Fisk.)	{ Corrosive Sub....1-30 gr. }	1 10
	{ Strych. Sulph....1-60 gr. }	
	{ Iron Powd.....1 gr. }	
	<b>Chocolate Coated</b> .....	1 20

One or two after meals for anæmic and cachectic conditions, chronic bronchitis and malaria.

<b>TONIC CHALYBEATE</b> ..... (Dr. Flint.)	<b>Plain or Chocolate Coated</b>	2 00
	1-2 size.....	1 50

From an analysis of the blood, containing an excess of iron and chloride of sodium.

Two tablets, three times a day after eating. A tonic made from an analysis of the blood; useful in anæmia, Bright's disease and nervous exhaustion.

<b>TONIC CHLORIDES</b> .....	{ Iron Chloride.....1 gr. }	1 25
	{ Merc. Bichlor....1-48 gr. }	
	{ Arsenic Chlor....1-64 gr. }	

One after meals for anæmia, and as a general tonic in convalescence.

<b>TONIC, GENERAL</b> ..... (Dr. Frank E. West.)	{ Acid Arsenous....1-30 gr. }	3 00
	{ Quinine Sulph....1-2 gr. }	
	{ Strych. Sulph....1-60 gr. }	
	{ Salicin.....1 1-2 gr. }	
	<b>Plain or Chocolate Coated</b>	

One tablet after each meal as a nerve and general tonic.

TONIC LAXATIVE (Dr. Miller.)	Ext. Colocy. Co...1-2 gr.	2 80
	Ext. Belladonna. 1-10 gr.	
	Quinine Sulph....1 gr.	

One to three at night, for chronic atony of the bowels and constipation of malarial and anæmic subjects.

TONIC SYPHILITIC (Dr. Feltwell.)	Mercury Protiod...1-5 gr.	1 50
	Morph. Sulph....1-24 gr.	
	Tartar Emetic....1-24 gr.	

TONIC TREATMENT.....	Capsicum.....1-60 gr.	1 25
	Opium Powd....1-15 gr.	
	Mercury Protiod..1-4 gr.	

One or two at a dose, two to three times a day.

TONSILLITIS .....	Tr. Aconite.....1-5 min.	1 00
	Tr. Bryonia....1-10 min.	
	Tr. Belladonna. 1-10 min.	
	Merc. Iod. Red...1-100 gr.	

Dissolve one on the tongue every twenty minutes for first hour, then every thirty minutes for the second hour, after which, one every hour, avoiding all acids or fruit; for sore throat from cold, also in mild cases of diphtheria, and scarlatinal sore throat.

TONSILLITIS, ACUTE (Dr. Newman.)	Opium Denarc. .1-10 gr.	1 25
	Tr. Verat. Viride .1-4 min.	
	Calomel.....1-10 gr.	
	Sugar Milk.....q. s.	
	Oil Anise .....1-20 min.	

One every hour for adults or children.

TONSILLITIS No. 3 (Dr. Bierhoff.)	Corros. Sublim...1-96 gr.	1 50
	Tr. Chloride Iron..3 min.	
	Oil Cinnamon.....q. s.	

One dissolved in water every hour, for diphtheria or tonsillitis.

TONSILLITIS No. 4 (Dr. Koyle.)	Merc. Protiod....1-50 gr.	1 00
	Licorice Powd .....q. s.	

One every fifteen minutes for twelve hours, in acute follicular tonsillitis, also as a disinfectant in diphtheria.

TONSILLO PHARYNGITIS.... (Dr. J. C. Applegate.)	Tr. Verat. Vir. .1-2 min.	1 35
	Tr. Belladonna. 1-2 min.	
	Tartar Emetic....1-50 gr.	
	Morphine Sulph..1-50 gr.	
	Merc. Iod. Red..1-100 gr.	

Two tablets every hour until improved, then one or two every two hours; in all inflammations of the tonsils, pharynx and larynx.

TRIFOLIUM ALTERNATIVE...	Ext. Trifolium....1-2 gr.	1 75
	Ext. Stillingia....1-2 gr.	
	Ext. P'kly Ash B'k.1-8 gr.	
	Ext. Burdock.....1-8 gr.	
	Ext. Poke Root...1-2 gr.	
	Ext. Blue Flag....1-4 gr.	

Two tablets, three times a day, as an alterative and anti-syphilitic.

TRIONAL.....	5 grs.....	30 00
--------------	------------	-------

One every hour, for three doses in insomnia and delirium, and to replace morphine where it disagrees, or in the cure of the habit.

TRIPLE BROMIDES No. 1. Sedative.	Sodium Brom....2 1-2 gr.	1 35
	Potass. Brom...2 1-2 gr.	
	Ammon. Brom..2 1-2 gr.	

One to four tablets dissolved in water, every three hours until the patient is quiet.

<b>TRIPLE BROMIDES No. 2</b> .....	{ Sodium Bromide....5 gr. Potass. Bromide....5 gr. Ammon. Bromide....5 gr. }	2 25
------------------------------------	--	------

One or two dissolved in water, three times a day, as a nerve sedative.

<b>TRIPLE SULPHOCARBONATES</b> .....	{ Zinc Sulphocarb. 1 1-2 gr. Sod. Sulphocarb. 1 1-2 gr. Cal. Sulphocarb. 1 1-2 gr. }	2 00
--------------------------------------	--	------

One tablet after meals, for catarrh of the bowels, and as an intestinal antiseptic.

<b>TRIPLEX</b> .....	{ Aloes.....2 gr. Pil. Hydrarg.....1 gr. Podophyllin.....1-4 gr. }	1 75
	<b>Chocolate Coated</b> .....	1 85

One for a laxative, two for a cathartic, three for a purge; best taken just before retiring, for biliousness and constipation.

<b>TULLY'S COMP. No. 2</b> .....	{ Tully's Powder. 2 1-2 gr. Podophyllin.... 1-10 gr. }	1 75
----------------------------------	---	------

A tablet every two or three hours, for fever, pain and cough, and to induce sleep.

<b>TULLY'S POWDER</b> .....	1-5 gr.....	75
-----------------------------	-------------	----

A tablet every two hours for painful diarrhoea of children.

<b>TULLY'S POWDER</b> .....	1-2 gr.....	75
-----------------------------	-------------	----

One, repeated in an hour, for painful cough of children.

<b>TULLY'S POWDER</b> .....	1 gr.....	90
-----------------------------	-----------	----

One every hour for three or four doses, as a diaphoretic.

<b>TULLY'S POWDER</b> .....	5 grs.....	2 25
	<b>Plain or Chocolate Coated</b> .....	

One tablet repeated in an hour and a half, as a sedative and diaphoretic.

<b>TULLY'S POWDER</b> .....	10 grs.....	3 25
	<b>Plain or Chocolate Coated</b> .....	

One at night, for insomnia due to pain, and for delirium and coma vigil.

<b>TULLY'S TABLETS</b> .....	2 1-2 grs.....	1 50
	<b>Chocolate Coated</b> .....	1 60

A tablet as often as every two hours, as a sedative and anodyne.

<b>TURPETH MINERAL</b> .....	1 gr.....	75
------------------------------	-----------	----

Emetic.

One every half hour for three doses, to produce vomiting; aided by warm water.

<b>TURPETH MINERAL</b> .....	2 grs.....	1 15
------------------------------	------------	------

A tablet, repeated in half an hour, to empty an overloaded stomach and for croup.

<b>TYPHOID FEVER No. 1</b> .....	{ Podoph. Resin.. 1-960 gr. Merc. Chlor. Mild. 1-16 gr. Guaiacol Carb... 1-16 gr. Menthol... 1-16 gr. Eucalyptol..... q. s. }	1 50
----------------------------------	---	------

One every fifteen minutes during the first forty-eight hours, while awake; if bowels move too freely decrease the dose. At the end of twenty-four hours begin with the Typhoid Fever Tablets No. 2.

<b>TYPHOID FEVER No. 2</b> .....	{ Podoph. Resin.. 1-960 gr. Merc. Chlor. Mild. 1-16 gr. Guaiacol Carb... 1-4 gr. Menthol... 1-16 gr. Thymol.... 1-16 gr. Eucalyptol..... q. s. }	3 25
----------------------------------	---	------

One or more tablets every fifteen minutes while awake, either in connection with the Typhoid Fever Tablets No. 1, or alone, until bowels move freely, then decrease the dose until the movements become reduced to one or two a day, and the temperature has touched the normal point.

<b>UTERINE SEDATIVE</b> ..... (Dr. R. G. H. Hayes.)	{ Helonin.....1-8 gr. }	<b>Plain or Chocolate Coated</b> 2 50
	{ Viburnin.....1-8 gr. }	
	{ Caulophyllin.....1-4 gr. }	
	{ Mitchella Rep...1 1-2 gr. }	

One to three tablets every three hours, for membranous or neuralgic dysmenorrhœa, and as a uterine tonic or antispasmodic.

<b>UTERINE SEDATIVE No 2</b> ....	{ Ext. Viburnum Opul.1 gr. }	<b>Chocolate Coated only</b> 2 50
	{ Ext. Hydras. Can.3-4 gr. }	
	{ Ext. Jam. Dogwood.1-2 gr. }	
	{ Ext. Pulsatilla....1-4 gr. }	

One to three tablets every three hours, for membranous or neuralgic dysmenorrhœa, and as a uterine tonic or antispasmodic.

<b>UTERINE TONIC</b> .....	{ Ergotin.....1-4 gr. }	1 30
	{ Hydrastin.....1-100 gr. }	
	<b>Chocolate Coated</b> .....	1 40

One to be given three or four times a day, in menorrhagia, metorrhagia and as a general uterine tonic.

<b>UTERINE TONIC No. 2</b> .....	{ Helonin.....1-6 gr. }	<b>Chocolate Coated only</b> 4 25
	{ Caulophyllin.....1-6 gr. }	
	{ Macrotin.....1-6 gr. }	
	{ Hyoscy. A morph.1-250 gr. }	

Indicated in functional pelvic derangements, accompanied by pain and melancholia.

<b>URETHRITIS No. 1</b> ..... For 2 oz. ( $\frac{1}{8}$ pint) bottle.	{ Zinc Acetate.....2 gr. }	1 50
	{ Corrosive Sub.....1-5 gr. }	
	Per doz. bottles net.....	2 25

Dissolve a tablet in four tablespoonfuls of hot water and use as an injection as hot as can be borne, after each urination.

<b>URETHRITIS No. 2</b> ..... For 4 oz. ( $\frac{1}{4}$ pint) bottle.	{ Zinc Acetate.....4 gr. }	1 75
	{ Corros. Sub.....2-5 gr. }	

Dissolve a tablet in eight tablespoonfuls of hot water and use as an injection as hot as can be borne, after each urination.

<b>VAGINAL ASTRINGENT</b> .....	{ Alum.....3 gr. }	2 00
	{ Zinc Sulph.....2 gr. }	
	{ Acid Tannic.....2 gr. }	
	{ Acid Boracic.....6 gr. }	
	{ Ext. Hydrastis.....1 gr. }	
	(Colorless)	

Dissolve two tablets in a quart of water and use as an injection.

<b>VAGINAL WAFER</b> .....	{ Zinc Sulph.....3 3-4 gr. }	1 75
	{ Sodium Sulph...3 1-2 gr. }	
	{ Sodium Borate....4 gr. }	
	{ Acid Boric.....3-4 gr. }	

Insert one every third night, followed by irrigation, for leucorrhœa.

<b>VALERIANATE SEDATIVE</b> ....	{ Quin. Valerianate..1 gr. }	<b>Chocolate Coated only</b> 4 25
	{ Iron Valerianate...1 gr. }	
	{ Zinc Valerianate...1 gr. }	
	{ Ext. Cannabis Ind.1-4 gr. }	

One tablet every four hours, as a nerve sedative and tonic.

<b>VERATRIA SULPHATE</b> .....	1-20 gr.....	1 30
--------------------------------	--------------	------

A tablet every four hours, for pertussis, and neuralgia.

<b>VERATRIA SULPHATE</b> .....	1-10 gr.....	2 25
--------------------------------	--------------	------

One every three or four hours, for gout, rheumatism, neuralgia and paralysis agitans.

<b>VERATRUM COMP.</b> .....	{ Antifebrin.....1 gr. Tr. Verat. Viride. 1-2 min. Wine Ipecac ....1-2 min. Codeine.....1-24 gr. Sugar Milk.....q. s. }	2 50
(Dr. F. R. Newman.)		

One every hour until six are taken; then one every two hours for fever.

<b>VIBURNUM SEDATIVE</b> .....	{ Ext. Viburn. Prun...1 gr. Ext. Viburnum Op...1 gr. Ext. Aletris Far. .1-2 gr. Ext. Helon. Dioica.1-2 gr. Ext. Mitchel. Rep.1-2 gr. Caulophyllin.....1-4 gr. }	2 75
	<b>Plain or Chocolate Coated</b>	

One to three tablets every three hours, or one tablet every hour, with a swallow of hot water or tea, as a uterine tonic and antispasmodic.

<b>VIBURNUM SEDATIVE</b> .....	{ Ext. Viburn. Prun..1-2 gr. Ext. Viburn. Op...1-2 gr. Ext. Aletris Far. .1-4 gr. Ext. Helon. Dioica.1-4 gr. Ext. Mitchel. Rep..1-4 gr. Caulophyllin.....1-8 gr. }	2 00
<b>One-half strength.</b>	<b>Chocolate Coated only</b>	

One to three tablets every two or three hours, as a uterine tonic and antispasmodic.

<b>VIBURNUM SEDATIVE No. 2.</b>	{ Fl. Ex. Viburn. Op.1 min. Camphor.....1-2 gr. Opium Powd.....1-4 gr. Oil Anise.....1-8 min. }	1 60
	<b>Plain or Chocolate Coated</b>	1 70

Begin one day before the period, with a tablet night and morning; the next day take three, and the next four; for scanty and painful menstruation and ovarian colic.

<b>VIBURNUM SEDATIVE No. 3.</b>	{ Black Haw.....3 3-4 gr. Golden Seal.....3 3-4 gr. Jam. Dogwood. .1 3-4 gr. }	2 00
(Dr. Fisher.)		
<b>Each tablet represents:</b>	<b>Plain or Chocolate Coated</b>	

One after meals for the week before the catamenia, increase if necessary during the flow; for dysmenorrhœa, menorrhagia, and hyperæmia of the pelvic organs, and as an antispasmodic.

<b>VIBURNUM SEDATIVE No. 4.</b>	{ Ext. Black Haw....2 gr. Ext. Hyoscyam...1-6 gr. Camphor Powd...1-4 gr. Morphine Sulph..1-6 gr. }	4 50
(Dr. Frank Hollister.)	<b>Plain or Chocolate Coated</b>	

One every four hours, commencing the day before the period is expected and continued until normal action is established; used in cases of dysmenorrhœa.

<b>VIBURNUM SEDATIVE</b> .....	{ Ext. Viburn. Prun..1 gr. Ext. Viburnum Op..1 gr. Ext. Aletris Far. .1-2 gr. Ext. Helon. Dioica.1-2 gr. Ext. Mitchel. Rep.1-2 gr. Caulophyllin.....1-4 gr. Ext. Pulsatilla.....1 gr. }	3 50
<b>Modified.</b>	<b>Chocolate Coated only</b>	

One or two tablets every two or three hours, as a uterine tonic and antispasmodic.

<b>WARBURG'S TINCTURE</b> .....1 teaspoonful.....		
<b>Chocolate Coated only</b>	4	00
One to two tablets three or four times a day, for chronic malarial fevers; if bowels move too freely, decrease the dose, or alternate with the Warburg Tincture Tablet without aloes.		
<b>WARBURG'S TINCTURE</b> .....1 teaspoonful.....		
<b>Without Aloes. Chocolate Coated only</b>	4	00
One to four, three or four times a day, in chronic malarial poisoning; if bowels are constipated, give the Warburg Tincture Tablet with aloes until laxative action is obtained.		
<b>WHOOPING COUGH</b> .....	{ Antipyrine.....1-2 gr. } { Tr. Drosera.....1 min. } { Belladonna.....1-500 gr. } { Ipecac Powd.....1-200 gr. }	1 65
One tablet every hour, for whooping cough.		
<b>WHOOPING COUGH No. 2</b> .....	{ Acetanilid.....1-2 gr. } { Tr. Drosera.....1 min. } { Belladonna.....1-400 gr. } { Ipecac Powd.....1-200 gr. }	1 50
One tablet every hour, for whooping cough.		
<b>WHOOPING COUGH No. 3</b> .....	{ Tr. Drosera.....2 min. } { (Dr. J. D. McGaughey.) Fl. Ext. Bellad. 1-12 min. }	1 50
For children over one year, one tablet every two hours for whooping cough.		
<b>WINTER COUGH</b> .....	{ Terpin Hydrate.....2 gr. } { Ammon. Muriate.....1 gr. } { Ext. Licorice.....1-2 gr. } { Ipecac Powd.....1-10 gr. }	1 50
One every two hours, for persistent cough.		
<b>YELLOW TABLETS</b> (Sugar Milk).....		50
<b>ZINC ACETATE</b> .....1 gr.....		75
One dissolved in one ounce of water, as an injection.		
<b>ZINC COMP.</b> .....	{ Lead Acetate.....2 gr. } { Zinc Acetate.....2 gr. }	1 50
Two dissolved in four ounces of water, for injection.		
<b>ZINC PHOSPHIDE</b> .....1-100 gr.....		75
Nerve tonic. One after meals, increasing gradually, for chorea and other spasmodic diseases.		
<b>ZINC PHOSPHIDE</b> .....1-50 gr.....		75
Three or four a day, increasing slowly, for chorea, epilepsy and neuralgia.		
<b>ZINC PHOSPHIDE</b> .....1-25 gr.....		75
A tablet every four hours, for hysteria and spasmodic affections of the muscles.		
<b>ZINC PHOSPHIDE</b> .....1-10 gr.....		75
One three times a day, for chronic nervous disease of an ataxic nature, and for neuralgia.		
<b>ZINC PHOSPHIDE</b> .....1-4 gr.....		1 10
One after meals, as a tonic to the cerebro-spinal centres, in neurasthenia, hysteria and epilepsy.		
<b>ZINC PHOS. &amp; NUX VOMICA</b> {	{ Zinc Phos.....1-10 gr. } { (Dr. Hammond.) Ext. Nux Vomica.1-4 gr. }	1 10
Nerve tonic. <b>Chocolate Coated</b> .....		1 20
One every three or four hours, as a general tonic to the nervous system.		

Please specify Fraser's Tablets

<b>ZINC SULPHATE</b> .....1 gr.....	75
Dissolve one in a tablespoonful of water; the resulting solution has a strength of about 1 to 250. For solution.	
<b>ZINC SULPHOCARBOLATE</b> ...1 gr.....	1 10.
One every hour, as an antiseptic and antipyretic in diarrhoea and dysentery of infants; dissolve it in a teaspoonful of hot water.	
<b>ZINC SULPHOCARBOLATE</b> ...2 grs.....	1 50
One every two hours in a little hot water, as an intestinal antiseptic in enteritis.	
<b>ZINC SULPHOCARBOLATE</b> ...2 1-2 grs.....	1 50
A tablet every two or three hours, for gastro enteritis with fever, and for dysentery.	
<b>ZINC SULPHOCARBOLATE</b> ...5 grs.....	2 00
One every three hours, as an antipyretic and intestinal disinfectant in typhoid fever and enteritis.	

## FRASER'S LOZENGES

<b>AMMONIATED LICORICE</b> ....	{ Ammonia Mur.....1 gr. Ext. Licorice....3 3-4 gr. Oil Anise.....1-20 min.	1 50
One or two dissolved in the mouth every hour, or as occasion requires, for hard dry cough with hoarseness and scanty expectoration.		
<b>ANTISEPTIC MOUTH AND THROAT LOZENGES</b> .....	{ Acid Carbolic.....1-40 gr. Menthol.....1-30 gr. Oil Sassafras.....q. s. Acacia.....q. s. Sugar.....q. s.	1 50
One dissolved in mouth as required, for sore throat or mouth; useful in correcting offensive breath, due to decayed teeth or disordered stomach.		
<b>ASTRINGENT THROAT</b> .....	{ Acid Benzoic.....1-5 gr. Opium Powd.....1-50 gr. Catechu Powd...1-4 gr. Camphor Powd...1-20 gr. Ext. Yerba Santa..1-4 gr. Sugar.....q. s. Aromatics.....q. s.	1 50
One allowed to dissolve on the tongue every half hour to hour, for sore and inflamed throat with cough, where a mild astringent is desired.		
<b>BROWN MIXTURE</b> .....	{ Acid Benzoic.....1-20 gr. Opium Powd.....1-20 gr. Camphor Powd...1-20 gr. Tartar Emetic....1-40 gr. Oil Anise.....q. s. Licorice.....q. s. Sugar.....q. s.	1 50
One on the tongue every hour for bronchial cough. Stop when oppression of chest is relieved and secretion well established.		
<b>BROWN MIXTURE COMP</b> ....	{ Acid Benzoic.....1-20 gr. Opium Powd.....1-20 gr. Camphor Powd...1-20 gr. Tartar Emetic....1-40 gr. Ammon. Muriate...1 gr. Oil Anise.....q. s. Licorice.....q. s. Sugar.....q. s.	1 50
One on the tongue, for bronchial cough with scanty expectoration and hoarseness. Reduce dose as oppression of chest is relieved.		

When ordering or prescribing

COMPOUND AMMONIA.....	{ Potass Chlorate...1-4 gr. Ammon. Muriate...1-4 gr. Cubebs Powd.....1-8 gr. Sugar.....q. s. Aromatics.....q. s.	1 50
-----------------------	--	------

One allowed to dissolve in the mouth every fifteen to thirty minutes as occasion requires, to relieve tickling and irritation of throat.

COUGH ACUTE.....	{ Terpin Hydrate...2 gr. Codeine.....1-8 gr.	4 00
------------------	---	------

One every hour or two in the first stages of cough, with scanty and painful expectoration.

CUBEBS COMP. with Cocaine..	{ Cocaine Mur.....1-50 gr. Acid Benzoic.....1-10 gr. Oleoresin Cubebs, 1-10 gr. Potass. Chlorate...1-2 gr. Sugar.....q. s. Ext. Licorice.....q. s.	1 50
-----------------------------	---	------

One dissolved in the mouth every half hour for four doses, then every hour or two as needed, to relieve tickling of the throat and bronchial irritation and cough.

CUBEBS CO. with Codeine.....	{ Ammonium Mur...1-4 gr. Oleoresin Cubebs, 1-10 gr. Codeine.....1-25 gr. Ext. Licorice.....q. s. Sugar.....q. s.	1 85
------------------------------	--	------

One on the tongue every fifteen or thirty minutes, as a sedative and expectorant, allaying tickling of the throat and relieving bronchial irritation and hoarseness.

MENTHOLIC THROAT. ....	{ Menthol.....1-35 gr. Cocaine Mur...1-250 gr. Oil Anise.....1-80 min. Acid Benzoic.....1-12 gr. Eucalyptol.....1-15 gr. Sugar.....q. s.	1 50
------------------------	---	------

To be used as a voice or throat tablet by allowing one to dissolve on the tongue and repeat in fifteen minutes to half an hour if necessary.

PALMETTO COMP.....	{ Ext. Saw Palmetto...1 gr. Ext. Yerba Santa...1 gr. Oil Wintergreen...q. s. Sugar.....q. s.	2 25
--------------------	---	------

Dissolve one in the mouth every three hours, for phthisical cough.

PEPSIN CHARCOAL COMP....	{ Pepsin Sacch.....1 gr. Charcoal.....1-2 gr. Magnesia.....1 gr. Ginger.....1-2 gr.	1 50
--------------------------	--	------

One, half an hour before meals to stimulate the appetite; one or two after meals repeated at intervals of an hour, to correct acid stomach and to relieve indigestion accompanied with flatulence and fermentation.

RED GUM COMP.....	{ Menthol.....1-20 gr. Cocaine Mur...1-20 gr. Red Gum.....q. s.	2 00
-------------------	---	------

One every hour for two doses, then every two to three hours, to allay painful symptoms accompanying sore throat with cough.

RED GUM LOZENGES.....		1 50
-----------------------	--	------

One allowed to dissolve in the mouth as occasion requires.

THROAT SEDATIVE.....	{ Morph. Hydroch.1-50 gr. Cocaine Hydroch.1-50 gr. Tr. Aconite.....1 min.	1 50
----------------------	---	------

One every hour during the day for an adult, for quinsy sore throat, and to relieve inflammation in throat troubles.

# FRASER'S

## Tablets from Official Tinctures.

		Per Thousand.
ACONITE.....	{ 1-4 minim, 1-2 minim, { 3-4 minim, 1 minim... }	75
ACONITE.....	2 minims.....	1 00
BELLADONNA.....	1-2 minim, 1 minim....	75
BELLADONNA.....	2 minims.....	1 00
BRYONIA.....	1-2 minim, 1 minim....	75
BRYONIA.....	2 minims.....	1 00
CANTHARIDES.....	1-2 minim, 1 minim....	75
CANTHARIDES.....	2 minims.....	1 00
CANNABIS INDICA.....	1 minim.....	75
CANNABIS INDICA.....	2 minims.....	1 10
CATECHU.....	1 minim.....	75
CATECHU.....	2 minims.....	1 00
CALABAR BEAN.....	1 minim.....	75
CALABAR BEAN.....	2 minims.....	1 10
CIMICIFUGA.....	1-2 minim, 1 minim....	75
CIMICIFUGA.....	2 minims.....	1 00
CINCHONA.....	1 minim.....	75
CINCHONA.....	2 minims.....	1 00
COLCHICUM.....	1 minim.....	75
COLCHICUM.....	2 minims.....	1 00
CONIUM.....	1 minim.....	75
CONIUM.....	2 minims.....	1 00
CUBEBS.....	1-2 minim, 1 minim....	75
CUBEBS.....	2 minims.....	1 00
DIGITALIS.....	1-2 minim, 1 minim....	75
DIGITALIS.....	2 minims.....	1 00
DIGITALIS.....	5 minims.....	1 00
GELSEMIUM.....	1 minim.....	75
GELSEMIUM.....	2 minims.....	1 00
GUAIAAC.....	1 minim.....	75
GUAIAAC.....	2 minims.....	1 00
HYDRASTIS.....	1 minim.....	75
HYDRASTIS.....	2 minims.....	1 00
HYOSCYAMUS.....	1 minim.....	75
HYOSCYAMUS.....	2 minims.....	1 00
HAMAMELIS.....	1 minim.....	75
HAMAMELIS.....	2 minims.....	1 00
IGNATIA.....	1-2 minim, 1 minim....	75
IGNATIA.....	2 minims.....	1 00
IPECAC.....	1-2 minim, 1 minim....	75

When ordering or prescribing

IPECAC.....	2 minims.....	1 00
KRAMERIA. ....	1 minim.....	75
KRAMERIA.....	2 minims.....	1 00
NUX VOMICA.....	1-2 minim, 1 minim....	75
NUX VOMICA.....	2 minims.....	1 00
NUX VOMICA .....	5 minims.....	1 10
OPIUM DEOD .....	1 minim.....	1 35
OPIUM DEOD.....	2 minims.....	1 85
SANGUINARIA. ....	1 minim.....	75
SANGUINARIA.....	2 minims.....	1 00
SQUILL.....	1 minim.....	75
SQUILL.....	2 minims.....	1 00
STRAMONIUM.....	1 minim.....	75
STRAMONIUM.....	2 minims.....	1 00
STROPHANTHUS.....	1 minim.....	1 00
STROPHANTHUS.....	2 minims.....	1 25
STROPHANTHUS.....	5 minims.....	1 75
VERATRUM VIRIDE.....	1 minim.....	75
VERATRUM VIRIDE.....	2 minims.....	1 00
VERATRUM VIR. (Norwood's).	1 minim.....	1 50
VERATRUM VIR. (Norwood's).	2 minims.....	3 00

## Tablets from Mother Tinctures.

ACONITE. ....	} 1-4 min., 1-2 min., 3-4 min. }	75
ACONITE.....		
ACTÆA RACEMOSA.....	2 minims.....	1 10
ACTÆA RACEMOSA.....	1-2 minim.....	75
ACTÆA RACEMOSA.....	1 minim.....	1 00
ACTÆA RACEMOSA.....	2 minims.....	1 10
AILANTHUS.....	1 minim.....	1 30
ALOES.....	1 minim.....	75
APIS.....	1-2 minim, 1 minim....	75
APOCYNUM.....	1 minim.....	75
APOCYNUM.....	2 minims.....	1 10
ARNICA.....	1 minim.....	1 00
ASCLEPIAS TUB .....	1 minim.....	75
ASCLEPIAS TUB .....	2 minims.....	1 10
BAPTISIA.....	1-2 minim, 1 minim....	75
BAPTISIA.....	2 minims.....	1 10
BELLADONNA.....	1-2 minim, 1 minim....	75
BELLADONNA.....	2 minims.....	1 10
BRYONIA.....	1-2 minim, 1 minim....	75
BRYONIA.....	2 minims.....	1 10
CACTUS.....	1 minim.....	1 10

Please specify Fraser's Tablets

CALABAR.....	1-2 minim.....	75
CALABAR.....	1 minim.....	1 00
CALABAR.....	2 minims.....	1 10
CALENDULA.....	1 minim.....	75
CANNABIS INDICA.....	1 minim.....	75
CANNABIS SATIVA.....	1 minim.....	75
CANTHARIDES.....	1-2 minim, 1 minim....	75
CANTHARIDES.....	2 minims.....	1 00
CAPSICUM.....	1-2 minim, 1 minim....	75
CAPSICUM.....	2 minims.....	1 00
CAUSTICUM.....	1-2 minim.....	75
CAUSTICUM.....	1 minim.....	1 00
CAUSTICUM.....	2 minims.....	1 10
CHAMOMILLA.....	1 minim.....	75
CHAMOMILLA.....	2 minims.....	1 10
CHELIDONIUM.....	1 minim.....	75
CHIMAPHILLA.....	1 minim.....	1 00
CHINA.....	1 minim.....	75
CHINA.....	2 minims.....	1 00
CIMICIFUGA.....	1-2 minim, 1 minim....	75
CIMICIFUGA.....	2 minims.....	1 10
CISTUS CAN.....	1 minim.....	75
CLEMATIS.....	1 minim.....	75
COCCULUS.....	1-2 minim, 1 minim....	75
COCCULUS.....	2 minims.....	1 10
COFFEA.....	1 minim.....	75
COFFEA.....	2 minims.....	1 10
COLCHICUM.....	1-2 minim, 1 minim....	75
COLCHICUM.....	2 minims.....	1 10
COLLINSONIA.....	1-2 minim, 1 minim....	75
COLLINSONIA.....	2 minims.....	1 10
COLOCYNTHIS.....	1-2 minim, 1 minim....	75
COLOCYNTHIS.....	2 minims.....	1 10
CONIUM.....	1 minim.....	75
CONIUM.....	2 minims.....	1 10
CORYDALIS.....	1 minim.....	75
CUBEBS.....	1-2 minim, 1 minim....	75
CUBEBS.....	2 minims.....	1 00
DIGITALIS.....	1-2 minim, 1 minim....	75
DIGITALIS.....	2 minims.....	1 10
DIOSCOREA.....	1-2 minim, 1 minim....	75
DIOSCOREA.....	2 minims.....	1 10
DROSERA.....	1-2 minim, 1 minim....	75
DROSERA.....	2 minims.....	1 10
DULCAMARA.....	1 minim.....	75
EUPHORBIA.....	1 minim.....	75
EUPHRASIA.....	1 minim.....	75

When ordering or prescribing

EUONYMUS	1 minim	75
GELSEMIUM	1-2 minim, 1 minim	75
GELSEMIUM	2 minims	1 15
GUAIAC	1 minim	75
HAMAMELIS	1 minim	75
HAMAMELIS	2 minims	1 00
HYDRASTIS	1 minim	75
HYDRASTIS	2 minims	1 00
HYOSCYAMUS	1 minim	75
HYOSCYAMUS	2 minims	1 10
IGNATIA	1-2 minim, 1 minim	75
IGNATIA	2 minims	1 00
IPECAC	1-2 minim, 1 minim	75
IPECAC	2 minims	1 00
IRIS	1 minim	75
KALMIA	1 minim	75
LEPTANDRA	1 minim	75
LOBELIA	1 minim	75
NUX VOMICA	1-2 minim, 1 minim	75
NUX VOMICA	2 minims	1 00
OPIUM	1-2 minim	1 00
OPIUM	1 minim	1 30
OPIUM	2 minims	1 85
PHOSPHORUS	1 minim	75
PHOSPHORUS	2 minims	1 00
PHYTOLACCA	1 minim	75
PODOPHYLLUM	1 minim	75
PODOPHYLLUM	2 minims	1 15
PULSATILLA	1-2 minim, 1 minim	75
PULSATILLA	2 minims	1 15
RHUS TOX	1 minim	75
RHUS TOX	2 minims	1 15
RUMEX	1 minim	75
SAMBUCUS	1 minim	75
SENECIO	1 minim	75
SPIGELIA	1 minim	75
SPONGIA	1-2 minim, 1 minim	75
SPONGIA	2 minims	1 15
STICTA	1 minim	75
THUJA	1 minim	75
THUJA	2 minims	1 15
URTICA URENS	1 minim	1 15
VERATRUM ALB	1-2 minim, 1 minim	75
VERATRUM ALB	2 minims	1 15
VERATRUM VIR	1-2 minim, 1 minim	75
VERATRUM VIR	2 minims	1 15
VIBURNUM OPULUS	1 minim	75
VIBURNUM PRUNIFOL	1 minim	75

Please specify Fraser's Tablets

## TINCTURES.


	Per Pint.
TINCT. ACONITE.....	75
“ ACONITE (Fleming's).....	90
“ ARNICA.....	60
“ BELLADONNA.....	75
“ BENZOIN COMP.....	80
“ BRYONIA.....	70
“ CALENDULA.....	1 00
“ CALUMBA.....	60
“ CANNABIS INDICA.....	1 50
“ CANTHARIDES.....	75
“ CAPSICUM.....	50
“ CARDAMOM COMP.....	70
“ CIMICIFUGA.....	70
“ CINCHONA COMP.....	75
“ CUBEBS.....	60
“ DIGITALIS.....	75
“ GELSEMIUM.....	65
“ GENTIAN COMP.....	60
“ GINGER.....	65
“ GREEN SOAP.....	65
“ GREEN SOAP COMP.....	85
“ HYDRASTIS.....	75
“ HYOSCYAMUS.....	50
“ IGNATIA.....	85
“ IODINE.....	1 00
“ IODINE DECOLORIZED.....	1 00
“ IODINE COMP.....	1 25
“ IODINE (Churchill's).....	1 50
“ IPECAC AND OPIUM.....	1 50
“ IRON CHLORIDE.....	75
“ MATRICARIA.....	75
“ NUX VOMICA.....	75
“ OPIUM, 10 per cent. of Opium.....	1 50
“ OPIUM CAMPHORATED (Paregoric.).....	50
“ PHOSPHORUS.....	1 00
“ PULSATILLA.....	75
“ SANGUINARIA.....	60
“ STROPHANTHUS (Fraser).....	1 00
SPTS. AMMON. AROMAT.....	75
“ CAMPHOR.....	75
“ ETHER COMP. Hoffman's Anodyne).....	1 00
“ LAVENDER COMP.....	75
“ NITRE.....	75

Please specify Fraser's Tablets

FRASER'S  
Soluble Hypodermic Tablets.

The medicines in these tablets are combined with an unobjectionable base. They do not cause abscesses, and may be given internally as well as administered hypodermically.

These tablets are easily soluble in ten minims of warm water\*; are hard enough to retain their form, and will not pulverize or crumble in the bottle. They also possess the advantage of accuracy of dose and perfect preservation of the drug.



	Per Hundred.
ACONITINE NITRATE.....	1-50 gr..... 75
ACONITINE NITRATE.....	1-100 gr..... 60
ACONITINE NITRATE.....	1-150 gr..... 50
ACONITINE NITRATE.....	1-200 gr..... 40
APOMORPHINE MUR.....	1-10 gr..... 75
APOMORPHINE MUR.....	1-20 gr..... 55
ARSENITE OF POTASS.....	1-50 gr., 1-30 gr., 1-20 gr.. 30
ATROPINE SULPHATE.....	1-50 gr..... 35
ATROPINE SULPHATE.....	1-200 gr. 1-150 gr. 1-100 gr. 30
CAFFEINE CITRATE.....	1-2 gr..... 40
CAFFEINE & SODIUM BENZ.....	1-2 gr..... 40
CAFFEINE & SODIUM BENZ.....	1 gr..... 60
COCAINE HYDROCHLORATE.....	1-20 gr..... 35
COCAINE HYDROCHLORATE.....	1-10 gr..... 55
COCAINE HYDROCHLORATE.....	1-8 gr..... 65
COCAINE HYDROCHLORATE.....	1-5 gr..... 95
COCAINE HYDROCHLORATE.....	1-4 gr..... 1 10
CODEINE PHOSPHATE.....	1-4 gr..... 1 10
CODEINE SULPHATE.....	1-4 gr..... 75
CORROSIVE SUBLIMATE.....	1-50 gr., 1-30 gr., 1-20 gr.. 30
DIGITALIA.....	1-100 gr..... 40
DIGITALIA.....	1-50 gr..... 60
DIGITALIA.....	1-25 gr..... 1 00
DIGITALIA & STRYCHNINE.	{ Digitalia..... 1-100 gr. } { Strvch. Sulph..... 1-60 gr. } 60

\*It is unnecessary to use warm water for the Morphine, Atropine or Hyoscyamia Tablets ; but better to use hot water for the others.

DIGITALIA & STRYCHNINE COMP.....	{ Digitalia.....1-100 gr. Strych. Nitrate...1-60 gr. Nitroglycerin...1-100 gr. }	65
DIGITALIA & STRYCHNINE COMP. No. 2.....	{ Digitalia.....1-100 gr. Strych. Nitrate...1-50 gr. Nitroglycerin...1-100 gr. Atropine Sulph. 1-150 gr. }	75
ERGOTIN.....	1-10 gr.....	30
HUBBARD'S BROMIDES No. 1	{ Morphine Brom...1-4 gr. Hyosc. Hydrobr..1-100 gr. }	1 25
HUBBARD'S BROMIDES No. 2	{ Hyosc. Hydrobr..1-100 gr. Morphine Brom...1-8 gr. }	1 00
HYOSCINE COMP..... (Dr. Cutter.)	{ Hyosc. Hydrobr. 1-200 gr. Atropine Sulph. 1-200 gr. Morphine Sulph...1-4 gr. }	95
HYOSCINE HYDROBROM.....	1-200 gr.....	75
HYOSCINE HYDROBROM.....	1-100 gr.....	95
HYOSCYAMIA SULPHATE..... (White Cryst.)	1-200 gr.....	50
HYOSCYAMIA SULPHATE..... (White Cryst.)	1-100 gr.....	75
HYOSCYAMIA SULPHATE..... (White Cryst.)	1-50 gr.....	1 50
HYOSCYAMIA SULPHATE..... (White Cryst.)	1-25 gr.....	3 00
MORPHINE SULPHATE.....	1-2 gr.....	75
MORPHINE SULPHATE.....	1-3 gr.....	55
MORPHINE SULPHATE.....	1-4 gr.....	45
MORPHINE SULPHATE.....	1-10 gr., 1-8 gr., 1-6 gr....	35
MORPHINE & ATROPINE....	{ Morphine Sulph...1-2 gr. Atropine Sulph...1-120 gr. }	75
MORPHINE & ATROPINE....	{ Morphine Sulph...1-3 gr. Atropine Sulph...1-150 gr. }	55
MORPHINE & ATROPINE....	{ Morphine Sulph...1-4 gr. Atropine Sulph...1-120 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph...1-4 gr. Atropine Sulph...1-150 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph...1-4 gr. Atropine Sulph...1-180 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph...1-4 gr. Atropine Sulph...1-200 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph...1-6 gr. Atropine Sulph...1-150 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph...1-6 gr. Atropine Sulph...1-180 gr. }	45
MORPHINE & ATROPINE....	{ Morphine Sulph...1-8 gr. Atropine Sulph...1-200 gr. }	40
MORPHINE & ATROPINE....	{ Morphine Sulph...1-8 gr. Atropine Sulph...1-150 gr. }	40

When ordering or prescribing

MORPHINE PHTHALATE*.....	1-8 gr.....	75
MORPHINE PHTHALATE.....	1-4 gr.....	1 00
MORPHINE PHTHALATE & ATROPINE.....	{ Morph. Phthalate. 1-4 gr. } { Atropine Sulph. 1-150 gr. }	1 10
MORPHINE PHTHALATE & ATROPINE.....	{ Morph. Phthalate. 1-6 gr. } { Atropine Sulph. 1-180 gr. }	1 00
MORPHINE PHTHALATE & ATROPINE.....	{ Morph. Phthalate. 1-8 gr. } { Atropine Sulph. 1-200 gr. }	90
MORPHINE & HYOSCYAMIA	{ Morphine Sulph. 1-4 gr. } { Hyoscy. Sulph. 1-100 gr. }	1 10
NUCLEIN SOLUTION.....	1 min.....	1 75
(Dr. Aulde.)		
NITROGLYCERIN.....	1-200 gr. 1-100 gr. 1-50 gr.	30
NITROGLYCERIN & STRYCHNINE.....	{ Nitroglycerin... 1-100 gr. } { Strych. Nitrate... 1-50 gr. }	40
NITROGLYCERIN & STRYCHNINE COMP.....	{ Nitroglycerin... 1-100 gr. } { Strych. Nitrate... 1-60 gr. } { Atropine Sulph. 1-100 gr. }	50
PILOCARPINE MUR.....	1-10 gr.....	75
PILOCARPINE MUR.....	1-20 gr.....	50
POTASS. PERMANG.....	1 gr.....	30
QUININE BISULPHATE.....	1-2 gr.....	30
QUININE BI-MUR. & UREA.....	2 1-2 grs.....	75
(For Hypodermic use.)		
SPARTEINE.....	1-10 gr.....	35
SPARTEINE.....	1-4 gr.....	50
STRYCHNINE NITRATE.....	{ 1-200 gr. 1-100 gr. 1-60 gr. } { 1-50 gr. 1-40 gr. 1-30gr. } { 1-25 gr. 1-20 gr..... }	30
STRYCHNINE SULPHATE....	{ 1-200 gr. 1-100 gr. 1-60 gr. } { 1-50 gr. 1-40 gr. 1-30 gr. } { 1-25 gr. 1-20 gr..... }	30
LOCAL ANÆSTHETIC, No. 1 (Strong.)	{ Cocaine Hydrochl. 1-5 gr. } { Morph. Hydrochl. 1-40 gr. } { Sodium Chloride.. 1-5 gr. }	1 25
LOCAL ANÆSTHETIC, No. 2 (Normal.)	{ Cocaine Hydrochl. 1-10 gr. } { Morph. Hydrochl. 1-40 gr. } { Sodium Chloride .. 1-5 gr. }	80
LOCAL ANÆSTHETIC, No. 3 (Weak.)	{ Cocaine Hydrochl. 1-100gr. } { Morph. Hydrochl. 1-40gr. } { Sodium Chloride .. 1-5gr. }	40

\*Phthalate of Morphine is more soluble than the sulphate, and has been specially recommended for hypodermic use, causing less nausea and general Morphine disturbance.

## SYRINGE FOR HYPODERMIC TABLETS.



This syringe is made by THE FRASER TABLET TRITURATE MFG. CO., for convenience in dissolving and using their Hypodermic Tablets. It is provided with a movable end or cap to protect the packing. For use the cap is removed and the tablet dropped into the syringe, together with ten minims of water and shaken for a few moments, until the tablet is dissolved.

Alligator Case.....	Net \$2 50
Aluminum " .....	" 2 75
Morocco " .....	" 2 50
Nickel " .....	" 2 50
Wood " .....	" 2 00

For.....

Take..... Tablets

.....

.....

Dr. ....

### DIRECTION LABELS.

For Dispensing Bottles or Boxes,  
printed like illustration and gummed.

Price per 1000 ..... 50

Please specify Fraser's Tablets

# MISCELLANEOUS.

The attention of physicians is respectfully directed to the list of standard preparations from our own laboratory, which we present with confidence as containing in each case a first-class specimen of the drug it represents faithfully and skillfully compounded, and of uniform strength.

## PILLS.

BLAUD'S, Silver Coated .....	3 gr.....	per 1000	\$2 50
" " " .....	3 gr.....	per 100	35
" " " .....	5 gr.....	per 1000	3 00
" " " .....	5 gr.....	per 100	45
EXT. CANNABIS INDICA..	1-4 gr. . . .	per 1000	5 00
" " " . . . .	1-4 gr. . . .	per 100	75

Perfectly soluble in the stomach.

These pills are made from the Extract of Cannabis, which has given us such a reputation for quality and uniformity of the drug that we ship preparations of it all over the world. Every sample of the Extract is tested clinically before we make a preparation of it.

## SPECIAL PREPARATIONS.

FRASER'S CHAMOMILLA COMP. (8 oz.) Per Doz. \$10 50

(The retail price of this preparation is \$1.25 per bottle, containing 64 adult doses.)

CHAMOMILLA COMP. is a diffusible nerve tonic, from the formula of Dr. Thos. A. McBride, and is a combination of the mother tinctures of Cinchona, Chamomilla, Ignatia and Phosphorus with Aromatics.

Samples of Chamomilla Comp. and literature referring to its use will be forwarded upon application.

## FRASER'S GRANULES.

	Per Doz.	Per Gross.
PEPSIN AND SODA MINT GRANULES,	\$1 05	\$12 50
RHUBARB AND SODA MINT GRANULES,	1 05	12 50
SODA MINT GRANULES, - - -	90	10 50
CAMPBOR GLOBULES, - - -	1 05	12 50

THESE PREPARATIONS ARE PUT UP IN SCREW-CAP BOTTLES OF PATENTED DESIGN, and are packed in DOZENS for COUNTER DISPLAY with a neat advertising card with each dozen. They are superior to any similar preparations ever offered.

Please specify Fraser's Tablets

## POWDER PAPERS.

POWDER PAPERS, No. 1, $2\frac{1}{2} \times 3\frac{1}{4}$	Per 1000.....	30
" " No. 2, $2\frac{3}{4} \times 3\frac{7}{8}$	" " .....	45
" " No. 3, $3\frac{5}{8} \times 5\frac{1}{8}$	" " .....	65

## FRASER'S PIPETTES.



Fig. 2.

PLAIN, 4 inch.....	Each.....	Fig. 2	10
" 7 inch.....	" .....	"	10
" 12 inch.....	" .....	"	10
GRADUATED, 4 inch, tested....	20 minims.....		10
" 7 " " ....	30 " .....		25
" 12 " " ....	40 " .....		25



Fig. 3.

GRADUATED, 4 inch, with Rubber suction, 20 min. Fig. 3,	20
" 7 " " " " 30 min. "	25
" 12 " " " " 40 min. "	30

## DR. KEYES' ACID APPLICATOR.



Price.....	10
------------	----


## URINE TESTS.

These tests are sold in two-ounce glass stoppered bottles, and their reliability is guaranteed.

ACETIC ACID.....	25
CARBOL-FUCHSIN SOLUTION, For Stain.....	25
FEHLING'S TEST, Alkaline Solution.....	25
FEHLING'S TEST, Copper Solution.....	25
INDIGO-CARMINE, Acidulated Solution.....	25
INDIGO-CARMINE, Sodium Solution.....	25
MILLARD'S TEST .....	25
METHYLENE BLUE SOLUTION, For Stain.....	25
SULPHURIC ACID, For Stain.....	25
TANRET'S TEST.....	25

When ordering or prescribing

## BOTTLES FOR DISPENSING TABLETS.

No. 50.  
Full Size.No. 25.  
Full Size.No. 10.  
Full Size.No. 50 A.  
Full Size.No. 50 B.  
Full Size.

THE NUMBERS OF THESE BOTTLES CORRESPOND TO THE NUMBER OF TABLETS EACH BOTTLE HOLDS. FOR INSTANCE, NO. 50 HOLDS ABOUT 50 TABLETS.

		Per Gross.
No. 50,	without corks.....	\$1 10
" 50,	with "	1 50
" 25,	without "	1 00
" 25,	with "	1 40
" 10,	without "	90
" 10,	with "	1 25
" 50A,	without "	1 10
" 50A,	with "	1 50
" 25A,	without "	1 00
" 25A,	with "	1 40
" 10A,	without "	90
" 10A,	with "	1 25
" 50B,	without "	1 10
" 50B,	with "	1 50
" 25B,	without "	1 00
" 25B,	with "	1 40
" 10B,	without "	90
" 10B,	with "	1 25

Please specify Fraser's Tablets



**PAPER DISPENSING VIALS.****No. 50,  
Full Size.****No. 25,  
Full Size.****No. 15,  
Full Size.****No. 10,  
Full Size.**

THE NUMBERS OF THESE VIALS CORRESPOND TO THE NUMBER OF TABLETS EACH VIAL HOLDS. THEY ARE PACKED IN PAPER, HALF GROSS RED AND HALF GROSS WHITE IN EACH PACKAGE.

<b>No. 50.</b>	.....	<b>Per Gross, \$1 00</b>
" 25.	.....	" " 75
" 15.	.....	" " 70
" 10.	.....	" " 60

**STOCK CABINETS.**

For Glass Stoppered Bottles of 1000 tablets.




**No. 1.—Office Cabinet, to hold 25 Bottles. .... \$2 00**

Please specify Fraser's Tablets



No. 5.—Cabinet, to hold 50 Bottles ..... \$8 00



No. 3.—Wall Cabinet, to hold 108 Bottles..... \$7 00  
 No. 4. " " " " 192 " ..... 10 00

## Fraser's Cases.

---

We add to our Catalogue, for the first time, a price list of Medicine Cases.

Our experience in having goods of this kind made by various houses has taught us the need of a manufacturer of first-class medicine containers for both physicians and druggists, and we mean to make quality as much of a specialty in this line as we always have in medicine.

We intend to pay particular attention to special cases for physicians, who either have new ideas of their own, or who wish the cases on the list modified.

We also make sample cases, gun cases, rod cases, reel cases, fishing tackle cases, cartridge cases, mathematical instrument cases, surgical instrument cases, and all kinds of strap work.

Our friends will find us prompt and reliable in our work.

THE FRASER TABLET TRITURATE  
· MANUFACTURING CO

## FRASER'S BOTTLE CASES.



## FOR LIQUIDS.

A neat and compact leather case for holding bottles used in traveling. Made of handsome smooth-grain leather.

No. 1	½-ounce size, with flint-glass bottle.....	\$ 50
No. 2	1-ounce " " " " .....	60
No. 3	2-ounce " " " " .....	70
No. 4	3-ounce " " " " .....	80
No. 5	4-ounce " " " " .....	90
No. 6	6-ounce " " " " .....	1 00
No. 7	8-ounce " " " " .....	1 10
No. 8	12-ounce " " " " .....	1 25
No. 9	16-ounce " " " " .....	1 50

---

Special Cases made to order promptly.

## FRASER'S BOTTLE CASES.



FOR TABLETS, POWDERS, OR SALTS.

A neat and compact leather case for holding bottles used in traveling. Made of handsome smooth-grain leather.

No. 50	1 $\frac{1}{2}$ -ounce size, with screw-cap, wide-mouth bottle	\$	50
No. 51	1-ounce " " " " "		60
No. 52	2-ounce " " " " "		70
No. 53	3-ounce " " " " "		80
No. 54	4-ounce " " " " "		90
No. 55	6-ounce " " " " "		1 00
No. 56	8-ounce " " " " "		1 10
No. 57	12-ounce " " " " "		1 25
No. 58	16-ounce " " " " "		1 50

---

Interior Fittings can be rearranged at forty-eight hours' notice

## FRASER'S SEAMLESS CASE, No. 100.



No. 100 Case Closed.



No. 100 Case Open.

Seamless Black Leather Hand Case,  $9\frac{1}{2}$  inches long,  $3\frac{1}{2}$  inches wide, and  $2\frac{1}{4}$  inches thick, with patent rounded corners, strong leather handle, and durable spring lock. Contains sixteen  $\frac{1}{2}$ -ounce French square bottles, with cork-stop or screw-cap, as ordered. Each bottle will hold about 100 Tablet Triturates.

Price empty.....\$5 00

Special Cases made to order promptly.

## FRASER'S SEAMLESS CASE, No. 101.



Same general description as No. 100,  $9\frac{1}{2}$  inches long, 7 inches wide, and  $2\frac{1}{4}$  inches thick. Contains thirty-two  $\frac{1}{2}$ -ounce bottles.  
 Price.....\$7 50

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S SEAMLESS CASE, No. 102.



Seamless Hand or Buggy Case, 9 $\frac{3}{4}$  inches long, 6 $\frac{3}{4}$  inches deep, and 2 inches thick. Made of durable, smooth, black leather, has all rounded edges, two round sliding handles, and good substantial lock. Contains twenty-four 2 $\frac{1}{2}$ -drachm, seven 6-drachm, and five 3-drachm, cork-stoppered vials, and compartment for hypodermic syringe.


Price.....\$7 50

Special Cases made to order promptly.

## FRASER'S SEAMLESS CASE, No. 103.



Dr. Hunter's Case, No. 103, Closed.



Dr. Hunter's Case, No. 103, Open.

Seamless Hand or Buggy Case,  $9\frac{1}{2}$  inches long,  $6\frac{1}{2}$  inches deep, and  $3\frac{1}{2}$  inches thick. Made of durable, smooth, black leather, has all rounded edges, with handle and spring lock. Contains sixty cork-stoppered vials in six rows, each vial holding about fifty Tablet Triturates; also a space for instruments and sundries.


Price .....\$10 00

Interior Fittings can be rearranged at forty-eight hours' notice.

---

**FRASER'S SEAMLESS CASE, No. 104.**


No. 104 Case Closed.



No. 104 Case Open.

Seamless Hand or Buggy Case, 9 $\frac{1}{4}$  inches long, 6 inches high, and 5 inches thick. Made of durable, smooth, black leather, has all rounded edges, with handle and spring lock. Contains twenty 3-drachm, twenty 1-ounce, and three 2-ounce cork-stoppered vials; also a box for sundries, size of which is 4 $\frac{1}{2}$  inches long, 4 $\frac{1}{4}$  inches wide, and 1 $\frac{1}{4}$  inches deep.

Price.....\$6 00


---

Special Cases made to order promptly.

## FRASER'S SEAMLESS CASE, No. 105.



No. 105 Case Closed.



No. 105 Case Open.

Seamless Hand or Buggy Case,  $9\frac{1}{2}$  inches long,  $6\frac{1}{2}$  inches deep, and 5 inches thick. Made of durable, smooth, black leather, has all rounded edges, with handle and spring lock. Contains sixty-four  $\frac{1}{2}$ -ounce square bottles, either cork-stoppered or screw-cap, as ordered, in two separate compartments of four rows of eight bottles each, holding 100 tablets.

Price .....\$12 50

Interior Fittings can be rearranged at forty-eight hours' notice.

**FRASER'S SEAMLESS CASE, No. 106.****No. 106 Case Open.**

Seamless Hand or Buggy Case,  $11\frac{1}{2}$  inches long,  $5\frac{3}{4}$  inches high, and  $4\frac{3}{4}$  inches thick. Made of durable, smooth, black leather, has all rounded edges, re-enforced where necessary, with nickel-plated metallic protectors, with handle and spring lock. Contains nine 1-ounce, forty-five  $\frac{1}{2}$ -ounce cork-stoppered vials, with loops for instruments and space for sundries.

Price.....\$9 00

**FRASER'S SEAMLESS CASE, No. 107.**

Seamless Hand or Buggy Case (same size and general description as No. 106). Contains thirty  $\frac{1}{2}$ -ounce, and sixty 2-drachm cork-stoppered vials.

Price.....\$9 00

**FRASER'S SEAMLESS CASE, No. 108.**

Seamless Hand or Buggy Case (same size and general description as No. 106). Contains ten 1-ounce, twenty-eight 6-drachm, and four 2-ounce cork-stoppered vials; also box for sundries, 5 inches long,  $4\frac{1}{2}$  inches wide, and  $1\frac{1}{4}$  inches deep.


Price.....\$7 50

Special Cases made to order promptly.

## FRASER'S BUGGY CASE, No. 150.



No. 150 Case Closed.




No. 150 Case Open.

Black seal-grain leather, on a strong and durable frame, 9 $\frac{1}{4}$  inches long, 4 inches high, and 3 inches wide. Contains fifty-seven 2-drachm cork-stoppered vials in four rows, each of which will hold about fifty Tablet Triturates; also a syringe case containing a hypodermic syringe, two needles, and four vials, filled with hypodermic tablets.

Price with syringe complete, and empty bottles.....\$7 50

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S STIMSON CASE, No. 200.



Emergency Case, suitable for traveling or family use,  $6\frac{1}{2}$  inches long,  $2\frac{3}{4}$  inches high, and  $1\frac{3}{8}$  inches thick, with button clasp. Made of handsome morocco-grained leather. Contains six  $\frac{1}{2}$ -ounce Fraser's French square screw-cap bottles, arranged in a safely packed but very compact manner. The front of the case drops forward, exposing all the labels. Lined throughout with leather.

Price.....\$1 80

No. 201.—Style same as No. 200, 9 inches long,  $2\frac{3}{4}$  inches high, and  $1\frac{3}{8}$  inches thick. Contains eight  $\frac{1}{2}$ -ounce bottles.

Price.....\$2 00

No. 202.—Style same as No. 200, 11 inches long,  $2\frac{3}{4}$  inches high, and  $1\frac{3}{8}$  inches thick. Contains ten  $\frac{1}{2}$ -ounce bottles.

Price.....\$2 15

No. 203.—Style same as No. 200, 13 inches long,  $2\frac{3}{4}$  inches high, and  $1\frac{3}{8}$  inches thick. Contains twelve  $\frac{1}{2}$ -ounce bottles.

Price.....\$2 35

---

Special Cases made to order promptly.

**FRASER'S STIMSON CASE, No. 204.**

Double Case. No. 204 Case Closed.



Double Case. No. 204 Case Open.

**FRASER'S STIMSON CASE, No. 204.**

Emergency Case, suitable for traveling or family use,  $6\frac{1}{2}$  inches long,  $2\frac{3}{4}$  inches high, and  $2\frac{1}{2}$  inches thick, with button clasp. Made of handsome morocco-grained leather. Contains twelve  $\frac{1}{2}$ -ounce Fraser's French square screw-cap bottles in two rows, arranged in a safely packed but very compact manner. The front of the case drops forward, exposing all the labels. Lined throughout with leather.

Price.....\$2 35

---

Interior Fittings can be rearranged at forty-eight hours' notice.

**FRASER'S STIMSON CASE, No. 205.**

Emergency Case (same general description as No. 204), 9 inches long,  $2\frac{3}{4}$  inches high, and  $2\frac{1}{2}$  inches thick. Contains sixteen  $\frac{1}{2}$ -ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$2 60

**FRASER'S STIMSON CASE, No. 206.**

Emergency Case (same general description as No. 204), 11 inches long,  $2\frac{3}{4}$  inches high, and  $2\frac{1}{2}$  inches thick. Contains twenty  $\frac{1}{2}$ -ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$2 85

**FRASER'S STIMSON CASE, No. 207.**

Emergency Case (same general description as No. 204), 13 inches long,  $2\frac{3}{4}$  inches high, and  $2\frac{1}{2}$  inches thick. Contains twenty-four  $\frac{1}{2}$ -ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$3 00

**FRASER'S STIMSON CASE, No. 208.**

Emergency Case (same general description as No. 200),  $7\frac{1}{4}$  inches long,  $3\frac{1}{2}$  inches high, and 2 inches thick. Contains five 1-ounce Fraser's French square screw-cap bottles.

Price.....\$1 80

**FRASER'S STIMSON CASE, No. 209.**

Emergency Case (same general description as No. 200),  $8\frac{1}{2}$  inches long,  $3\frac{1}{2}$  inches high, and 2 inches thick. Contains six 1-ounce Fraser's French square screw-cap bottles.

Price.....\$2 00

**FRASER'S STIMSON CASE, No. 210.**

Emergency Case (same general description as No. 200), 11 inches long,  $3\frac{1}{2}$  inches high, and 2 inches thick. Contains eight 1-ounce Fraser's French square screw-cap bottles.

Price.....\$2 25

**FRASER'S STIMSON CASE, No. 211.**

Emergency Case (same general description as No. 204),  $7\frac{1}{4}$  inches long,  $3\frac{1}{2}$  inches high, and  $3\frac{1}{4}$  inches thick. Contains ten 1-ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$2 60

**FRASER'S STIMSON CASE, No. 212.**

Emergency Case (same general description as No. 204),  $8\frac{1}{2}$  inches long,  $3\frac{1}{2}$  inches high, and  $3\frac{1}{4}$  inches thick. Contains twelve 1-ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$3 00

**FRASER'S STIMSON CASE, No. 213.**

Emergency Case (same general description as No. 204), 11 inches long,  $3\frac{1}{2}$  inches high, and  $3\frac{1}{4}$  inches thick. Contains sixteen 1-ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$3 40

---

Special Cases made to order promptly.

**FRASER'S STIMSON CASE, No. 214.**

Emergency Case (same general description as No. 200), 7 inches long, 4 inches high, and  $2\frac{1}{4}$  inches thick. Contains four 2-ounce Fraser's French square screw-cap bottles.

Price.....\$2 00

**FRASER'S STIMSON CASE, No. 215.**

Emergency Case (same general description as No. 200),  $8\frac{1}{2}$  inches long, 4 inches high, and  $2\frac{1}{4}$  inches thick. Contains five 2-ounce Fraser's French square screw-cap bottles.

Price.....\$2 25

**FRASER'S STIMSON CASE, No. 216.**

Emergency Case (same general description as No. 200), 10 inches long, 4 inches high, and  $2\frac{1}{4}$  inches thick. Contains six 2-ounce Fraser's French square screw-cap bottles.

Price.....\$2 50

**FRASER'S STIMSON CASE, No. 217.**

Emergency Case (same general description as No. 204), 7 inches long, 4 inches high, and  $3\frac{3}{4}$  inches thick. Contains eight 2-ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$2 95

**FRASER'S STIMSON CASE, No. 218.**

Emergency Case (same general description as No. 204),  $8\frac{1}{2}$  inches long, 4 inches high, and  $3\frac{3}{4}$  inches thick. Contains ten 2-ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$3 25

**FRASER'S STIMSON CASE, No. 219.**

Emergency Case (same general description as No. 204), 10 inches long, 4 inches high, and  $3\frac{3}{4}$  inches thick. Contains twelve 2-ounce Fraser's French square screw-cap bottles in two rows.

Price.....\$3 60

**FRASER'S STIMSON CASE, No. 220.**

Emergency Case (same general description as No. 200),  $4\frac{1}{4}$  inches long,  $4\frac{3}{4}$  inches high, and  $2\frac{1}{2}$  inches thick. Contains two 3-ounce Fraser's French square screw-cap bottles.

Price.....\$1 75

**FRASER'S STIMSON CASE, No. 221.**

Emergency Case (same general description as No. 200),  $6\frac{1}{2}$  inches long,  $4\frac{3}{4}$  inches high, and  $2\frac{1}{2}$  inches thick. Contains three 3-ounce Fraser's French square screw-cap bottles.

Price.....\$2 25

**FRASER'S STIMSON CASE, No. 222.**

Emergency Case (same general description as No. 200), 8 inches long,  $4\frac{3}{4}$  inches high, and  $2\frac{1}{2}$  inches thick. Contains four 3-ounce Fraser's French square screw-cap bottles.

Price.....\$2 65

---

Interior Fittings can be rearranged at forty-eight hours' notice.

**FRASER'S STIMSON CASE, No. 223.**

Emergency Case (same general description as No. 200), 4 $\frac{1}{2}$  inches long, 5 $\frac{1}{4}$  inches high, and 2 $\frac{5}{8}$  inches thick. Contains two 4-ounce Fraser's French square screw-cap bottles.

Price.....\$2 00

**FRASER'S STIMSON CASE, No. 224.**

Emergency Case (same general description as No. 200), 6 $\frac{1}{4}$  inches long, 5 $\frac{1}{4}$  inches high, and 2 $\frac{5}{8}$  inches thick. Contains three 4-ounce Fraser's French square screw-cap bottles.

Price.....\$2 30

**FRASER'S STIMSON CASE, No. 225.**

Emergency Case (same general description as No. 200), 8 $\frac{1}{2}$  inches long, 5 $\frac{1}{4}$  inches high, and 2 $\frac{5}{8}$  inches thick. Contains four 4-ounce Fraser's French square screw-cap bottles.

Price.....\$2 75

**FRASER'S STIMSON CASE, No. 226.**

Emergency Case (same general description as No. 200), 5 inches long, 6 inches high, and 2 $\frac{3}{4}$  inches thick. Contains two 6-ounce Fraser's French square screw-cap bottles.

Price.....\$2 30

**FRASER'S STIMSON CASE, No. 227.**

Emergency Case (same general description as No. 200), 7 inches long, 6 inches high, and 2 $\frac{3}{4}$  inches thick. Contains three 6-ounce Fraser's French square screw-cap bottles.

Price.....\$2 70

**FRASER'S STIMSON CASE, No. 228.**

Emergency Case (same general description as No. 200), 9 inches long, 6 inches high, and 2 $\frac{3}{4}$  inches thick. Contains four 6-ounce Fraser's French square screw-cap bottles.

Price.....\$3 00

**FRASER'S STIMSON CASE, No. 229.**

Emergency Case (same general description as No. 200), 5 $\frac{1}{2}$  inches long, 6 inches high, and 3 inches thick. Contains two 8-ounce Fraser's French square screw-cap bottles.

Price.....\$2 50

**FRASER'S STIMSON CASE, No. 230.**

Emergency Case (same general description as No. 200), 7 $\frac{3}{4}$  inches long, 6 inches high, and 3 inches thick. Contains three 8-ounce Fraser's French square screw-cap bottles.

Price.....\$3 00

**FRASER'S STIMSON CASE, No. 231.**


Emergency Case (same general description as No. 200), 10 inches long, 6 inches high, and 3 inches thick. Contains four 8-ounce Fraser's French square screw-cap bottles.

Price.....\$3 50

---

Special Cases made to order promptly.

## FRASER'S POCKET VIAL CASE, No. 300.



No. 300 Case Closed.



No. 300 Case Open.

Folding Hypodermic Case for vest pocket,  $3\frac{1}{4}$  inches long,  $1\frac{3}{4}$  inches wide, and  $\frac{1}{2}$  inch thick. Soft, flexible, case, made of black or dark brown morocco leather, with spring-button catch. Contains six tube vials, each holding twenty-five Hypodermic Tablets


Price empty.....\$1 00

Price of vials used in case.....per doz..15c. Gross.. 1 50

---

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S POCKET VIAL CASE, No. 301.



No. 301 Case Closed.



No. 301 Case Open.

Folding Vest Pocket Case,  $3\frac{3}{8}$  inches long,  $2\frac{1}{2}$  inches wide, and  $\frac{9}{16}$  inches thick. Soft, flexible case, made of black or dark brown morocco leather. Contains six tube vials, each holding eighteen Tablet Triturates.

Price empty.....\$1 25


Price of vials used in case.....per doz. 15c. Gross 1 50

Special Cases made to order promptly.

## FRASER'S POCKET VIAL CASE, No. 302.



No. 302 Case Closed.



No. 302 Case Open.

Folding Pocket Emergency Case,  $4\frac{3}{4}$  inches long and  $3\frac{1}{2}$  inches wide, when closed. Soft, flexible case, made of black or dark brown morocco leather. Contains one row of twelve tube vials, each holding eighteen Tablet Triturates.

Price empty.....\$1 50  
 Filled with tablets, as per following list..... 2 00

Aconite Tr., 1 min.  
 Aloin and Belladonna Comp.  
 Ammonium Muriate Comp.  
 Belladonna Tincture, 1 min.  
 Bronchitis.  
 Calomel, 1-10 gr.

Calomel, 2 grs.  
 Digitalis, Tr., 2 mins.  
 Dover's Tablets,  $2\frac{1}{2}$  grs.  
 Morphine Sulphate,  $\frac{1}{4}$  gr.  
 Tartar Emetic, 1-50 gr.  
 Turpeth Mineral, 2 grs


Price of vials.....per doz..15c. Gross..\$1 50

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S POCKET VIAL CASE, No. 303.



No. 303 Case Closed.



No. 303 Case Open.

Folding Pocket Vial Case,  $6\frac{1}{2}$  inches long by  $3\frac{1}{2}$  inches wide, when closed. Soft, flexible case, made of black or dark brown morocco leather. Contains one row of ten tube vials, each holding forty Tablet Triturates.


Price empty.....\$2 50

Price of vials.....per doz..15c. Gross.. 1 50


---

Special Cases made to order promptly.

## FRASER'S POCKET VIAL CASE, No. 304.



No. 304 Case Closed.



No. 304 Case Open.


Folding Pocket Case, with clasp, 7¾ inches long by 3½ inches wide, when closed. Soft, flexible case, made of black or dark brown morocco leather. Contains one row of twenty tube vials, each holding eighteen Tablet Triturates.

Price empty.....\$2 50


Price of vials used in case.....per doz..15c. Gross..1 50

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S POCKET VIAL CASE, No. 305.



No. 305 Case Closed.




No. 305 Case Open.

Folding Pocket Case, with clasp,  $7\frac{3}{4}$  inches long by  $3\frac{1}{2}$  inches wide, when closed. Soft, flexible case, made of black or dark brown morocco leather. Contains two rows of twenty tube vials, each holding eighteen Tablet Triturates.

Price empty.....\$3 00  
 Price of vials used in case.....per doz..15c. Gross.. 1 50

Special Cases made to order promptly.

## FRASER'S POCKET VIAL CASE, No. 306.



No. 306 Case Closed.



No. 306 Case Open.


Folding Pocket Case, with clasp,  $7\frac{3}{4}$  inches long by  $3\frac{1}{2}$  inches wide, when closed. Soft, flexible case, made of black or dark brown morocco leather, containing one row of twelve tube vials, each holding forty Tablet Triturates.

Price empty.....\$2 50

Price of vials used in case.....per doz..15c. Gross.. 1 50

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S POCKET CASE, No. 307.



No. 307 Case Closed.



No. 307 Case Open.

Folding Pocket Case, with clasp,  $7\frac{1}{2}$  inches long by  $3\frac{1}{2}$  inches wide. Soft, flexible case, made of black or dark brown morocco leather, containing two rows of twelve tube vials, each vial holding forty Tablet Triturates.

Price empty.....\$3 00

Price of vials used in case.....per doz..15c. Gross.. 1 50

---

Special Cases made to order promptly.

## FRASER'S FOLDING POCKET CASE, No. 308.



No. 308 Case Closed.



No. 308 Case Open.


Folding Pocket Case with clasp,  $7\frac{1}{2}$  inches long by  $3\frac{1}{2}$  inches wide. Soft, flexible case, made of black or dark brown morocco leather, containing one row of twelve tube vials, each holding forty Tablet Triturates, and one row of twenty tube vials, each holding eighteen Tablet Triturates.

Price empty.....\$3 00


Price of vials used in case.....per doz..15c. Gross.. 1 50

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S VIAL CASE, No. 309.



No. 309 Case Closed.



No. 309 Case Open.

Folding Pocket Case with clasp, 6 $\frac{1}{4}$  inches long by 3 $\frac{1}{3}$  inches wide. Soft, flexible case, made of black or dark brown morocco leather, containing one row of twelve bottles, each holding eighteen Tablet Triturates, and four bottles, each holding twelve to fifteen Hypodermic Tablets.

Price with bottles only.....\$1 75

Price with Hypodermic Syringe..... 3 00

---

Special Cases made to order promptly.

**FRASER'S MOULDED VIAL CASE, No. 310.**

Folding Pocket Case, 10 inches long,  $3\frac{1}{4}$  inches wide, and  $1\frac{1}{2}$  inches thick, lined with velvet.

Our No. 310 is an ideal pocket vial case. The leather is moulded into proper shape, making it perfectly smooth on the outside, and very handsome in appearance. It is the most durable pocket case ever placed on the market. Contains twenty-four 2-drachm screw-cap vials.

Price empty.....\$3 00  
Price of vials used in case.....per doz..30c. Gross.. 3 00

**FRASER'S MOULDED VIAL CASE, No. 311.**

Folding Pocket Case (same general description as No. 310),  $8\frac{1}{2}$  inches long,  $3\frac{1}{4}$  inches wide, and  $1\frac{1}{2}$  inches thick. Contains twenty 2-drachm screw-cap vials.

Price empty.....\$2 75  
Price of vials used in case.....per doz..30c. Gross.. 3 00

**FRASER'S MOULDED VIAL CASE, No. 312.**

Folding Pocket Case (same general description as No. 310), 7 inches long,  $3\frac{1}{4}$  inches wide, and  $1\frac{1}{2}$  inches thick. Contains sixteen 2-drachm screw-cap vials.

Price empty.....\$2 50  
Price of vials used in case.....per doz..30c. Gross.. 3 00

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S BUGGY CASE, No. 400.



No. 400 Case Closed.



No. 400 Case Open.

Buggy Case,  $10\frac{1}{4}$  inches long,  $4\frac{1}{2}$  inches wide, and  $4\frac{1}{2}$  inches high. Best oil-tanned black sole leather, manufactured expressly for us. Handle made of same material as case, pressed smooth, so that it will not wear out glove, and firmly riveted to the case. Bottles held in place by nickel-plated metallic springs, and so arranged that every label can be seen at a glance when case is in use. Contains twenty 5-drachm, and thirty  $2\frac{1}{2}$ -drachm screw-capped vials; also a box for Hypodermic Syringe.

Price with Fraser's screw-cap jars for tablets.....\$5 00

Price of  $2\frac{1}{2}$ -drachm jars, used in case, per doz. 30c. Gross.. 3 00

Price of 5-drachm jars, used in case, per doz. 35c. Gross.. 4 20

Special Cases made to order promptly.

## FRASER'S BUGGY CASE, No. 401.



Buggy Case, 11 inches long, 6 inches wide, and 6 inches high. Best oil tanned black sole leather, manufactured expressly for us. Handle made of same material as case, pressed smooth, so that it will not wear out glove, and firmly riveted to the case. Bottles held in place by nickel-plated metallic springs, and so arranged that every label can be seen at a glance when case is in use. Contains seven 1½-ounce glass-stoppered bottles, twenty-four 6-drachm, and thirty-two 4-drachm screw-capped vials. Price with Fraser's screw-cap jars for tablets,.....\$6 50

Interior Fittings can be rearranged at forty-eight hours' notice

## FRASER'S BUGGY CASE, No. 402.



No. 402 Case Open.

Buggy Case, 11 inches long, 7 inches high, and 6 inches wide. Best oil-tanned black sole leather, manufactured expressly for us. Handle made of same material as case, pressed smooth, so that it will not wear out glove, and firmly riveted to the case. Bottles held in place by nickel-plated metallic springs, and so arranged that every label can be seen at a glance when case is in use. Contains seven 1½-ounce glass-stoppered bottles, twenty-four 6-drachm, thirty-two 4-drachm, and fifteen 2½-drachm screw-capped vials; also a box for sundries in top of the case, 9¼ inches long, 4 inches wide, and 1 inch deep.


Price with Fraser's screw-cap jars for tablets.....	\$7 50
Price of 6-drachm jars, used in case..per doz..45c.	Gross.. 5 00
Price of 4-drachm jars, used in case..per doz..35c.	Gross.. 3 75
Price of 2½-drachm jars, used in case..per doz..30c.	Gross.. 3 00
Price of 1½-oz. glass-stoppered bottles..each..10c.	Per doz. 1 00

Special Cases made to order promptly.

## FRASER'S BUGGY CASE No. 403.



No. 403 Case Closed.



No. 403 Case Open.

Buggy Case,  $9\frac{1}{2}$  inches long,  $4\frac{1}{2}$  inches high, and  $3\frac{1}{2}$  inches wide. Best oil-tanned black sole leather, manufactured expressly for us. Contains twenty-four 4-drachm and thirty  $2\frac{1}{2}$ -drachm Fraser's specially made screw-cap jars, so arranged that every label can be seen at a glance when case is in use.

Price complete.....\$6 50


Price of 4-drachm jars used in case..per doz..40c. Gross.. 4 50

Price of  $2\frac{1}{2}$ -drachm jars used in case..per doz..35c. Gross.. 3 75

---

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S CASE, No. 404.



Trunk or box-shaped case, 11 inches long, 6 inches wide, and 6½ inches high. Best oil-tanned black sole leather, manufactured expressly for us. Has a nickel-plated lock with key.

The inside is divided into three spaces. One space contains eight 1½-ounce vials and twenty-four 3-drachm vials; the other side contains twenty-four 1-ounce vials; the intermediate space contains a lacquered tray for instruments or sundries. This tray is 10½ inches long, 6 inches high, and 2 inches wide.

Price complete.....\$8 00

## FRASER'S CASE, No. 405.




Buggy and Portable Case,  $8\frac{1}{2}$  inches long,  $5\frac{1}{4}$  inches high, and  $4\frac{1}{2}$  inches wide. Sole leather interchangeable leaf case made of solid leather, and so arranged that the trays or leaves holding the bottles, etc., can be easily removed, and other leaves with different size bottles, or variety of medicine, can be put in their places. Extra leaves, with vials, supplied with this case.

The following leaves, with bottles, go with each case :


- |  |  |
|--|--|
| One leaf with ten 1-ounce vials.               | One leaf with fourteen 3-drachm vials. |
| One leaf with twelve 4-drachm vials.           | One leaf with eleven 6-drachm vials.   |
| One leaf with eight 6-drachm vials.            | One box for hypodermic syringe.        |
| One leaf with box for instruments and sundries |  |

Price, with six leaves complete, .....\$10 00

## FRASER'S BUGGY CASE, No. 406.



No. 406 Case Closed.




No. 406 Case Open.

Buggy Case,  $9\frac{1}{2}$  inches long,  $4\frac{1}{2}$  inches high, and  $3\frac{3}{4}$  inches wide. Best oil-tanned black sole leather, manufactured expressly for us. Contains ten 6-drachm, eight 4-drachm, and thirty  $2\frac{1}{2}$ -drachm Fraser's specially made screw-cap jars, so arranged that every label can be seen at a glance when case is in use; also a box for hypodermic syringe and pocket for sundries.


Price complete.....	\$5 50
Price of 6-drachm jars used in case..per doz..	.35c. Gross.. 4 20
Price of 4-drachm jars used in case..per doz..	.30c. Gross.. 3 00
Price of $2\frac{1}{2}$ -drachm jars used in case..per doz..	.30c. Gross.. 3 00

Special Cases made to order promptly.

## FRASER'S MEDICINE CASE, No. 407.



No. 407 Case Closed.



No. 407 Case Open.


Medicine Case, 7 inches long,  $4\frac{1}{2}$  inches high, and  $1\frac{1}{2}$  inches wide. Made of best oil-tanned black sole leather, manufactured expressly for us. Contains six 1-ounce cork-stoppered vials held in place by metallic clamps.

Price.....	\$2 00
Price of 1-ounce bottles used in case... ..per doz..	35


---

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S BICYCLE CASE, No. 408.



No. 408 Case Closed.




No. 408 Case Open.

Bicycle Case, 9 inches long, 5½ inches high, and 2½ inches wide. Best oil-tanned black sole leather, manufactured expressly for us, all hand-stitched. Contains twenty-four 3-drachm and sixteen 2-drachm cork-stoppered vials.

Price of 408 Case.....	\$5 00
Price of 3-drachm vials used in case..per doz..	.25c. Gross.. 2 00
Price of 2-drachm vials used in case..per doz..	.20c. Gross.. 1 75

---

Special Cases made to order promptly.

**FRASER'S BUGGY CASE, No. 409.**

Buggy Case,  $10\frac{1}{2}$  inches long, 6 inches wide, and  $4\frac{1}{2}$  inches high. Solid russet sole leather, moulded into shape, and made in such a manner that the edges of the leather on the ends extend beyond the surface of the case, preventing any wearing of the stitches, and adding materially to the handsome appearance of the case. Contains ten 10-drachm, twelve 6-drachm, and twenty-eight 4-drachm cork-stoppered vials, and loops for instruments under bottles.

Price.....\$10 00

**FRASER'S BUGGY CASE, No. 410.**

Buggy Case (same general description as No. 409.) Contains ten 10-drachm, twenty-four 6-drachm, and twenty-eight 2-drachm cork-stoppered vials.


Price.....\$10 00

Interior Fittings can be rearranged at forty-eight hours' notice.

## FRASER'S OBSTETRICAL BAG, No. 500.



No. 500 Bag Closed.



No. 500 Bag Open.

## FRASER'S OBSTETRICAL BAG, No. 500.

Made of strong, black seal-grain leather, lined throughout with leather or chamois. Suitable for carrying obstetrical and other long instruments. Contains six 2-ounce glass-stoppered bottles arranged in loops on one side.


15-inch bag, without bottles.....	\$3 50
16 " " " " " " .....	3 75
17 " " " " " " .....	4 00
18 " " " " " " .....	4 25
15 " " with bottles.....	4 00
16 " " " " " " .....	4 25
17 " " " " " " .....	4 50
18 " " " " " " .....	4 75

Special Cases made to order promptly.

## FRASER'S COMBINATION BAG, No. 501.



No. 501 Bag Closed.



No. 501 Bag Open.

Satchel Buggy Case, made of black seal-grain leather, 15 inches long, 6 inches wide, and 5 inches deep, lined throughout with leather. Contains eleven 1-ounce, nineteen 3-drachm, and twenty-four 2-drachm vials on one side, and a space on the other side  $14\frac{1}{2}$  inches long,  $4\frac{1}{2}$  inches wide, and  $2\frac{1}{2}$  inches deep, for instruments or appliances.

The ounce and 3-drachm bottles are arranged on a leaf which may be entirely removed from the bag.

Price.....\$11 00

---

Interior Fittings can be rearranged at forty-eight hours' notice.

**FRASER'S MEDICINE BAG, No. 502.****No. 502 Bag Open.****FRASER'S MEDICINE BAG, No. 502.**

Buggy Case (cabin bag style), 10½ inches long, 7 inches high, and 6 inches wide. Made of strong, black seal-grain leather. Contains thirty-six 4-drachm vials, and a pocket; also an empty space in the centre 10 inches long, 6 inches high, and 2 inches wide.

The bottles are arranged on leaves which may be entirely removed from the bag.

Price .....\$10 00

**FRASER'S MEDICINE BAG, No. 503.**

Same general description as No. 502, excepting that it contains twenty-four 4-drachm and twenty-four 2-drachm bottles.

Price .....\$10 00

**FRASER'S MEDICINE BAG, No. 504.**

Same general description as No. 502, excepting that it contains nine 1-ounce and twenty-four 4-drachm bottles.

Price .....\$10 00

**FRASER'S MEDICINE BAG, No. 505.**


Same general description as No. 502, excepting that it contains forty-eight 4-drachm bottles, with no pocket.

Price .....\$10 00

---


Special Cases made to order promptly.

## FRASER'S TRAVELING BAG, STYLE No. 506.



Made of black sea-grain leather, 10 inches long,  $6\frac{1}{2}$  inches high,  $5\frac{1}{2}$  inches wide, and lined throughout with leather. Contains seven rows, each containing eight  $\frac{1}{2}$ -ounce bottles, and a compartment for instruments 8 inches long, 2 inches wide, and 1 inch high; also pockets for powder papers, etc.

Price .....\$12 50

**FRASER'S EMERGENCY BAG, No. 507.**

Fraser's Cabin Style Emergency Bag, with nickel-plated trimmings, 16 inches long, 8 inches high, and 7½ inches wide. Made of strong, black seal-grain leather, and lined throughout with leather. Contains ten 2-ounce glass-stoppered bottles, a pocket, and loops for instruments; also a sterilizing tray, held by straps to the bottom of the bag.


Price .....\$9 00

**FRASER'S EMERGENCY BAG, No. 508.**

Same description as No. 507, excepting that the dimensions are 17 inches long, 11 inches high, and 8½ inches wide, and has brass trimmings.

Price .....\$10 00

Special Cases made to order promptly.

**FRASER'S MEDICINE CHEST, No. 600.****No. 600 Medicine Chest Closed.**

Leather Medicine Chest, 12 $\frac{3}{4}$  inches long, 8 $\frac{1}{4}$  inches wide, and 9 $\frac{1}{4}$  inches high, with substantial lock. Made of heavy russet leather, without seams or stitching on the edges, leaving the full strength of the leather where it is most needed. A stout leather strap is attached to the chest in such a manner that it may be used as a handle or shoulder strap, as required. Contains two 4-ounce, sixteen 2-ounce, and nineteen 1-ounce glass-stoppered bottles; also one 1-ounce graduated measure, one 1-ounce glass mortar and pestle, and four  $\frac{1}{2}$ -ounce metallic-covered ointment jars.

Tray for scales and space for appliances, 11 inches long, 3 $\frac{1}{4}$  inches wide, and 3 inches high, under bottles.

Price .....\$16 00

**FRASER'S MEDICINE CHEST, No. 601.**

Leather Medicine Chest, with substantial lock (same general description as No. 600), 15 $\frac{1}{2}$  inches long, 9 $\frac{1}{4}$  inches wide, and 9 $\frac{1}{4}$  inches high. Contains four 4-ounce, eighteen 2-ounce, twenty 1-ounce, and eight  $\frac{1}{2}$ -ounce glass-stoppered bottles; also one 1-ounce graduated measure, one 1-ounce glass mortar and pestle, and four  $\frac{1}{2}$ -ounce metallic-covered ointment jars.

Tray for scales and space for appliances, 14 $\frac{1}{2}$  inches long, 4 $\frac{1}{4}$  inches wide, and 3 inches high, under bottles.

Price .....\$19 50

**FRASER'S MEDICINE CHEST, No. 602.**

Leather Medicine Chest, with substantial lock (same general description as number 600), 14 $\frac{1}{2}$  inches long, 9 $\frac{1}{4}$  inches wide, and 9 $\frac{1}{4}$  inches high. Contains four 4-ounce, sixteen 2-ounce, eighteen 1-ounce, and six  $\frac{1}{2}$ -ounce glass-stoppered bottles; also one 1-ounce graduated measure, one 1-ounce glass mortar and pestle, and four  $\frac{1}{2}$ -ounce metallic-covered ointment jars.

Tray for scales and space for appliances, 13 $\frac{1}{2}$  inches long, 4 $\frac{1}{4}$  inches wide, and 3 inches high, under bottles.

Price .....\$18 00



No. 600 Medicine Chest Open.

**FRASER'S MEDICINE CHEST, No. 603.**

Leather Medicine Chest, with substantial lock (same general description as No. 600), 11 inches long,  $8\frac{1}{4}$  inches wide, and  $9\frac{1}{4}$  inches high. Contains two 4-ounce, fourteen 2-ounce, and sixteen 1-ounce glass-stoppered bottles; also one 1-ounce graduated measure, one 1-ounce glass mortar and pestle, and four  $\frac{1}{2}$ -ounce metallic-covered ointment jars.

Tray for scales and space for appliances,  $9\frac{1}{2}$  inches long,  $3\frac{1}{4}$  inches wide, and 3 inches high, under bottles.

Price .....\$14 50

**FRASER'S MEDICINE CHEST, No. 604.**

Leather Medicine Chest, with substantial lock (same general description as No. 600),  $9\frac{3}{4}$  inches long,  $8\frac{1}{4}$  inches wide, and  $9\frac{1}{4}$  inches high. Contains two 4-ounce, twelve 2-ounce, and thirteen 1-ounce glass-stoppered bottles; also one 1-ounce graduated measure, one 1-ounce glass mortar and pestle, and four  $\frac{1}{2}$ -ounce metallic-covered ointment jars.

Tray for scales and space for appliances,  $8\frac{1}{4}$  inches long,  $3\frac{1}{4}$  inches wide, and 3 inches high, under bottles.

Price .....\$13 50

Special Cases made to order promptly.

**FRASER'S MEDICINE CHEST, No. 605.**

Leather Medicine Chest, 13 inches long, 9 inches wide, and 5 $\frac{3}{4}$  inches high, with substantial lock, and with a tray in front of bottles, and a space under tray for appliances. Made of heavy russet leather, without seams or stitching on the edges, leaving the full strength of the leather where it is most needed. Contains four 4-ounce, nine 2-ounce, twelve 1-ounce, and six  $\frac{1}{2}$ -ounce glass-stoppered bottles.

Space under tray for appliances and sundries, 8 $\frac{1}{2}$  inches long 3 $\frac{1}{2}$  inches wide, and 1 $\frac{3}{4}$  inches high. Space in tray, 5 $\frac{1}{2}$  inches long, 3 $\frac{1}{4}$  inches wide, and 2 $\frac{1}{2}$  inches high.

Price .....\$13 00

**FRASER'S MEDICINE CHEST, No. 606.**

Leather Medicine Chest, with substantial lock (same general description as No. 605), 11 inches long, 7 $\frac{3}{4}$  inches wide, and 5 $\frac{3}{4}$  inches high. Contains five 4-ounce, seven 2-ounce, and six 1 ounce glass-stoppered bottles.

Space under tray for appliances and sundries, 7 inches long, 3 $\frac{1}{2}$  inches wide, and 2 inches high. Space in tray, 6 $\frac{3}{4}$  inches long, 3 inches wide, and 2 $\frac{1}{4}$  inches high.


Price .....\$10 00

**FRASER'S MEDICINE CHEST, No. 607.**

Leather Medicine Chest, with substantial lock (same general description as No. 605), 9 inches long, 6 $\frac{1}{2}$  inches wide, and 5 $\frac{3}{4}$  inches high. Contains ten 2-ounce and four 1-ounce glass-stoppered bottles.

Space under tray for appliances and sundries, 4 $\frac{1}{2}$  inches long, 2 $\frac{1}{2}$  inches wide, and 2 inches high. Space in tray, 4 $\frac{1}{2}$  inches long, 2 inches wide, and 2 inches high.

Price .....\$8 50

**FRASER'S MEDICINE CHEST, No. 608.**

No. 608 Medicine Chest Open.

**FRASER'S MEDICINE CHEST, No. 608.**

Leather Medicine Chest, 7 inches long, 5 $\frac{3}{4}$  inches wide, and 5 inches high. Made of heavy russet leather, without seams or stitching on the edges, leaving the full strength of the leather where it is most needed. Contains two 4-ounce and six 2-ounce glass-stoppered bottles, and space for sundries.

Price.....\$5 25

**FRASER'S MEDICINE CHEST, No. 609.**

Leather Medicine Chest (same general style as No. 608), 7 inches long, 4 $\frac{3}{4}$  inches wide, and 4 $\frac{3}{4}$  inches high. Made of heavy russet leather, without seams or stitching on the edges, leaving the full strength of leather where it is most needed. Contains nine 1-ounce glass-stoppered bottles, and space for sundries.

Price.....\$4 75


**FRASER'S MEDICINE CHEST, No. 610.**

Leather Medicine Chest (same general style as No. 608), 7 inches long, 4 $\frac{3}{4}$  inches wide, and 4 $\frac{3}{4}$  inches high. Made of heavy russet leather, without seams or stitching on the edges, leaving the full strength of the leather where it is most needed. Contains fifteen 1-ounce glass-stoppered bottles.

Price.....\$5 25

---

Special Cases made to order promptly.

**FRASER'S MEDICINE CASE, No. 611.**

No. 611 Medicine Case Closed.



No. 611 Medicine Case Open.

**FRASER'S MEDICINE CASE, No. 611.**

Leather Medicine Case with strap, flat shape,  $8\frac{1}{2}$  inches long,  $5\frac{3}{4}$  inches wide, and  $2\frac{1}{4}$  inches high. Made of heavy russet leather, without seams or stitching on the edges, leaving the full strength of the leather where it is most needed. Contains six 2-ounce glass-stoppered bottles.

Price.....\$4 75

**FRASER'S MEDICINE CASE, No. 612.**

Leather Medicine Case (same general style as No. 611),  $7\frac{1}{2}$  inches long, 5 inches wide, and  $2\frac{1}{4}$  inches high. Made of heavy russet leather, without seams or stitching on the edges, leaving the full strength of the leather where it is most needed. Contains six 1-ounce glass-stoppered bottles.

Price.....\$4 25

---

Interior Fittings can be rearranged at forty-eight hours' notice.

**FRASER'S MEDICINE CASE, No. 651.**

No. 651 Case Closed.



No. 651 Case Open.

**FRASER'S MEDICINE CASE, No. 651.**

Seamless Leather Case, made of black or russet leather,  $8\frac{1}{4}$  inches long,  $4\frac{3}{4}$  inches wide, and  $2\frac{1}{2}$  inches high. Flat pattern, round corners and ends; has a strap and substantial lock. Contains six 1-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price .....\$5 35

---

Special Cases made to order promptly.

**FRASER'S MEDICINE CASE, No. 652.**

Style same as No. 651, 10 $\frac{3}{4}$  inches long, 4 $\frac{3}{4}$  inches wide, and 2 $\frac{1}{2}$  inches high. Contains eight 1-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$5 75

**FRASER'S MEDICINE CASE, No. 653.**

Style same as No. 651, 7 $\frac{1}{4}$  inches long, 5 inches wide, and 2 $\frac{3}{4}$  inches high. Contains four 2-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$5 00

**FRASER'S MEDICINE CASE, No. 654.**

Style same as No. 651, 10 inches long, 5 inches wide, and 2 $\frac{3}{4}$  inches high. Contains six 2-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$5 75

**FRASER'S MEDICINE CASE, No. 655.**

Style same as No. 651, 13 inches long, 5 inches wide, and 2 $\frac{3}{4}$  inches high. Contains eight 2-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$6 25

**FRASER'S MEDICINE CASE, No. 656.**

Style same as No. 651, 6 $\frac{1}{4}$  inches long, 6 $\frac{1}{4}$  inches wide, and 2 $\frac{1}{2}$  inches high. Contains three 4-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$5 00

**FRASER'S MEDICINE CASE, No. 657.**

Style same as No. 651, 8 inches long, 6 $\frac{1}{4}$  inches wide, and 2 $\frac{1}{2}$  inches high. Contains four 4-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$5 75

**FRASER'S MEDICINE CASE, No. 658.**

Style same as No. 651, 12 inches long, 6 $\frac{1}{4}$  inches wide, and 2 $\frac{1}{2}$  inches high. Contains six 4-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$6 25

**FRASER'S MEDICINE CASE, No. 659.**

Style same as No. 651, 7 $\frac{1}{2}$  inches long, 5 $\frac{3}{4}$  inches wide, and 2 $\frac{1}{4}$  inches high. Contains two 8-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$5 75

**FRASER'S MEDICINE CASE, No. 660.**

Style same as No. 651, 8 $\frac{3}{4}$  inches long, 7 $\frac{1}{2}$  inches wide, and 2 $\frac{1}{4}$  inches high. Contains three 8-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$6 50

**FRASER'S MEDICINE CASE, No. 661.**

Style same as No. 651, 11 $\frac{1}{2}$  inches long, 7 $\frac{1}{2}$  inches wide, and 2 $\frac{1}{4}$  inches high. Contains four 8-ounce Fraser's square flint-glass bottles, with flat glass stopper.

Price.....\$7 00

---

Interior Fittings can be rearranged at forty-eight hours' notice.

**FRASER'S MEDICINE CASE, No. 663.****FRASER'S MEDICINE CASE, No. 662.**

Double Seamless Leather Case, made of black or russet leather, 7 inches long, 4 $\frac{1}{4}$  inches wide, and 3 $\frac{1}{2}$  inches high. Flat pattern, round corners and ends; has a strap and substantial lock. Contains ten 1-ounce Fraser's square flint-glass bottles, with flat glass stopper, in two rows, with pocket.

Price.....\$7 00

Special Cases made to order promptly.

**FRASER'S MEDICINE CASE, No. 663.**

Double Seamless Leather Case, made of black or russet leather, 8 $\frac{1}{4}$  inches long, 4 $\frac{3}{4}$  inches wide, and 3 $\frac{1}{2}$  inches high. Flat pattern, round corners and ends; has a strap and substantial lock. Contains twelve 1-ounce Fraser's square flint-glass bottles, with flat glass stopper, in two rows, with pocket.

Price.....\$7 50

**FRASER'S MEDICINE CASE, No. 664.**

Double Seamless Leather Case, made of black or russet leather, 10 $\frac{3}{4}$  inches long, 4 $\frac{3}{4}$  inches wide, and 3 $\frac{1}{2}$  inches high. Flat pattern, round corners and ends; has a strap and substantial lock. Contains sixteen 1-ounce Fraser's square flint-glass bottles, with flat glass stopper, in two rows, with pocket.

Price.....\$8 75

**FRASER'S MEDICINE CASE, No. 665.**

Double Seamless Leather Case, made of black or russet leather, 8 $\frac{1}{2}$  inches long, 5 $\frac{1}{4}$  inches wide, and 4 $\frac{1}{4}$  inches high. Flat pattern, round corners and ends; has a strap and substantial lock. Contains ten 2-ounce Fraser's square flint-glass bottles, with flat glass stopper, in two rows, with pocket.

Price.....\$8 25

**FRASER'S MEDICINE CASE, No. 666.**

Double Seamless Leather Case, made of black or russet leather, 10 inches long, 5 $\frac{1}{4}$  inches wide, and 4 $\frac{1}{4}$  inches high. Flat pattern, round corners and ends; has a strap and substantial lock. Contains twelve 2-ounce Fraser's square flint-glass bottles, with flat glass stopper, in two rows, with pocket.

Price.....\$8 75

**FRASER'S MEDICINE CASE, No. 667.**


Double Seamless Leather Case, made of black or russet leather, 13 inches long, 5 $\frac{1}{4}$  inches wide, and 4 $\frac{1}{4}$  inches high. Flat pattern, round corners and ends; has a strap and substantial lock. Contains sixteen 2-ounce Fraser's square flint-glass bottles, with flat glass stopper, in two rows, with pocket.

Price.....\$9 50

**INDEX.**

Chamomilla Comp.....	127
Dispensing Conveniences and Cabinets.....	129-132
Fraser's Granules .....	127
Hypodermic Syringes .....	126
Lozenges .....	116-117
Miscellaneous .....	127-128
Physicians' Cases .....	133-183
Pills.....	127
Tablets, Compressed.....	1-116
Tablets, Hypodermic .....	123-125
Tablets, Mother Tincture .....	119-121
Tablets, Tincture.....	118-119
Tablet Triturates.....	1-116
Tinctures .....	122
Urine Tests.....	128

Interior Fittings can be rearranged at forty-eight hours' notice.



THE name "FRASER'S TABLETS" is recognized by medical practitioners everywhere as the synonym for Tablet Triturates, which were first developed, manufactured, and brought to their present state of perfection by us upon the suggestion of DR. ROBERT M. FULLER. (See *Medical Record*, March 9th, 1878, and March 25th, 1882.)

This method of dispensing medicines that are given in small quantities has been attended, since its introduction by us, by the success due :

1st. To the care which we have taken in selecting and manufacturing the drugs and chemicals employed.

2d. To the thoroughness of the trituration and division of the remedy, and consequent perfect accuracy of the dose.

3d. To the quickness and ease with which our tablets are dissolved or diffused in water, and in the acids of the stomach.

4th. To their elegance of form and novelty of appearance.

---

## CAUTION.

---

The manufacture of Tablet Triturates is apparently so simple that many have attempted to imitate our goods. We desire, therefore, to warn practitioners that dangers are possible in the use of tablets which may resemble ours, but made without the painstaking care and skill, the high class drugs, the close adherence to the formula and above all the thorough trituration, which lead to the uniformity of product that has been a recognized characteristic of our goods from the first. In prescribing the small doses represented by many of the preparations in tablet triturate form, these characteristics are indispensable to the success of the remedies.

*Laboratories: 454 to 474 Eighteenth Street, Brooklyn, New York.*

